



Research article

Chronic treatment with escitalopram and venlafaxine affects the neuropeptide S pathway differently in adult Wistar rats exposed to maternal separation

Milosz Golyszny^{1,*}, Michał Zieliński¹, Monika Paul-Samojedny², Artur Palasz³ and Ewa Obuchowicz¹

¹ Department of Pharmacology, Faculty of Medical Sciences in Katowice, Medical University of Silesia, Medyków 18, 40-752 Katowice, Poland

² Department of Medical Genetics, Faculty of Pharmaceutical Sciences in Sosnowiec, Medical University of Silesia, Jedności 8, 41-200 Sosnowiec, Poland

³ Department of Histology, Faculty of Medical Sciences in Katowice, Medical University of Silesia, Medyków 18, 40-752 Katowice, Poland

* **Correspondence:** Email:miloszgolyszny@o2.pl; Tel: +48 32 2088522; Fax: +48 32 2523835.

Supplementary materials




Figure S1. The individual data points from Δ Ct values obtained in qPCR. (A) Plots for each groups in the NPS and NPSR mRNA level analyses in brainstem. (B) Δ Ct values from each groups in the NPSR mRNA expression analysis in the hypothalamus and amygdala.



AIMS Press

©2022 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)