



Research article

Modeling portfolio loss by interval distributions

Bill Huajian Yang^{1,*}, Jenny Yang² and Haoji Yang³

¹ Royal Bank of Canada, 155 Wellington St W, ON M5V 3H6 Toronto, Canada

² Scotiabank, 4 King St W, ON M5H 1A1 Toronto, Canada

³ Department of statistics, Shenzhen University, 518000 Shenzhen, China

* **Correspondence:** Email: h_y02@yahoo.ca.

Abstract: Models for a continuous risk outcome has a wide application in portfolio risk management and capital allocation. We introduce a family of interval distributions based on variable transformations. Densities for these distributions are provided. Models with a random effect, targeting a continuous risk outcome, can then be fitted by maximum likelihood approaches assuming an interval distribution. Given fixed effects, regression function can be estimated and derived accordingly when required. This provides an alternative regression tool to the fraction response model and Beta regression model.

Keywords: interval distribution; model with a random effect; tailed index; expected shortfall; heteroscedasticity; beta regression model; fraction response model; maximum likelihood

1. Introduction

For a continuous risk outcome $0 < y < 1$, a model with a random effect has potentially a wide application in portfolio risk management, especially, for stress testing [1,2,7,16,19], capital allocation, conditional expected shortfall estimation [3,11,17].

Given fixed effects $x = (x_1, x_2, \dots, x_k)$, two widely used regression models to estimate the expected value $E(y|x)$ are: the fraction response model [10] and Beta regression model [4,6,8]. There are cases, however, where tail behaviours or severity levels of the risk outcome are relevant. In those cases, a regression model may no longer fit in for the requirements. In addition, a fraction response model of the form $E(y|x) = \Phi(a_0 + a_1x_1 + \dots + a_kx_k)$ may not be adequate when data exhibits significant heteroscedasticity, where Φ is a map from R^1 to the open interval $(0, 1)$.

In this paper, we assume that the risk outcome y is driven by a model:

$$y = \Phi(a_0 + a_1x_1 + \dots + a_kx_k + bs), \quad (1.1)$$

where s is a random continuous variable following a known distribution, independent of fixed effects (x_1, x_2, \dots, x_k) . Parameters a_0, a_1, \dots, a_k are constant, while parameter b can be chosen to be

dependent on (x_1, x_2, \dots, x_k) when required, for example, for addressing data heteroscedasticity.

Given random effect model (1.1), the expected value $E(y|x)$ can be deduced accordingly. It is given by the integral $\int_{\Omega} \Phi(a_0 + a_1x_1 + \dots + a_kx_k + bs)f(s)ds$ over the domain Ω of s , where f is the probability density of s . Given the routine QUAD implemented in SAS and Python, this integral can be evaluated as quickly as other function calls. Relative error tolerance for QUAD is $1.49e-8$ in Python and is $1e-7$ in SAS. But one can rescale the default tolerance to a desired level when necessary. This leads to an alternative regression tool to the fraction response model and Beta regression model.

We introduce a family of interval distributions based on variable transformations. Probability densities for these distributions are provided (Proposition 2.1). Parameters of model (1.1) can then be estimated by maximum likelihood approaches assuming an interval distribution. In some cases, these parameters get an analytical solution without the needs for a model fitting (Proposition 4.1). We call a model with a random effect, where parameters are estimated by maximum likelihood assuming an interval distribution, an interval distribution model.

In its simplest form, the interval distribution model $y = \Phi(a + bs)$, where a and b , are constant, can be used to model the loss rate as a random distribution for a homogeneous portfolio. Let y_{α} and s_{α} denote the α -quantiles for y and s at level α , $0 < \alpha < 1$. Then $y_{\alpha} = \Phi(a + bs_{\alpha})$. The conditional expected shortfall for loss rate y , at level α , can then be estimated as the integral $\frac{1}{1-\alpha} \int_{[s_{\alpha}, +\infty)} \Phi(a + bs)f(s)ds$, where f is the density of s . Meanwhile, a stress testing loss estimate, derived from a model on a specific scenario, can be compared in loss rate to severity y_{α} ($= \Phi(a + bs_{\alpha})$), to position its level of severity. A loss estimate may not have reached the desired, for example, 99% level yet, if it is far below $y_{0.99}$, and far below the maximum historical loss rate. In which case, further recalibrations for the model may be required.

The paper is organized as follows: in section 2, we introduce a family of interval distributions. A measure for tail fatness is defined. In section 3, we show examples of interval distributions and investigate their tail behaviours. We propose in section 4 an algorithm for estimating the parameters in model (1.1).

2. Interval distributions generated by transformations

Interval distributions introduced in this section are defined for a risk outcome over a finite open interval (c_0, c_1) , where $c_0 < c_1$ are finite numbers. These interval distributions can potentially be used for modeling a risk outcome over an arbitrary finite interval, including interval $(0,1)$, by maximum likelihood approaches.

Let $D = (d_0, d_1)$, $d_0 < d_1$, be an open interval, where d_0 can be finite or $-\infty$ and d_1 can be finite or $+\infty$.

Let

$$\Phi: D \rightarrow (c_0, c_1) \quad (2.1)$$

be a transformation with continuous and positive derivatives $\Phi'(x) = \phi(x)$. A special example is $(c_0, c_1) = (0, 1)$, and $\Phi: D \rightarrow (0, 1)$ is the cumulative distribution function (CDF) of a random variable with a continuous and positive density.

Given a continuous random variable s , let f and F be respectively its density and CDF. For constants a and $b > 0$, let

$$y = \Phi(a + bs), \quad (2.2)$$

where we assume that the range of variable $(a + bs)$ is in the domain D of Φ . Let $g(y, a, b)$ and $G(y, a, b)$ denote respectively the density and CDF of y in (2.2).

Proposition 2.1. Given $\Phi^{-1}(y)$, functions $g(y, a, b)$ and $G(y, a, b)$ are given as:

$$g(y, a, b) = U_1/(bU_2) \quad (2.3)$$

$$G(y, a, b) = F\left[\frac{\Phi^{-1}(y)-a}{b}\right]. \quad (2.4)$$

where

$$U_1 = f\{[\Phi^{-1}(y) - a]/b\}, \quad U_2 = \phi[\Phi^{-1}(y)]. \quad (2.5)$$

Proof. A proof for the case when $(c_0, c_1) = (0, 1)$ can be found in [18]. The proof here is similar. Since $G(y, a, b)$ is the CDF of y , it follows:

$$\begin{aligned} G(y, a, b) &= P[\Phi(a + bs) \leq y] \\ &= P\{s \leq [\Phi^{-1}(y) - a]/b\} \\ &= F\{[\Phi^{-1}(y) - a]/b\}. \end{aligned}$$

By chain rule and the relationship $\Phi[\Phi^{-1}(y)] = y$, the derivative of $\Phi^{-1}(y)$ with respect to y is

$$\frac{\partial \Phi^{-1}(y)}{\partial y} = \frac{1}{\phi[\Phi^{-1}(y)]}. \quad (2.6)$$

Taking the derivative of $G(y, a, b)$ with respect to y , we have

$$\frac{\partial G(y, a, b)}{\partial y} = \frac{f\{[\Phi^{-1}(y)-a]/b\}}{b\phi[\Phi^{-1}(y)]} = \frac{U_1}{bU_2}.$$

One can explore into these interval distributions for their shapes, including skewness and modality. For stress testing purposes, we are more interested in tail risk behaviours for these distributions.

Recall that, for a variable X over $(-\infty, +\infty)$, we say that the distribution of X has a fat right tail if there is a positive exponent $\alpha > 0$, called tailed index, such that $P(X > x) \sim x^{-\alpha}$. The relation \sim refers to the asymptotic equivalence of functions, meaning that their ratio tends to a positive constant. Note that, when the density is a continuous function, it tends to 0 when $x \rightarrow +\infty$. Hence, by L'Hospital's rule, the existence of tailed index is equivalent to saying that the density decays like a power law, whenever the density is a continuous function.

For a risk outcome over a finite interval (c_0, c_1) , $c_0 < c_1$, however, its density can be $+\infty$ when approaching boundaries c_0 and c_1 . Let y_0 be the largest lower bound for all values of y under (2.2), and y_1 the smallest upper bound. We assume $y_0 = c_0$ and $y_1 = c_1$.

We say that an interval distribution has a fat right tail if the limit $\lim_{y \rightarrow y_1^-} g(y, a, b) = +\infty$, and a fat left tail if $\lim_{y \rightarrow y_0^+} g(y, a, b) = +\infty$, where $y \rightarrow y_0^+$ and $y \rightarrow y_1^-$ denote respectively y approaching y_0 from the right-hand-side, and y_1 from the left-hand-side. For simplicity, we write $y \rightarrow y_0$ for $y \rightarrow y_0^+$, and $y \rightarrow y_1$ for $y \rightarrow y_1^-$.

Given $\alpha > 0$, we say that an interval distribution has a fat right tail with tailed index α if $\lim_{y \rightarrow y_1} g(y, a, b)(y_1 - y)^\beta = +\infty$ whenever $0 < \beta < \alpha$, and $\lim_{y \rightarrow y_1} g(y, a, b)(y_1 - y)^\beta = 0$ for $\beta > \alpha$. Similarly, an interval distribution has a fat left tail with tailed index α if $\lim_{y \rightarrow y_0} g(y, a, b)(y - y_0)^\beta = +\infty$ whenever $0 < \beta < \alpha$, and $\lim_{y \rightarrow y_0} g(y, a, b)(y - y_0)^\beta = 0$ for $\beta > \alpha$. Here the status at $\beta = \alpha$ is left open. There are examples (Remark 3.4), where an interval distribution has a fat right tail with tailed index α , but the limit $\lim_{y \rightarrow y_1} g(y, a, b)(y_1 - y)^\alpha$ can

either be $+\infty$ or 0. Under this definition, a tailed index of an interval distribution with a continuous density is always larger than 0 and less or equal to 1, if it exists.

Recall that, for a Beta distribution with parameters $\alpha > 0$ and $\beta > 0$, its density is given by $f(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)}$, where $B(\alpha,\beta)$ is the Beta function. Under the above definition, Beta distribution has a fat right tail with tailed index $(1 - \beta)$ when $0 < \beta < 1$, and a fat left tail with tailed index $(1 - \alpha)$ when $0 < \alpha < 1$.

Next, because the derivative of Φ is assumed to be continuous and positive, it is strictly monotonic. Hence $\Phi^{-1}(y)$ is defined. Let

$$z = \Phi^{-1}(y) \quad (2.7)$$

Then $\lim_{y \rightarrow y_0} z$ exists (can be $-\infty$), and the same for $\lim_{y \rightarrow y_1} z$ (can be $+\infty$). Let $\lim_{y \rightarrow y_0} z = z_0$, and $\lim_{y \rightarrow y_1} z = z_1$. Rewrite $g(y, a, b)$ as $g(\Phi(z), a, b)$ by (2.7). Let $\partial[g(\Phi(z), a, b)]^{\frac{1}{\beta}}/\partial z$ denote the derivative of $[g(\Phi(z), a, b)]^{-1/\beta}$ with respect to z .

Lemma 2.2. Given $\beta > 0$, the following statements hold:

(i) $\lim_{y \rightarrow y_0} g(y, a, b)(y - y_0)^\beta = \lim_{z \rightarrow z_0} g(\Phi(z), a, b)(\Phi(z) - y_0)^\beta$ and $\lim_{y \rightarrow y_1} g(y, a, b)(y_1 - y)^\beta = \lim_{z \rightarrow z_1} g(\Phi(z), a, b)(y_1 - \Phi(z))^\beta$.

(ii) If $\lim_{y \rightarrow y_0} g(y, a, b) = +\infty$ and $\lim_{z \rightarrow z_0} \{\partial[g(\Phi(z), a, b)]^{\frac{1}{\beta}}/\partial z\}/\phi(z)$ is 0 (resp. $+\infty$), then $\lim_{y \rightarrow y_0} g(y, a, b)(y - y_0)^\beta = +\infty$ (resp. 0).

(iii) If $\lim_{y \rightarrow y_1} g(y, a, b) = +\infty$ and $\lim_{z \rightarrow z_1} -\{\partial[g(\Phi(z), a, b)]^{\frac{1}{\beta}}/\partial z\}/\phi(z)$ is 0 (resp. $+\infty$), then $\lim_{y \rightarrow y_1} g(y, a, b)(y_1 - y)^\beta = +\infty$ (resp. 0).

Proof. The first statement follows from the relationship $y = \Phi(z)$. For statements (ii) and (iii), we show only (iii). The proof for (ii) is similar. Notice that

$$[g(y, a, b)(y_1 - y)^\beta]^{-1/\beta} = \frac{[g(y, a, b)]^{-1/\beta}}{y_1 - y} = \frac{[g(\Phi(z), a, b)]^{-1/\beta}}{y_1 - \Phi(z)}. \quad (2.8)$$

By L'Hospital's rule and taking the derivatives of the numerator and the denominator of (2.8) with respect to z , we have $\lim_{y \rightarrow y_1} [g(y, a, b)(y_1 - y)^\beta]^{-1/\beta} = 0$ (resp. $+\infty$) if $\lim_{z \rightarrow z_0} -\{\partial[g(\Phi(z), a, b)]^{-1/\beta}/\partial z\}/\phi(z)$ is 0 (resp. $+\infty$). Hence $\lim_{y \rightarrow y_1} g(y, a, b)(y_1 - y)^\beta = +\infty$ (resp. 0).

For tail convexity, we say that the right tail of an interval distribution is convex if $g(y, a, b)$ is convex for $y_1 - \epsilon < y < y_1$ for sufficiently small $\epsilon > 0$. Similarly, the left tail is convex if $g(y, a, b)$ is convex for $y_0 < y < y_0 + \epsilon$ for sufficiently small $\epsilon > 0$. One sufficient condition for convexity for the right (resp. left) tail is $g''_{yy}(y, a, b) \geq 0$ when y is sufficiently close to y_1 (resp. y_0).

Again, write $g(y, a, b) = g(\Phi(z), a, b)$. Let

$$h(z, a, b) = \log[g(\Phi(z), a, b)], \quad (2.9)$$

where $\log(x)$ denotes the natural logarithmic function. Then

$$g(y, a, b) = \exp[h(z, a, b)]. \quad (2.10)$$

By (2.9), (2.10), using (2.6) and the relationship $z = \Phi^{-1}(y)$, we have

$$g'_y = [h'_z(z)/\phi(z)]\exp[h(\Phi^{-1}(y), a, b)],$$

$$g''_{yy} = \left[\frac{h''_{zz}(z)}{\phi^2(z)} - \frac{h'_z(z)\phi'_z(z)}{\phi^3(z)} + \frac{h'_z(z)h'_z(z)}{\phi^2(z)} \right] \exp[h(\Phi^{-1}(y), a, b)]. \quad (2.11)$$

The following lemma is useful for checking tail convexity, it follows from (2.11).

Lemma 2.3. Suppose $\phi(z) > 0$, and derivatives $h'_z(z)$, $h''_{zz}(z)$, and $\phi'_z(z)$, with respect to z , all exist. If $h''_{zz}(z) \geq 0$ and $h'_z(z)\phi'_z(z) \leq 0$, then $g''_{yy}(y, a, b) \geq 0$.

3. Examples of interval distributions and their tail behaviours

In this section, we focus on the case where $(c_0, c_1) = (0, 1)$, and $\Phi: D \rightarrow (0, 1)$ in (2.2) is the CDF of a continuous distribution. This includes, for example, the CDFs for standard normal and standard logistic distributions.

One can explore into a wide list of densities with different choices for Φ and s under (2.2). We consider here only the following four interval distributions:

A. $s \sim N(0,1)$ and Φ is the CDF for the standard normal distribution.

B. s follows the standard logistic distribution and Φ is the CDF for the standard normal distribution.

C. s follows the standard logistic distribution and Φ is its CDF.

D. $s \sim N(0,1)$ and Φ is the CDF for standard logistic distribution.

Densities for cases A, B, C, and D are given respectively in (3.3) (section 3.1), (A.1), (A.3), and (A5) (Appendix A). Tail behaviour study is summarized in Propositions 3.3, 3.5, and Remark 3.6. Sketches of density plots are provided in Appendix B for distributions A, B, and C.

3.1. Case A: the Vasicek distribution and its tail behaviours

Using the notations of section 2, we have $\phi = f$ and $\Phi = F$. We claim that $y = \Phi(a + bs)$ under (2.2) follows the Vasicek distribution [13,14].

By (2.5), we have

$$\log\left(\frac{U_1}{U_2}\right) = \frac{-z^2 + 2az - a^2 + b^2z^2}{2b^2} \quad (3.1)$$

$$= \frac{-(1-b^2)\left(z - \frac{a}{1-b^2}\right)^2 + \frac{b^2}{1-b^2}a^2}{2b^2}. \quad (3.2)$$

Therefore, we have

$$g(y, a, b) = \frac{1}{b} \exp\left\{ \frac{-(1-b^2)\left(z - \frac{a}{1-b^2}\right)^2 + \frac{b^2}{1-b^2}a^2}{2b^2} \right\}. \quad (3.3)$$

Again, using the notations of section 2, we have $y_0 = 0$ and $y_1 = 1$. With $z = \Phi^{-1}(y)$, we have $\lim_{y \rightarrow 0} z = -\infty$ and $\lim_{y \rightarrow 1} z = +\infty$. Recall that a variable $0 < y < 1$ follows a Vasicek distribution [13,14] if its density has the form:

$$g(y, p, \rho) = \sqrt{\frac{1-\rho}{\rho}} \exp\left\{ -\frac{1}{2\rho} \left[\sqrt{1-\rho} \Phi^{-1}(y) - \Phi^{-1}(p) \right]^2 + \frac{1}{2} [\Phi^{-1}(y)]^2 \right\}, \quad (3.4)$$

where p is the mean of y , and ρ is a parameter called asset correlation.

Proposition 3.1. Density (3.3) is equivalent to (3.4) under the relationships:

$$a = \frac{\Phi^{-1}(p)}{\sqrt{1-\rho}} \quad \text{and} \quad b = \sqrt{\frac{\rho}{1-\rho}}. \quad (3.5)$$

Proof. A similar proof can be found in [19]. By (3.4), we have

$$\begin{aligned} g(y, p, \rho) &= \sqrt{\frac{1-\rho}{\rho}} \exp\left\{-\frac{1-\rho}{2\rho} [\Phi^{-1}(y) - \Phi^{-1}(p)/\sqrt{1-\rho}]^2 + \frac{1}{2} [\Phi^{-1}(y)]^2\right\} \\ &= \frac{1}{b} \exp\left\{-\frac{1}{2} \left[\frac{\Phi^{-1}(y) - a}{b}\right]^2\right\} \exp\left\{\frac{1}{2} [\Phi^{-1}(y)]^2\right\} \\ &= U_1/(bU_2) = g(y, a, b). \end{aligned}$$

The following relationships are implied by (3.5):

$$\rho = \frac{b^2}{1+b^2}, \quad (3.6)$$

$$a = \Phi^{-1}(p)\sqrt{1+b^2}. \quad (3.7)$$

Remark 3.2. The mode of $g(y, p, \rho)$ in (3.4) is given in [14] as $\Phi\left(\frac{\sqrt{1-\rho}}{1-2\rho}\Phi^{-1}(p)\right)$. We claim this is the same as $\Phi\left(\frac{a}{1-b^2}\right)$. By (3.6), $1-2\rho = \frac{1-b^2}{1+b^2}$ and $\sqrt{1-\rho} = \frac{1}{\sqrt{1+b^2}}$. Therefore, we have

$$\frac{\sqrt{1-\rho}}{1-2\rho}\Phi^{-1}(p) = \frac{\sqrt{1+b^2}}{1-b^2}\Phi^{-1}(p) = \frac{a}{1-b^2}.$$

This means $\Phi\left(\frac{\sqrt{1-\rho}}{1-2\rho}\Phi^{-1}(p)\right) = \Phi\left(\frac{a}{1-b^2}\right)$.

Proposition 3.3. The following statements hold for $g(y, a, b)$ given in (3.3):

(i) $g(y, a, b)$ is unimodal if $0 < b < 1$ with mode given by $\Phi\left(\frac{a}{1-b^2}\right)$, and is in U-shape if $b > 1$.

(ii) If $b > 1$, then $g(y, a, b)$ has a fat left tail and a fat right tail with tailed index $(1 - 1/b^2)$.

(iii) If $b > 1$, both tails of $g(y, a, b)$ are convex, and is globally convex if in addition $a = 0$.

Proof. For statement (i), we have $-(1-b^2) < 0$ when $0 < b < 1$. Therefore by (3.2) function $\log\left(\frac{U_1}{U_2}\right)$ reaches its unique maximum at $z = \frac{a}{1-b^2}$, resulting in a value for the mode at $\Phi\left(\frac{a}{1-b^2}\right)$. If $b > 1$, then $-(1-b^2) > 0$, thus by (3.2), $g(y, a, b)$ is first decreasing and then increasing when y varying from 0 to 1. This means (y, a, b) is in U-shape.

Consider statement (ii). First by (3.3), if $b > 1$, then $\lim_{y \rightarrow 1} g(y, a, b) = +\infty$ and $\lim_{y \rightarrow 0} g(y, a, b) = +\infty$. Thus $g(y, a, b)$ has a fat right and a fat left tail. Next for tailed index, we use Lemma 2.2 (ii) and (iii). By (3.1),

$$[g(\Phi(z), a, b)]^{-1/\beta} = b^{1/\beta} \exp\left(-\frac{(b^2-1)z^2+2az-a^2}{2\beta b^2}\right) \quad (3.8)$$

By taking the derivative of (3.8) with respect to z and noting that $\phi(z) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{z^2}{2}\right)$, we have

$$-\left\{\partial[g(\Phi(z), a, b)]^{-\frac{1}{\beta}}/\partial z\right\}/\phi(z) = \sqrt{2\pi} b^{\frac{1}{\beta}} \frac{(b^2-1)z+a}{\beta b^2} \exp\left(-\frac{(b^2-1)z^2+2az-a^2}{2\beta b^2} + \frac{z^2}{2}\right). \quad (3.9)$$

Thus $\lim_{z \rightarrow +\infty} -\left\{\partial[g(\Phi(z), a, b)]^{-\frac{1}{\beta}}/\partial z\right\}/\phi(z)$ is 0 if $\frac{b^2-1}{\beta b^2} > 1$, and is $+\infty$ if $\frac{b^2-1}{\beta b^2} < 1$. Hence by Lemma 2.2 (iii), $g(y, a, b)$ has a fat right tail with tailed index $(1 - 1/b^2)$. Similarly, for the left tail, we have by (3.9)

$$\left\{ \partial [g(\Phi(z), a, b)]^{-\frac{1}{\beta}} / \partial z \right\} / \phi(z) = -\sqrt{2\pi} b^{\frac{1}{\beta}} \frac{(b^2-1)z+a}{\beta b^2} \exp\left(-\frac{(b^2-1)z^2+2az-a^2}{2\beta b^2} + \frac{z^2}{2}\right). \quad (3.10)$$

Thus $\lim_{z \rightarrow -\infty} \left\{ \partial [g(\Phi(z), a, b)]^{-\frac{1}{\beta}} / \partial z \right\} / \phi(z)$ is 0 if $\frac{b^2-1}{\beta b^2} > 1$, and is $+\infty$ if $\frac{b^2-1}{\beta b^2} < 1$.

Hence $g(y, a, b)$ has a fat left tail with tailed index $(1 - 1/b^2)$ by Lemma 2.2 (ii).

For statement (iii), we use Lemma 2.3. By (2.9) and using (3.2), we have

$$h(z, a, b) = \log\left(\frac{U_1}{bU_2}\right) = \frac{-(1-b^2)\left(z - \frac{a}{1-b^2}\right)^2 + \frac{b^2}{1-b^2}a^2}{2b^2} - \log(b).$$

When $b > 1$, it is not difficult to check out that $h''_{zz}(z) \geq 0$ and $h'_z(z)\phi'_z(z) \leq 0$ when $z \rightarrow \pm\infty$ or when $a = 0$.

Remark 3.4. Assume $\beta = (1 - 1/b^2)$ and $b > 1$. By (3.9), we see

$$\lim_{z \rightarrow +\infty} -\left\{ \partial [g(\Phi(z), a, b)]^{-\frac{1}{\beta}} / \partial z \right\} / \phi(z)$$

is $+\infty$ for $a = 0$, and is 0 for $a > 0$. Hence for this β , the limit $\lim_{y \rightarrow 1} g(y, a, b)(1 - y)^\beta$ can be either 0 or $+\infty$, depending on the value of a .

3.2. Tail behaviours for interval distributions for cases B-D

For these distributions, we again focus on their tail behaviours. A proof for the next proposition can be found in Appendix A.

Proposition 3.5. The following statements hold:

(a) Density $g(y, a, b)$ has a fat left tail and a fat right tail for case B for all $b > 0$, and for case C if $b > 1$. For case D, it does not have a fat right tail nor a fat left tail for any $b > 0$.

(b) The tailed index of $g(y, a, b)$ for both right and left tails is 1 for case B for all $b > 0$, and is $(1 - \frac{1}{b})$ for case C for B for $b > 1$.

Remark 3.6. Among distributions A, B, C, and Beta distribution, distribution B gets the highest tailed index of 1, independent of the choices of $b > 0$.

4. Algorithms for fitting interval distribution models

In this section, we assume that Φ in (2.2) is a function from R^1 to $(0, 1)$ with positive continuous derivatives. We focus on parameter estimation algorithms for model (1.1).

First, we consider a simple case, where risk outcome y is driven by a model:

$$y = \Phi(v + bs), \quad (4.1)$$

where $b > 0$ is a constant, $v = a_0 + a_1x_1 + \dots + a_kx_k$, and $s \sim N(0, 1)$, independent of fixed effects $x = (x_1, x_2, \dots, x_k)$. The function Φ does not have to be the standard normal CDF. But when Φ is the standard normal CDF, the expected value $E(y|x)$ can be evaluated by the formula $E_S[\Phi(a + bs)] = \Phi\left(\frac{a}{\sqrt{1+b^2}}\right)$ [12].

Given a sample $\{(x_{1i}, x_{2i}, \dots, x_{ki}, y_i)\}_{i=1}^n$, where $(x_{1i}, x_{2i}, \dots, x_{ki}, y_i)$ denotes the i^{th} data point of the sample, let $z_i = \Phi^{-1}(y_i)$. and $v_i = a_0 + a_1x_{1i} + \dots + a_kx_{ki}$. By (2.3), the log-likelihood function for model (4.1) is:

$$LL = \sum_{i=1}^n \left\{ \log f\left(\frac{z_i - v_i}{b}\right) - \log \phi(z_i) - \log b \right\}, \quad (4.2)$$

where f is the density of s . The part of $\sum_{i=1}^n \log \phi(z_i)$ is constant, which can be dropped off from the maximization.

Recall that the least squares estimators of a_0, a_1, \dots, a_k , as a row vector, that minimize the sum squares

$$SS = \sum_{i=1}^n (z_i - v_i)^2 \quad (4.3)$$

has a closed form solution given by the transpose of $(X^T X)^{-1} X^T Z$ [5,9] whenever the design matrix X has a rank of k , where

$$X = \begin{bmatrix} 1 & x_{11} & \dots & x_{k1} \\ 1 & x_{12} & \dots & x_{k2} \\ \dots & \dots & \dots & \dots \\ 1 & x_{1n} & \dots & x_{kn} \end{bmatrix}, \quad Z = \begin{bmatrix} z_1 \\ z_2 \\ \dots \\ z_n \end{bmatrix}.$$

The next proposition shows there exists an analytical solution for the parameters of model (4.1).

Proposition 4.1. Given a sample $\{(x_{1i}, x_{2i}, \dots, x_{ki}, y_i)\}_{i=1}^n$, assume that the design matrix has a rank of k . If $s \sim N(0, 1)$, then the maximum likelihood estimates of parameters (a_0, a_1, \dots, a_k) , as a row vector, and parameter b are respectively given by the transpose of $(X^T X)^{-1} X^T Z$, and $b^2 = \frac{1}{n} \sum_{i=1}^n (z_i - v_i)^2$. In absence of fixed effects $\{x_1, x_2, \dots, x_k\}$, parameters a_0 and b^2 degenerate respectively to the sample mean and variance of z_1, z_2, \dots, z_n .

Proof. Dropping off the constant term from (4.2) and noting $f(z) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{z^2}{2}\right)$, we have

$$LL = -\frac{1}{2b^2} \sum_{i=1}^n (z_i - v_i)^2 - n \log b, \quad (4.4)$$

Hence the maximum likelihood estimates (a_0, a_1, \dots, a_k) are the same as least squares estimators of (4.3), which are given by the transpose of $(X^T X)^{-1} X^T Z$. By taking the derivative of (4.4) with respect to b and setting it to zero, we have $b^2 = \frac{1}{n} \sum_{i=1}^n (z_i - v_i)^2$.

Next, we consider the general case of model (1.1), where the risk outcome y is driven by a model:

$$y = \Phi[v + ws], \quad (4.5)$$

where parameter w is formulated as $w = \exp(u)$, and $u = b_0 + b_1 x_1 + \dots + b_k x_k$. We focus on the following two cases:

- (a) $s \sim N(0, 1)$,
- (b) s is standard logistic.

Given a sample $\{(x_{1i}, x_{2i}, \dots, x_{ki}, y_i)\}_{i=1}^n$, let $w_i = \exp(b_0 + b_1 x_{1i} + \dots + b_k x_{ki})$ and $u_i = b_0 + b_1 x_{1i} + \dots + b_k x_{ki}$. The log-likelihood functions for model (4.5), dropping off the constant part $\log(U_2)$, for cases (a) and (b) are given respectively by (4.6) and (4.7):

$$LL = \sum_{i=1}^n -\frac{1}{2} [(z_i - v_i)^2 / w_i^2 - u_i], \quad (4.6)$$

$$LL = \sum_{i=1}^n \{- (z_i - v_i) / w_i - 2 \log[1 + \exp\{- (z_i - v_i) / w_i\}] - u_i\}, \quad (4.7)$$

Recall that a function is log-concave if its logarithm is concave. If a function is concave, a local maximum is a global maximum, and the function is unimodal. This property is useful for searching maximum likelihood estimates.

Proposition 4.2. The functions (4.6) and (4.7) are concave as a function of (a_0, a_1, \dots, a_k) . As a function of (b_0, b_1, \dots, b_k) , (4.6) is concave.

Proof. It is well-known that, if $f(x)$ is log-concave, then so is $f(Az + b)$, where $Az + b : R^m \rightarrow R^1$

is any affine transformation from the m -dimensional Euclidean space to the 1-dimensional Euclidean space. For (4.6), the function $f(x) = -(z - v)^2 \exp(-2u)$ is concave as a function of v , thus function (4.6) is concave as a function of (a_0, a_1, \dots, a_k) . Similarly, this function $f(x)$ is concave as a function of u , so (4.6) is concave as a function of (b_0, b_1, \dots, b_k) .

For (4.7), the linear part $-(z_i - v_i) \exp(-u_i)$, as a function of (a_0, a_1, \dots, a_k) , in (4.7) is ignored. For the second part in (4.7), we know $-\log\{1 + \exp[-(z - v)/\exp(u)]\}$, as a function of v , is the logarithm of the CDF of a logistic distribution. It is well-known that the CDF for a logistic distribution is log-concave. Thus (4.7) is concave with respect to (a_0, a_1, \dots, a_k) .

In general, parameters (a_0, a_1, \dots, a_k) and (b_0, b_1, \dots, b_k) in model (4.5) can be estimated by the algorithm below.

Algorithm 4.3. Follow the steps below to estimate parameters of model (4.5):

- (a) Given (b_0, b_1, \dots, b_k) , estimate (a_0, a_1, \dots, a_k) by maximizing the log-likelihood function;
- (b) Given (a_0, a_1, \dots, a_k) , estimate (b_0, b_1, \dots, b_k) by maximizing the log-likelihood function;
- (c) Iterate (a) and (b) until a convergence is reached.

5. Conclusions

With the interval distributions introduced in this paper, models with a random effect can be fitted for a continuous risk outcome by maximum likelihood approaches assuming an interval distribution. These models provide an alternative regression tool to the Beta regression model and fraction response model, and a tool for tail risk assessment as well.

Acknowledgments

Authors are very grateful to the third reviewer for many constructive comments. The first author is grateful to Biao Wu for many valuable conversations. Thanks also go to Clovis Sukam for his critical reading for the manuscript.

We would like to thank you for following the instructions above very closely in advance. It will definitely save us lot of time and expedite the process of your paper's publication.

Conflict of interest

The views expressed in this article are not necessarily those of Royal Bank of Canada and Scotiabank or any of their affiliates. Please direct any comments to Bill Huajian Yang at h_y02@yahoo.ca.

References

1. Basel Committee on Banking Supervision, An Explanatory Note on the Basel II IRB Risk Weight Functions, 2005. Available from: <https://www.bis.org/bcbs/irbriskweight.htm>.
2. Basel Committee on Banking Supervision, Guidance on credit risk and accounting for expected credit loss, 2015. Available from: <https://www.bis.org/bcbs/publ/d350.htm>.
3. Basel Committee on Banking Supervision, Minimum capital requirements for market risk, 2019. Available from: <https://www.bis.org/bcbs/publ/d457.htm>.

4. Cribari-Neto F and Zeileis A, (2010) Beta Regression in R. *J Stat Software* 34: 1–24.
5. Friedman J, Hastie T and Tibshirani R, (2001) *The Elements of Statistical Learning*, 2 Eds., New York: Springer series in statistics.
6. Gouriéroux C, Monfort A and Trognon A, (1984) Pseudo maximum likelihood methods: Theory. *Econometrica* 52: 681–700.
7. Huang X, Oosterlee CW and Mesters M, (2007) Computation of VaR and VaR contribution in the Vasicek portfolio credit loss model: a comparative study. *J Credit Risk* 3: 75–96.
8. Mullahy J, (1990) Regression models and transformations for Beta-distributed outcomes.
9. Murphy K, (2012) *Machine learning: a probabilistic perspective*, MIT press.
10. Papke LE and Wooldridge JM, (1996) Econometric methods for fractional response variables with an application to 401 (k) plan participation rates. *J Appl Econometrics* 11: 619–632.
11. Ramponi FA and Campi MC, (2018) Expected shortfall: Heuristics and certificates. *Eur J Operational Res* 267: 1003–1013.
12. Rosen D and Saunders D, (2009) Analytical methods for hedging systematic credit risk with linear factor portfolios. *J Econ Dyn Control* 33: 37–52.
13. Vasicek O, (1991) Limiting loan loss probability distribution. KMV Corporation.
14. Vasicek O, (1991) The distribution of loan portfolio value. *Risk* 15: 160–162.
15. Wolfinger RD, Fitting nonlinear mixed models with the new NLMIXED procedure. Proceedings of the 24th Annual SAS Users Group International Conference (SUGI 24), 1999. Available from: <https://pdfs.semanticscholar.org/>.
16. Wu B, (2019) The Probability of Default Distribution of Heterogeneous Loan Portfolio. *Curr Anal Econ Finance* 1: 88–95.
17. Yamai Y and Yoshida T, (2002) Comparative analyses of expected shortfall and value-at-risk: their estimation error, decomposition, and optimization. *Monetary Econ Stud* 20: 87–121.
18. Yang BH, (2013) Estimating Long-Run PD, Asset Correlation, and Portfolio Level PD by Vasicek Models. *J Risk Model Validation* 7: 3–19.
19. Yang BH, Wu B, Cui K et al. (2020) IFRS9 Expected Credit Loss Estimation: Advanced Models for Estimating Portfolio Loss and Weighting Scenario Losses. *J Risk Model Validation* 14: 19–34.

Appendix A

Proof of Proposition 3.5 in section 3.

B. s follows the standard logistic distribution and Φ is the CDF for the standard normal distribution.

C. s follows the standard logistic distribution and Φ is its CDF.

D. $s \sim N(0,1)$ and Φ is the CDF for standard logistic distribution.

For case B, functions ϕ and Φ in Proposition 2.1 are respectively the standard normal density and CDF. Density f for the standard logistic distribution is given by $f(x) = \exp(-x) / (1 + \exp(-x))^2$. Recall the relationship $z = \Phi^{-1}(y)$. Then $g(y, a, b)$ in (2.3) is given by

$$g(y, a, b) = \frac{U_1}{bU_2} = c \exp\left(-\frac{z-a}{b} + \frac{z^2}{2}\right) \{1 + \exp[-\frac{z-a}{b}]\}^{-2}, \quad (\text{A.1})$$

where $c = \frac{\sqrt{2\pi}}{b}$. Rewrite $f(x)$ as $f(x) = \exp(x) / (1 + \exp(x))^2$. Then $g(y, a, b)$ has the form:

$$g(y, a, b) = \frac{U_1}{bU_2} = c \exp\left(\frac{z-a}{b} + \frac{z^2}{2}\right) \{1 + \exp[\frac{z-a}{b}]\}^{-2}. \quad (\text{A.2})$$

With $z = \Phi^{-1}(y)$, $\lim_{y \rightarrow 0} z = -\infty$ and $\lim_{y \rightarrow 1} z = +\infty$. Thus $\lim_{y \rightarrow 1} g(y, a, b) = +\infty$ by (A.1), and $\lim_{y \rightarrow 0} g(y, a, b) = +\infty$ by (A.2). Hence $g(y, a, b)$ has a fat right tail and a fat left tail for all $b > 0$.

Next for tailed index, we use Lemma 2. We note that $\lim_{y \rightarrow 1} z = +\infty$, and the part $\{1 + \exp[-\frac{z-a}{b}]\}^{-2}$ in (A.1) approaches 1 when $z \rightarrow +\infty$. Hence by (A.1) and using (2.8), we have for the right tail.

$$\begin{aligned} \lim_{y \rightarrow 1} \frac{[g(y, a, b)]^{-\frac{1}{\beta}}}{1-y} &= \lim_{z \rightarrow +\infty} \frac{c_1 \exp[-\frac{1}{\beta}(-\frac{z-a}{b} + \frac{z^2}{2})]}{1-\Phi(z)} \\ &= \lim_{z \rightarrow +\infty} \frac{c_1(-\frac{1}{b} + z) \exp[-\frac{1}{\beta}(-\frac{z-a}{b} + \frac{z^2}{2})]}{\phi(z)} \end{aligned}$$

by L'Hospital's rule, where $c_1 = c^{-\frac{1}{\beta}}$. Since $\phi(z) = \frac{1}{\sqrt{2\pi}} \exp(-\frac{z^2}{2})$, the above limit is 0 when $0 < \beta < 1$, and is $+\infty$ if $\beta > 1$. Similarly, for the left tail, by (A.2), we have

$$\begin{aligned} \lim_{y \rightarrow 0} \frac{[g(y, a, b)]^{-\frac{1}{\beta}}}{y} &= \lim_{z \rightarrow -\infty} \frac{c_1 \exp[-\frac{1}{\beta}(\frac{z-a}{b} + \frac{z^2}{2})]}{\Phi(z)} \\ &= \lim_{z \rightarrow -\infty} \frac{c_1(\frac{1}{b} - z) \exp[-\frac{1}{\beta}(\frac{z-a}{b} + \frac{z^2}{2})]}{\phi(z)} \end{aligned}$$

by L'Hospital's rule, which is 0 when $0 < \beta < 1$, and is $+\infty$ if $\beta > 1$. Thus the left and right tailed indexes for case B are 1 by Lemma 2.2.

For case C, we have $\phi = f$, where $f(x) = \frac{\exp(-x)}{(1+\exp(-x))^2}$. Density $g(y, a, b)$ in Proposition 2.1 is given by

$$g(y, a, b) = \frac{u_1}{bu_2} = \frac{1}{b} \exp\left(\left(1 - \frac{1}{b}\right)z + \frac{a}{b}\right) [1 + \exp(-z)]^2 [1 + \exp(-\frac{z-a}{b})]^{-2}. \quad (\text{A.3})$$

With $z = \Phi^{-1}(y)$, we have $\lim_{y \rightarrow 0} z = -\infty$ and $\lim_{y \rightarrow 1} z = +\infty$. If $b > 1$, then $(1 - \frac{1}{b}) > 0$, hence $\lim_{y \rightarrow 1} g(y, a, b) = +\infty$ by (A.3), and $g(y, a, b)$ has a fat right tail when $b > 1$. Rewrite $\phi(x)$ as $\phi(x) = \exp(x)/(1 + \exp(x))^2$, then $g(y, a, b)$ has the form:

$$g(y, a, b) = \frac{u_1}{bu_2} = \frac{1}{b} \exp\left(\left(1 - \frac{1}{b}\right)(-z) - \frac{a}{b}\right) [1 + \exp(z)]^2 \left[1 + \exp\left(\frac{z-a}{b}\right)\right]^{-2}. \quad (\text{A.4})$$

Hence $\lim_{y \rightarrow 0} g(y, a, b) = +\infty$, as both $[1 + \exp(z)]$ and $\left[1 + \exp\left(\frac{z-a}{b}\right)\right]$ approach 1 when $z \rightarrow -\infty$. Thus $g(y, a, b)$ has a fat left tail.

Next, we note that the part $[1 + \exp(-z)]$ and $[1 + \exp(-z + a)/b]$ in (A.3) both approach 1 when $z \rightarrow +\infty$. Hence by (A.3) and using (2.8), we have

$$\begin{aligned} \lim_{z \rightarrow +\infty} \frac{[g(\Phi(z), a, b)]^{-1/\beta}}{1 - \Phi(z)} &= \lim_{z \rightarrow +\infty} \frac{c \exp[-\frac{1}{\beta}(1 - \frac{1}{b})z]}{1 - \Phi(z)} \\ &= \lim_{z \rightarrow +\infty} \frac{\frac{c}{\beta}(1 - \frac{1}{b}) \exp[-\frac{1}{\beta}(1 - \frac{1}{b})z]}{\phi(z)} \end{aligned}$$

by L'Hospital's rule, where $c = b^{\frac{1}{\beta}} \exp(-\frac{a}{\beta b})$. Since $\phi(z) = \exp(-z) / (1 + \exp(-z))^2$, the above limit is 0 if $\frac{1}{\beta}(1 - \frac{1}{b}) > 1$, and is $+\infty$ if $\frac{1}{\beta}(1 - \frac{1}{b}) < 1$. This means $g(y, a, b)$ has a fat right tail with tailed index $(1 - 1/b)$ when $b > 1$. Similarly, for the left tail, we have by (A.4)

$$\lim_{y \rightarrow 0} \frac{[g(y, a, b)]^{-1/\beta}}{y} = \lim_{z \rightarrow -\infty} \frac{c \exp[-\frac{1}{\beta}(1 - \frac{1}{b})(-z)]}{\Phi(z)} = \lim_{z \rightarrow -\infty} \frac{\frac{c}{\beta}(1 - \frac{1}{b}) \exp[\frac{1}{\beta}(1 - \frac{1}{b})z]}{\phi(z)}$$

by L'Hospital's rule, where $c = b^{\frac{1}{\beta}} \exp(\frac{a}{\beta b})$. Using $\phi(z) = \frac{\exp(z)}{(1 + \exp(z))^2}$, we see the above limit is 0 for $\frac{1}{\beta}(1 - \frac{1}{b}) > 1$, and is $+\infty$ if $\frac{1}{\beta}(1 - \frac{1}{b}) < 1$. This means $g(y, a, b)$ has a tailed index $(1 - 1/b)$ for the left tail when $b > 1$.

For case D, functions ϕ and Φ are respectively the density and the CDF for the standard logistic distribution, and f is the standard normal density. Since $\phi(z) = \frac{\exp(-z)}{(1 + \exp(-z))^2}$, by Proposition 2.1,

$$g(y, a, b) = \frac{U_1}{bU_2} = c \exp\left(z - \frac{(z-a)^2}{2b^2}\right) [1 + \exp(-z)]^2, \quad (\text{A.5})$$

where $c = \frac{1}{b\sqrt{2\pi}}$. Write $\phi(z)$ as $\phi(z) = \exp(z) / (1 + \exp(z))^2$. Then $g(y, a, b)$ has another form:


$$g(y, a, b) = \frac{U_1}{bU_2} = c \exp\left(-z - \frac{(z-a)^2}{2b^2}\right) [1 + \exp(z)]^2. \quad (\text{A.6})$$

With $z = \Phi^{-1}(y)$, we have $\lim_{y \rightarrow 0} z = -\infty$ and $\lim_{y \rightarrow 1} z = +\infty$. Thus $\lim_{y \rightarrow 1} g(y, a, b) = 0$ by (A.5), and $\lim_{y \rightarrow 0} g(y, a, b) = 0$ by (A.6). Hence $g(y, a, b)$ does not have a fat right tail, neither a fat left tail.

Appendix B

Sketches of density plots for interval distributions A-C in section 3.

We focus on interval distributions given as the examples in section 3 with fat tails. This includes distributions A, B, and C. Figures B1, B2, and B3 below plot respectively the probability densities for these three types of interval distributions. Values for parameters a and b are given at the top of the chart. Two densities are included for each of three type distributions, with different values for parameter b but the same value for parameter a .



It is observed that, with $b > 1$, all curves have a fat right tail and a fat left tail, and the curve with a higher parameter b climbs up the tails more quickly than the other, as the latter has lower tailed index. The roughness in the middle is a consequence of squeezing the values in x -axis, in order to put together the left and right tails into the chart (see the value crossing in x -axis from left to right). It does not necessarily reflect the true behaviours of the curves.



AIMS Press

© 2020 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)