



---

*Research article*

## **A multiple objective programming approach to linear bilevel multi-follower programming**

**Habibe Sadeghi\* and Fatemeh Moslemi**

Department of Mathematics, Faculty of Mathematical Sciences and Computer, Shahid Chamran University of Ahvaz, Ahvaz, Iran

\* **Correspondence:** Email: [habibe.sadeghi@scu.ac.ir](mailto:habibe.sadeghi@scu.ac.ir).

**Abstract:** In this paper, we investigate the relationship between a certain class of linear bilevel multi-follower programming problems and multiple objective programming. We introduce two multiple objective linear programming problems with different objective functions and the same constraint region. We show that the extreme points of the set of efficient solutions for both problems are the same as those of the set of feasible solutions to the linear bilevel multi-follower programming problem. Based on this relationship, a new algorithm to find an optimal solution for the linear bilevel multi-follower programming problem is developed. Some numerical examples are presented to show the feasibility of the proposed algorithm.

**Keywords:** linear bilevel programming; multi-follower; multiple objective programming; efficient set

**Mathematics Subject Classification:** 90C29, 90C08

---

### **1. Introduction**

Multilevel programming (MLP) is developed to model the decentralized decision-making situations wherein decision makers are arranged within a hierarchical structure. In a MLP problem, when only two levels exist, the problem is referred to as the bilevel programming (BLP) problem. In actual, BLP problems involve two nested optimization problems where the constraint region of first-level problem contains the second-level optimization problem. In this model, a decision maker at the first-level is termed as a leader, and at the second-level, a follower. BLP problems have many applications in the real world, such as those in supply chain management [14], planning [16], and logistics [19]. Most of the research concerning MLP have focused on bilevel programming (BLP) [1, 5]. In this paper, we consider linear bilevel problems with one leader and multiple followers. This model is called linear bilevel multi-follower programming (LBLMFP) problem. Shi et al. extended the Kth-Best algorithm to solve LBLMFP problems [21]. Several researchers developed

heuristic algorithms to the BLP problems, for instance, see [10]. Lu et al, applied the extended Kuhn-Tucker approach to solve LBLMFP problem [13]. Also, one approach is developed to solve the LBLMFP problems based on multi-parametric programming [7]. In addition, some researchers have exploited multiple objective programming techniques to solve BLP problems [9, 18]. These kinds of solution approaches are developed based on some relationships between BLP and multiple objective programming that were first presented by Fülöp [8]. For example, Glackin et al. presented an algorithm to solve linear bilevel programming (LBLEP) problems based on multiple objective linear programming (MOLP) techniques [9].

In this paper, Motivated by the relationship between LBLEP and MOLP, a relationship between a class of LBLMFP problems and MOLP problems is introduced for the first time. Moreover, we present an algorithm for solving LBLMFP problems based on the proposed relation.

The paper is organized as follows: In the next section, we present the basic definitions, and some notions about LBLMFP problems and MOLP problems. In Section 3, we introduce two MOLP problems such that each feasible solution for the LBLMFP problem is efficient for both of the two MOLP problems. Next, we obtain results, based on which the LBLMFP problem can be reduced to optimize the first-level objective function over a certain efficient set. Based on this result, a new algorithm is developed in Section 4 for solving the LBLMFP problem. Furthermore, we obtain some results for special cases. Section 5 presents a number of numerical examples to illustrate the proposed algorithm. Finally, in Section 6, we present the conclusions.

## 2. Preliminaries

In this section, we present the formulation of LBLMFP problem which we shall investigate, accompanied by basic definitions. In addition, we recall some notions and theoretical properties about MOLP problems.

### 2.1. Linear bilevel multi-follower programming problem

In this article, we consider a linear bilevel programming (LBLEP) problem with two followers. There are individual decision variables in separate objective functions and constraints between the followers. Each follower takes other followers' decision variables as a reference. This is called a reference-uncooperative LBLMFP problem [23]. This model can be formulated as follows:

$$\begin{aligned} \min_{x^1 \in X^1} F(x^1, x^2, x^3) &= (c^{11})^T x^1 + (c^{12})^T x^2 + (c^{13})^T x^3, \\ \min_{x^2 \in X^2} f_1(x^2, x^3) &= (c^{22})^T x^2 + (c^{23})^T x^3, \\ \text{s.t. } A_1 x^1 + A_2 x^2 + A_3 x^3 &\leq b_1, \\ \min_{x^3 \in X^3} f_2(x^2, x^3) &= (c^{32})^T x^2 + (c^{33})^T x^3, \\ \text{s.t. } B_1 x^1 + B_2 x^2 + B_3 x^3 &\leq b_2, \end{aligned}$$

where  $x^i \in X^i \subset \mathbb{R}_+^{n_i}$ ,  $i = 1, 2, 3$ ,  $F : X^1 \times X^2 \times X^3 \rightarrow \mathbb{R}$ , and  $f_k : X^2 \times X^3 \rightarrow \mathbb{R}$ ,  $k = 1, 2$ ,  $c^{ij}$ , for  $i, j \in \{1, 2, 3\}$ , are the vectors of conformal dimension, and for each  $i \in \{1, 2, 3\}$ ,  $A_i$  is a  $m \times n_i$  matrix,  $B_i$  is a  $q \times n_i$  matrix,  $b_1 \in \mathbb{R}^m$ , and  $b_2 \in \mathbb{R}^q$ . Also, although  $x^i \geq 0$ , for  $i = 1, 2, 3$ , do not explicitly

appear in this problem, we assume that they exist in the set of constraints.

Notice that, the followers' objective functions are linear in  $x^2$  and  $x^3$  and for each follower, the value for the variable  $x^1$  is given. Thus, a problem equivalent to the LBLMFP problem is obtained if the followers' objective functions are replaced by  $\sum_{j=1}^3 (c^{ij})^T x^j$ , for  $i=2,3$ .

We need to introduce the following definitions that can be found in [23].

(1) Constraint region of the LBLMFP problem:

$$S := \{(x^1, x^2, x^3) \in X^1 \times X^2 \times X^3 : A_1 x^1 + A_2 x^2 + A_3 x^3 \leq b_1, B_1 x^1 + B_2 x^2 + B_3 x^3 \leq b_2\}.$$

We suppose that  $S$  is non-empty and compact.

(2) Projection of  $S$  onto the leader's decision space:

$$S(X^1) := \{x^1 \in X^1 : \exists (x^2, x^3) \in X^2 \times X^3, (x^1, x^2, x^3) \in S\}.$$

(3) Feasible sets for the first and second followers, respectively:

$$S_1(x^1, x^3) := \{x^2 \in X^2 : (x^1, x^2, x^3) \in S\},$$

$$S_2(x^1, x^2) := \{x^3 \in X^3 : (x^1, x^2, x^3) \in S\}.$$

The feasible region of each follower is affected by the leader's choice of  $x^1$ , and the other followers' decisions.

(4) The first and second followers' rational reaction sets, respectively:

$$P_1(x^1, x^3) := \{x^2 : x^2 \in \operatorname{argmin}[f_1(\hat{x}^2, x^3) : \hat{x}^2 \in S_1(x^1, x^3)]\},$$

where

$$\operatorname{argmin}[f_1(\hat{x}^2, x^3) : \hat{x}^2 \in S_1(x^1, x^3)] = \{x^2 \in S_1(x^1, x^3) : f_1(x^2, x^3) \leq f_1(\hat{x}^2, x^3), \\ \forall \hat{x}^2 \in S_1(x^1, x^3)\},$$

and

$$P_2(x^1, x^2) := \{x^3 : x^3 \in \operatorname{argmin}[f_2(x^2, \hat{x}^3) : \hat{x}^3 \in S_2(x^1, x^2)]\},$$

where

$$\operatorname{argmin}[f_2(x^2, \hat{x}^3) : \hat{x}^3 \in S_2(x^1, x^2)] = \{x^3 \in S_2(x^1, x^2) : f_2(x^2, x^3) \leq f_2(x^2, \hat{x}^3), \\ \forall \hat{x}^3 \in S_2(x^1, x^2)\}.$$

(5) Inducible region ( $IR$ ):

$$IR := \{(x^1, x^2, x^3) \in S : x^2 \in P_1(x^1, x^3), x^3 \in P_2(x^1, x^2)\}.$$

In actual, the inducible region is the set of feasible solutions of the LBLMFP problem. Therefore, determining the solution for the LBLMFP problem is equivalent to solving the following problem:

$$\min\{F(x^1, x^2, x^3) : (x^1, x^2, x^3) \in IR\}. \quad (2.1)$$

In order to assure that (2.1) has an optimal solution [23], we supposed that the following assumptions hold:

(1)  $IR$  is non-empty.

(2)  $P_1(x^1, x^3)$  and  $P_2(x^1, x^2)$  are point-to-point mappings with respect to  $(x^1, x^3)$  and  $(x^1, x^2)$ , respectively. In other words, for each leader's choice of  $x^1$ , there will be a unique solution to each follower.

### 2.1.1. Multiple objective linear programming problem

Assume that  $p \geq 2$  is an integer and  $c_i \in \mathbb{R}^n$ ,  $i = 1, 2, \dots, p$  are row vectors. Let  $C$  be a  $p \times n$  matrix whose  $i$ -th row is given by  $c_i$ ,  $i = 1, 2, \dots, p$ , and  $U$  is a non-empty, compact and convex polyhedral set in  $\mathbb{R}^n$ . A multiple objective linear programming (MOLP) problem is formulated in general as follows:

$$\min\{Cx : x \in U\}, \quad (\text{MOLP})$$

where  $U$  is called feasible region.

**Definition 1.** [22] A feasible point,  $\bar{x} \in U$ , is called an efficient solution if there exists no  $x \in U$  such that  $Cx \leq C\bar{x}$  and  $Cx \neq C\bar{x}$ .

An efficient solution is also called a Pareto-optimal solution.

Let  $E$  denote the set of all efficient solutions of the MOLP problem. Note that  $E \neq \emptyset$  [[6], Theorem 2.19].

Let  $d \in \mathbb{R}^n$ . Consider the following mathematical programming problem:

$$\min\{d^T x : x \in E\}. \quad (2.2)$$

This problem is a non-convex linear optimization problem over the efficient set  $E$  of the MOLP problem. Let  $E^*$  denote the set of optimal solutions to the problem (2.2) and  $U_{ex}$  denote the set of extreme points of the polyhedron  $U$ . Because  $U$  is a non-empty, compact and convex polyhedron in  $\mathbb{R}^n$ ,  $U_{ex}$  is non-empty [[3], Theorem 2.6.5]. So, the following result holds:

**Theorem 1.** Let  $U$  be a non-empty, compact, and convex polyhedral set, and let  $E^*$  be non-empty. Then, at least one element of  $E^*$  is an extreme point of  $U$  [[4], Theorem 4.5].

By considering the relation  $E \subseteq U$ , we immediately get the following corollary to Theorem 1.

**Corollary 1.** Let  $E^* \neq \emptyset$ . Then, there is an extreme point of  $E$ , which is an optimal solution to the problem (2.2).

From Theorem 3.40 in [6], because  $U$  is a non-empty, compact and convex polyhedron in  $\mathbb{R}^n$ , and the objective functions  $c_i x$ ,  $i = 1, 2, \dots, p$  are convex, we can conclude that the efficient set  $E$  is connected.

Moreover, it is well-known that the efficient set  $E$ , is comprised of a finite union of faces of  $U$  [6]. Therefore, we obtain the following corollary:

**Corollary 2.** The efficient set  $E$  is comprised of a finite union of connected faces of the polyhedron  $U$ .

The Corollary 2 obviously implies that the efficient set  $E$  is closed.

Also, the following definition is used in sequel:

**Definition 2.** [22] Let  $U \subseteq \mathbb{R}^n$ , and  $x_1, x_2, \dots, x_r \in U$ . The notation  $\gamma(x_1, x_2, \dots, x_r)$  denotes the set of all convex combinations of  $x_1, x_2, \dots, x_r$ .

### 3. Theoretical properties

In this section, we will introduce two MOLP problems in such a way that any extreme efficient solution for both problems simultaneously, is an extreme feasible solution (in other words, an extreme point of  $IR$ ) to LBLMFP problem.

Note that  $\bar{x} = (\bar{x}^1, \bar{x}^2, \bar{x}^3) \in IR$ , i.e.,  $\bar{x}^2 \in P_1(\bar{x}^1, \bar{x}^3)$  and  $\bar{x}^3 \in P_2(\bar{x}^1, \bar{x}^2)$ . Therefore,  $\bar{x}^2$  is an optimal solution for the following linear program (LP):

$$\min_{x^2 \in X^2} f_1(x^2, \bar{x}^3) = (c^{22})^T x^2 + (c^{23})^T \bar{x}^3, \quad (3.1)$$

$$s.t. A_2 x^2 \leq b_1 - A_1 \bar{x}^1 - A_3 \bar{x}^3, \quad (3.2)$$

and  $\bar{x}^3$  is an optimal solution for the following LP:

$$\min_{x^3 \in X^3} f_2(\bar{x}^2, x^3) = (c^{32})^T \bar{x}^2 + (c^{33})^T x^3, \quad (3.3)$$

$$s.t. B_3 x^3 \leq b_2 - B_1 \bar{x}^1 - B_2 \bar{x}^2. \quad (3.4)$$

Now, set  $n := \sum_{i=1}^3 n_i$ . We construct augmented matrix  $\mathcal{A} = [A_1 \ A_3]$ . Let  $rank(\mathcal{A}) = r_1$ . Without

loss of generality, we assume that augmented matrix  $\mathcal{A}$  can be rewritten as  $\begin{bmatrix} \bar{A}_1 & \bar{A}_3 \\ \hat{A}_1 & \hat{A}_3 \end{bmatrix}$ , where  $\begin{bmatrix} \bar{A}_1 & \bar{A}_3 \end{bmatrix}$

is  $r_1 \times (n_1 + n_3)$  matrix and  $rank(\begin{bmatrix} \bar{A}_1 & \bar{A}_3 \end{bmatrix}) = rank(\mathcal{A}) = r_1$ . [Notice that the augmented matrix  $\mathcal{A}$  contains an  $(n_1 + n_3) \times (n_1 + n_3)$  identity submatrix corresponding to non-negativity constraints. So,  $r_1 = n_1 + n_3$ ]. Similarly, we construct augmented matrix  $\mathcal{B} = [B_1 \ B_2]$ . Let  $rank(\mathcal{B}) = r_2$

and augmented matrix  $\mathcal{B}$  can be rewritten as  $\begin{bmatrix} \bar{B}_1 & \bar{B}_2 \\ \hat{B}_1 & \hat{B}_2 \end{bmatrix}$ , where  $\begin{bmatrix} \bar{B}_1 & \bar{B}_2 \end{bmatrix}$  is  $r_2 \times (n_1 + n_2)$  matrix and

$rank(\begin{bmatrix} \bar{B}_1 & \bar{B}_2 \end{bmatrix}) = rank(\mathcal{B}) = r_2$ . Let  $k_i = r_i + 2$ ,  $i = 1, 2$  and the  $k_i \times n$ , criterion matrices  $C_i$ ,  $i = 1, 2$  be defined as follows:

$$C_1 := \begin{bmatrix} \bar{A}_1 & O & \bar{A}_3 \\ -e^T \bar{A}_1 & 0 & -e^T \bar{A}_3 \\ 0 & (c^{22})^T & 0 \end{bmatrix}, \quad C_2 := \begin{bmatrix} \bar{B}_1 & \bar{B}_2 & O \\ -e^T \bar{B}_1 & -e^T \bar{B}_2 & 0 \\ 0 & 0 & (c^{33})^T \end{bmatrix}, \quad (3.5)$$

where  $O$  and  $0$ , are zero matrix and zero vectors of conformal dimension, respectively. Also,  $e$  is a vector having each entry equal to 1.

Next, consider the following two MOLP problems:

$$\min\{C_i x : x \in S\}, \quad i = 1, 2. \quad (\text{MOLP}_i)$$

Let  $E_i$  be the set of efficient solutions of the MOLP $_i$ ,  $i = 1, 2$ . We have the following result:

**Proposition 1.** Let  $IR$  and  $E_1$  be defined as above. Then,  $IR \subseteq E_1$ .

*Proof.* Assume that  $\bar{x} = (\bar{x}^1, \bar{x}^2, \bar{x}^3) \in IR$ . It suffices to show that  $\bar{x}$  is an efficient solution for the MOLP1. Let us suppose the contrary, i.e., there exists  $x = (x^1, x^2, x^3) \in S$  such that  $C_1x \leq C_1\bar{x}$  and  $C_1x \neq C_1\bar{x}$ . Because  $C_1x \leq C_1\bar{x}$ , using the structure of matrix  $C_1$ , one obtains:

$$\bar{A}_1x^1 + \bar{A}_3x^3 \leq \bar{A}_1\bar{x}^1 + \bar{A}_3\bar{x}^3, \quad (3.6)$$

$$-e^T\bar{A}_1x^1 - e^T\bar{A}_3x^3 \leq -e^T\bar{A}_1\bar{x}^1 - e^T\bar{A}_3\bar{x}^3. \quad (3.7)$$

From inequality (3.6), one obtains:

$$-e^T\bar{A}_1x^1 - e^T\bar{A}_3x^3 \geq -e^T\bar{A}_1\bar{x}^1 - e^T\bar{A}_3\bar{x}^3. \quad (3.8)$$

The last two inequalities imply that:

$$e^T\bar{A}_1x^1 + e^T\bar{A}_3x^3 = e^T\bar{A}_1\bar{x}^1 + e^T\bar{A}_3\bar{x}^3. \quad (3.9)$$

From equality (3.9) and inequality (3.6), it is easy to see that  $\bar{A}_1x^1 + \bar{A}_3x^3 = \bar{A}_1\bar{x}^1 + \bar{A}_3\bar{x}^3$ . So, we have:

$$A_1x^1 + A_3x^3 = A_1\bar{x}^1 + A_3\bar{x}^3. \quad (3.10)$$

Obviously,  $x^2$  is a feasible solution for the problem (3.1)-(3.2). Moreover, due to  $C_1x \neq C_1\bar{x}$  and in view of equalities (3.9) and (3.10), we deduce that  $(c^{22})^T x^2 < (c^{22})^T \bar{x}^2$  which contradicts  $\bar{x}^2$  is an optimal solution for the problem (3.1)-(3.2). This completes the proof.  $\square$

The proof of the following proposition is similar to that of Proposition 1, and we omit it.

**Proposition 2.** Let  $IR$  and  $E_2$  be defined as above. Then,  $IR \subseteq E_2$ .

Now, if we set  $E := E_1 \cap E_2$ , from Propositions 1 and 2, we can get:

**Remark 1.** Let  $IR$ ,  $E_1$ , and  $E_2$  be defined as above. Then,  $IR \subseteq E_1 \cap E_2 = E$ , and regarding to Assumption (1),  $E$  is a non-empty set.

Therefore, this allows us to prove the main result of this section.

**Theorem 2.** The extreme points of  $IR$  and  $E$  are identical.

*Proof.* Let  $x = (x^1, x^2, x^3) \in IR_{ex}$  be arbitrary. Then,  $x$  is an extreme point of  $S$  [[23], Corollary 4.9]. Because  $IR \subseteq E$ , we get  $x \in E$ . Therefore,  $x$  is an extreme point of  $E$  (Note that  $E \subseteq S$ ). Since  $x \in IR_{ex}$  was chosen arbitrarily, we obtain  $IR_{ex} \subseteq E_{ex}$ . Now, we will show that  $E_{ex} \subseteq IR_{ex}$ . Let  $x = (x^1, x^2, x^3) \in E_{ex}$  be chosen arbitrarily. There are two cases to be considered here.

Case 1:  $x \notin IR$ . In this case, we conclude from the definition of  $IR$  that at least one of the following two subcases should occur:

Subcase 1:  $x^2 \notin P_1(x^1, x^3)$ . It follows from Assumption (2) that there exists  $\hat{x}^2 \in P_1(x^1, x^3)$ , so that  $f_1(\hat{x}^2, x^3) < f_1(x^2, x^3)$ . As a consequence, we get:

$$(c^{22})^T \hat{x}^2 < (c^{22})^T x^2. \quad (3.11)$$

Set  $\hat{x} = (x^1, \hat{x}^2, x^3)$ . By definition of  $P_1(x^1, x^3)$ ,  $\hat{x} \in S$ . Due to the structure of matrix  $C_1$  and inequality (3.11), one has  $C_1 \hat{x} \leq C_1 x$ ,  $C_1 \hat{x} \neq C_1 x$  that contradicts  $x \in E \subseteq E_1$ .

Subcase 2:  $x^3 \notin P_2(x^1, x^2)$ . By Assumption (2), there exists  $\hat{x}^3 \in P_2(x^1, x^2)$  so that  $f_2(x^2, \hat{x}^3) < f_2(x^2, x^3)$ . As a consequence, we have:

$$(c^{33})^T \hat{x}^3 < (c^{33})^T x^3. \quad (3.12)$$

Set  $\hat{x} = (x^1, x^2, \hat{x}^3)$ . By definition of  $P_2(x^1, x^2)$ ,  $\hat{x} \in S$ . Due to the structure of matrix  $C_2$  and inequality (3.12), one has  $C_2 \hat{x} \leq C_2 x$ ,  $C_2 \hat{x} \neq C_2 x$ , which contradicts  $x \in E \subseteq E_2$ .

Case 2:  $x \in IR$ . Since  $x \in E_{ex}$  and  $IR \subseteq E$  (see Remark 1), we conclude that  $x \in IR_{ex}$ . This completes the proof.  $\square$

Now, let us consider the following problem:

$$\min\{F(x^1, x^2, x^3) = (c^{11})^T x^1 + (c^{12})^T x^2 + (c^{13})^T x^3 : (x^1, x^2, x^3) \in E\}. \quad (3.13)$$

**Remark 2.** Note that Corollary 2 obviously implies that the efficient set  $E$  is closed. Also, since  $E$  is a closed subset of the compact polyhedral set of  $S$ , itself is a compact set. Therefore, problem (3.13) involves the optimization of a linear function over a compact set. Hence, there exists an optimal solution to the problem (3.13), i.e.,  $E^* \neq \emptyset$ . Then, according to the Corollary 1, there is an extreme point of  $E$ , which is an optimal solution for the problem (3.13).

The relation between LBLMFP problem and problem (3.13) is stated in the following theorem.

**Theorem 3.** A point  $\bar{x} = (\bar{x}^1, \bar{x}^2, \bar{x}^3) \in IR_{ex}$  is an optimal solution to LBLMFP problem if, and only if, it is an optimal solution to the problem (3.13).

*Proof.* It is well-known that solving LBLMFP problem is equivalent to solving problem (2.1). Furthermore, there exists an extreme point of  $IR$ , which is an optimal solution for problem (2.1) [23]. Because the optimal solution of problem (3.13) occurs at an extreme point of  $E$  and the extreme points of  $E$  and  $IR$  are the same, we conclude that if  $\bar{x}$  is an optimal solution for problem (3.13), it is an optimal solution for LBLMFP problem, and vice versa. This completes the proof.  $\square$

In the next section, based on Theorem 3, we propose a new algorithm to solve LBLMFP problem.

#### 4. The Algorithm

Owing to the preceding discussion, we will propose a new algorithm for solving LBLMFP problems. First, one can solve two MOLPi,  $i = 1, 2$  in order to obtain the sets  $E_i$ ,  $i = 1, 2$ . Let  $E$  be the set of efficient points that have been discovered to be efficient solutions to both problems. Then, one can minimize the leader objective function over the set  $S$ . If the obtained extreme optimal solution lies in the set  $E$ , it solves the LBLMFP problem. Otherwise, one can solve problem (3.13). Then, by Theorem 3, the obtained extreme optimal solution solves the LBLMFP problem. The algorithm can be described as follows:

The Algorithm:

**Step 1.** Construct the MOLPi,  $i = 1, 2$ .

**Step 2.** Find efficient sets  $E_i$  with respect to the MOLP $_i$ ,  $i = 1, 2$ .

- For instance, approaches presented in [6, 17, 20, 22] can be used in Step 2.

**Step 3.** Set  $E = E_1 \cap E_2$ .

**Step 4.** Solve the following LP:

$$\min\{F(x^1, x^2, x^3) = (c^{11})^T x^1 + (c^{12})^T x^2 + (c^{13})^T x^3 : (x^1, x^2, x^3) \in S\}. \quad (4.1)$$

Let  $x^*$  be an optimal solution (One can use the common known solution methods for LPs in [2]). If  $x^* \in E$ , stop. Then,  $x^*$  is an optimal solution to LBLMFP problem. Otherwise, go to Step 5.

**Step 5.** Find an optimal solution to problem (3.13) using, for instance, approaches developed in [4, 11, 12, 15]. Let  $x^*$  be an optimal solution for problem (3.13). Then, it is an optimal solution to LBLMFP problem, too.

- Notice that, by Remark 2, there exists an extreme point of  $E$  which is an optimal solution for problem (3.13). Hence, in Step 5, for a few number of variables and extreme points, we can find an optimal solution by picking the minimum objective function value among all extreme points of  $E$ .

Note that because the efficient set  $E$  is not convex in general, the problem (3.13) is a non-convex optimization problem. Thus, in Step 5, we face a non-convex optimization problem that is generally difficult to solve. Moreover, the algorithm stops by obtaining an efficient optimal solution. The Step 4 solves the LP (4.1) and checks whether  $x^* \in E$ . If  $x^* \in E$ , the algorithm stops at Step 4, and does not enter Step 5. Therefore, Step 4 is necessary in this algorithm to reduce the computations in some cases.

Prior to applying the proposed algorithm for solving numerical examples, we will state the following lemma:

**Lemma 1.** Let  $E$  be defined as above. Then  $E \subseteq \text{conv}IR$ , where  $\text{conv}IR$  be the smallest convex set containing  $IR$ .

*Proof.* Let  $\hat{x} \in E$  be chosen arbitrarily. Suppose the contrary, that is  $\hat{x} \notin \text{conv}IR$ . Because  $IR$  is closed [[23], Theorem 4.9],  $\text{conv}IR$  is a convex and closed set. Therefore, there exists a non-zero vector  $P$  [[3], Theorem 2.4.4], such that  $P^T \hat{x} < P^T y$ , for all  $y \in \text{conv}IR$ . Since  $S$  is a compact set and  $\text{conv}IR$  is a closed subset of  $S$ ,  $\text{conv}IR$  is a compact set as well. Furthermore, because  $y \in \text{conv}IR$ , according to the Representation Theorem [3], we have  $y = \sum_{i=1}^n \lambda_i x_i$ ,  $\sum_{i=1}^n \lambda_i = 1$ ,  $0 \leq \lambda_i \leq 1$ ,  $x_i \in (\text{conv}IR)_{ex}$ . On the other hand,  $(\text{conv}IR)_{ex} = IR_{ex} = E_{ex}$ , thus,

$$P^T \hat{x} < P^T \left( \sum_{i=1}^n \lambda_i x_i \right), \quad \sum_{i=1}^n \lambda_i = 1, \quad 0 \leq \lambda_i \leq 1, \quad x_i \in E_{ex}. \quad (4.2)$$

Now, consider the problem  $\min\{P^T x : x \in E\}$ . By Remark 2, there exists  $x_j \in E_{ex}$ , which is an optimal solution for this problem. Then,  $P^T x_j \leq P^T x$ , for all  $x \in E$ . Because  $\hat{x} \in E$ , we have:

$$P^T x_j \leq P^T \hat{x}. \quad (4.3)$$

Since  $x_j \in E_{ex}$ , it follows from (4.2) that  $P^T \hat{x} < P^T x_j$ , which contradicts (4.3). Hence,  $\hat{x} \in \text{conv}IR$ .  $\square$



Note that the efficient set  $E$  and the inducible region  $IR$  are not convex sets, generally [1, 5, 6, 22]. Up to now, we have proved that  $IR \subseteq E \subseteq \text{conv}IR$ . In the following, we show that these inclusions convert to equality in some special cases, as convexity.

**Corollary 3.** *If the efficient set  $E$  is a convex set, then  $E = \text{conv}IR$ .*

*Proof.* By the convexity of  $E$  and from Remark 1, we obtain  $\text{conv}IR \subseteq E$ . Also, according to Lemma 1, we have  $E \subseteq \text{conv}IR$ . Then,  $E = \text{conv}IR$ .  $\square$

**Corollary 4.** *If the inducible region  $IR$  is a convex set, then  $E = IR$ .*

*Proof.* According to Remark 1,  $IR \subseteq E$ . Because  $IR$  is a convex set,  $\text{conv}IR = IR$ . It follows from Lemma 1 that  $E \subseteq IR$ . Then,  $E = IR$ .  $\square$

## 5. Numerical examples

In this section, we apply the proposed algorithm for solving some numerical examples.

**Example 1.** Consider the following LBLMFP problem:

$$\begin{aligned} \min_{x^1} F(x^1, x^2, x^3) &= 3x^1 + x^2 + x^3, \\ \min_{x^2} f_1(x^2, x^3) &= -x^2 + x^3, \\ \text{s.t. } x^1 + x^2 &\leq 8, \quad x^1 + 4x^2 \geq 8, \\ x^1 + 2x^2 &\leq 13, \quad 7x^1 - 2x^2 \geq 0, \\ x^1 &\geq 0, \quad x^2 \geq 0, \quad x^3 \geq 0, \\ \min_{x^3} f_2(x^2, x^3) &= 2x^3, \\ \text{s.t. } x^1 &\geq 0, \quad x^2 \geq 0, \quad 0 \leq x^3 \leq 2. \end{aligned}$$


In this problem,  $S = \{(x^1, x^2, x^3) : x^1 + x^2 \leq 8, x^1 + 4x^2 \geq 8, x^1 + 2x^2 \leq 13, 7x^1 - 2x^2 \geq 0, x^1 \geq 0, x^2 \geq 0, 0 \leq x^3 \leq 2\}$ , and the extreme points of  $S$  are as follows:

$$\begin{aligned} a_1 &= (8, 0, 0), \quad a_2 = (8, 0, 2), \quad a_3 = (3, 5, 0), \quad a_4 = (3, 5, 2), \\ a_5 &= \left(\frac{13}{8}, \frac{91}{16}, 0\right), \quad a_6 = \left(\frac{13}{8}, \frac{91}{16}, 2\right), \quad a_7 = \left(\frac{8}{15}, \frac{28}{15}, 0\right), \quad a_8 = \left(\frac{8}{15}, \frac{28}{15}, 2\right). \end{aligned}$$

By [7], one can obtain:

$$\begin{aligned} IR &= \{(x^1, x^2, x^3) : \frac{8}{15} \leq x^1 \leq \frac{13}{8}, x^2 = \frac{7}{2}x^1, x^3 = 0\} \\ &\cup \{(x^1, x^2, x^3) : \frac{13}{8} \leq x^1 \leq 3, x^2 = \frac{13}{2} - \frac{1}{2}x^1, x^3 = 0\} \\ &\cup \{(x^1, x^2, x^3) : 3 \leq x^1 \leq 8, x^2 = 8 - x^1, x^3 = 0\}. \end{aligned}$$

The Figure 1 displays  $S$  and  $IR$ . The inducible region  $IR$  is denoted by the hatched lines.



**Figure 1.** Constraint region and  $IR$  of Example 1.

Note that in this example,  $S$  and  $IR$  are non-empty and  $S$  is compact. Also, one can obtain:

$$\begin{aligned}
 P_1(x^1, x^3) &= \{(x^1, x^2, x^3) \in S : x^2 = \frac{7}{2}x^1, \frac{8}{15} \leq x^1 \leq \frac{13}{8}\} \\
 &\cup \{(x^1, x^2, x^3) \in S : x^2 = \frac{13}{2} - \frac{1}{2}x^1, \frac{13}{8} \leq x^1 \leq 3\} \\
 &\cup \{(x^1, x^2, x^3) \in S : x^2 = 8 - x^1, 3 \leq x^1 \leq 8\}, \\
 P_2(x^1, x^2) &= \{(x^1, x^2, x^3) \in S : x^3 = 0\}.
 \end{aligned}$$

Hence,  $P_1(x^1, x^3) \neq \emptyset$ ,  $P_2(x^1, x^2) \neq \emptyset$ , and  $P_1(x^1, x^3)$  and  $P_2(x^1, x^2)$  are point-to-point maps with respect to  $(x^1, x^3)$  and  $(x^1, x^2)$ , respectively. Therefore, the Assumptions (1) and (2) hold, and we can solve this problem by the proposed algorithm. Using the proposed algorithm, the process is as follows:

**Step 1.** The MOLP $_i$  problems, for  $i = 1, 2$  are constructed as follows, respectively:


$$\begin{aligned}
 &\min\{x^1, x^3, -x^1 - x^3, -x^2\} : (x^1, x^2, x^3) \in S, \\
 &\min\{x^1, x^2, -x^1 - x^2, 2x^3\} : (x^1, x^2, x^3) \in S.
 \end{aligned}$$

**Step 2.** In order to find the sets  $E_i$ ,  $i = 1, 2$ , using the approach described in [20], one can obtain:

$$\begin{aligned}
 E_1 &= \gamma(a_1, a_3, a_4, a_2) \cup \gamma(a_3, a_5, a_6, a_4) \cup \gamma(a_5, a_7, a_8, a_6), \\
 E_2 &= \gamma(a_1, a_3, a_5, a_7).
 \end{aligned}$$

The sets  $E_i$ ,  $i = 1, 2$  are shown in Figure 2 (a) and (b), by the hatched regions and gray area, respectively.

**Step 3.** Set  $E = E_1 \cap E_2$ . We obtain  $E = \gamma(a_1, a_3) \cup \gamma(a_3, a_5) \cup \gamma(a_5, a_7) = IR$  which is shown in Figure 1 by the hatched lines.



**Figure 2.** Constraint region,  $E_1$  and  $E_2$  of Example 1.

**Step 4.** Solve the following LP:

$$\min\{3x^1 + x^2 + x^3 : (x^1, x^2, x^3) \in S\}.$$

The optimal solution is  $x^* = (\frac{8}{15}, \frac{28}{15}, 0)$ . Because  $x^* \in E$ , stop. We deduce that  $x^*$  is an optimal solution for this problem. Also, we solve this example using the multi-parametric approach [7]. We obtain  $x^* = (\frac{8}{15}, \frac{28}{15}, 0)$  which is equal with the obtained optimal solution using the proposed algorithm.

**Example 2.** Consider the following LBLMFP problem:

$$\begin{aligned} \min_{x^1} F(x^1, x^2, x^3) &= 3x^1 + x^2 - x^3, \\ \min_{x^2} f_1(x^2, x^3) &= 2x^2 - 3x^3, \\ \text{s.t. } x^1 + x^2 &\geq 1, x^1 \geq 0, \\ &x^2 \geq 0, x^3 \geq 0, \\ \min_{x^3} f_2(x^2, x^3) &= -4x^2 + x^3, \\ \text{s.t. } 2x^1 + x^2 + x^3 &\leq 5, \\ &x^1 \geq 0, x^2 \geq 0, x^3 \geq 0. \end{aligned}$$


In this problem,  $S = \{(x^1, x^2, x^3) : 2x^1 + x^2 + x^3 \leq 5, x^1 + x^2 \geq 1, x^1 \geq 0, x^2 \geq 0, x^3 \geq 0\}$ , and the extreme points of  $S$  are as follows:

$$a_1 = (2.5, 0, 0), a_2 = (1, 0, 0), a_3 = (0, 1, 0), a_4 = (0, 5, 0), a_5 = (0, 1, 4), a_6 = (1, 0, 3).$$

By [7], one can obtain:

$$\begin{aligned} IR &= \{(x^1, x^2, x^3) : 0 \leq x^1 \leq 1, x^2 = 1 - x^1, x^3 = 0\} \\ &\cup \{(x^1, x^2, x^3) : 1 \leq x^1 \leq \frac{5}{2}, x^2 = 0, x^3 = 0\}. \end{aligned}$$

The sets  $S$  and  $IR$  are drawn in Figure 3. The inducible region  $IR$  is denoted by the hatched lines.



**Figure 3.** Constraint region and  $IR$  of Example 2.

Note that in this example,  $S$  and  $IR$  are non-empty and  $S$  is compact. Also, one can obtain:

$$P_1(x^1, x^3) = \{(x^1, x^2, x^3) \in S : x^2 = 1 - x^1, 0 \leq x^1 \leq 1\} \\ \cup \{(x^1, x^2, x^3) \in S : x^2 = 0, 1 \leq x^1 \leq \frac{5}{2} - \frac{1}{2}x^3\}, \\ P_2(x^1, x^2) = \{(x^1, x^2, x^3) \in S : x^3 = 0\}.$$

Hence,  $P_1(x^1, x^3) \neq \emptyset$ ,  $P_2(x^1, x^2) \neq \emptyset$ , and  $P_1(x^1, x^3)$  and  $P_2(x^1, x^2)$  are point-to-point maps with respect to  $(x^1, x^3)$  and  $(x^1, x^2)$ , respectively. Therefore, the Assumptions (1) and (2) hold, and we can solve this problem by the proposed algorithm. Applying the proposed algorithm to this example, we have:

**Step 1.** The MOLP $i$  problems, for  $i = 1, 2$  are constructed as follows, respectively:

$$\min\{(x^1, x^3, -x^1 - x^3, 2x^2) : (x^1, x^2, x^3) \in S\}, \\ \min\{(x^1, x^2, -x^1 - x^2, x^3) : (x^1, x^2, x^3) \in S\}.$$

**Step 2.** One can obtain the sets  $E_i$ ,  $i = 1, 2$ , using the approach described in [20]:

$$E_1 = \gamma(a_1, a_2, a_6) \cup \gamma(a_2, a_3, a_5, a_6), \quad E_2 = \gamma(a_1, a_2, a_3, a_4).$$

The sets  $E_i$ ,  $i = 1, 2$  are drawn in Figure 4 (a) and (b) by the hatched regions and gray area, respectively.

**Step 3.** Set  $E = E_1 \cap E_2$ . So, we obtain  $E = \gamma(a_1, a_2) \cup \gamma(a_2, a_3)$ . Here,  $E$  coincides with  $IR$  which is shown in Figure 3 by the hatched lines.


**Step 4.** Solve the following LP:

$$\min\{3x^1 + x^2 - x^3 : (x^1, x^2, x^3) \in S\}.$$

The optimal solution is  $x^* = (0, 1, 4)$ . Because  $x^* \notin E$ , go to step 5.

**Step 5.** Solve the following problem:

$$\min\{3x^1 + x^2 - x^3 : (x^1, x^2, x^3) \in E\}.$$



**Figure 4.** Constraint region,  $E_1$  and  $E_2$  of Example 2.

This problem is a non-convex optimization problem. According to Remark 2, we just consider the extreme points of  $E$ . Then, we conclude that the point  $x^* = (0, 1, 0)$  is an optimal solution. Also, one can solve this example using the multi-parametric approach and obtain  $x^* = (0, 1, 0)$ . The optimal solution is equal with the obtained optimal solution using the proposed algorithm.

**Example 3.** Consider the following LBLMFP problem:

$$\begin{aligned} \min_{x^1} F(x^1, x^2, x^3) &= -x^1 - 4x^2, \\ \min_{x^2} f_1(x^2, x^3) &= 3x^2 - 2x^3, \\ \text{s.t. } x^1 + x^2 &\leq 2, \quad x^1 \geq 0, \quad x^2 \geq 0, \\ \min_{x^3} f_2(x^2, x^3) &= -x^2 + 4x^3, \\ \text{s.t. } x^1 &\geq 0, \quad x^2 \geq 0, \quad 2 \leq x^3 \leq 4. \end{aligned}$$

In this problem,  $S = \{(x^1, x^2, x^3) : x^1 + x^2 \leq 2, x^1 \geq 0, x^2 \geq 0, 2 \leq x^3 \leq 4\}$ , and the extreme points of  $S$  are as follows:


$$a_1 = (2, 0, 2), \quad a_2 = (0, 0, 2), \quad a_3 = (0, 2, 2), \quad a_4 = (0, 2, 4), \quad a_5 = (0, 0, 4), \quad a_6 = (2, 0, 4).$$

By [7], one can obtain:

$$IR = \{(x^1, x^2, x^3) : 0 \leq x^1 \leq 2, x^2 = 0, x^3 = 2\}.$$

The Figure 5 displays constraint region  $S$  and inducible region  $IR$ . The inducible region is denoted by the hatched line. Note that in this example,  $S$  and  $IR$  are non-empty and  $S$  is compact. Also, one can obtain:

$$\begin{aligned} P_1(x^1, x^3) &= \{(x^1, x^2, x^3) \in S : x^2 = 0\}, \\ P_2(x^1, x^2) &= \{(x^1, x^2, x^3) \in S : x^3 = 2\}. \end{aligned}$$



**Figure 5.** Constraint region and  $IR$  of Example 3.

Hence,  $P_1(x^1, x^3) \neq \emptyset$ ,  $P_2(x^1, x^2) \neq \emptyset$ , and  $P_1(x^1, x^3)$  and  $P_2(x^1, x^2)$  are point-to-point maps with respect to  $(x^1, x^3)$  and  $(x^1, x^2)$ , respectively. Therefore, the Assumptions (1) and (2) hold, and we can solve this problem by the proposed algorithm. Applying the proposed algorithm to this example, we have:


**Step 1.** The MOLP $i$  problems, for  $i = 1, 2$  are constructed as follows, respectively:

$$\begin{aligned} & \min\{(x^1, x^3, -x^1 - x^3, 3x^2) : (x^1, x^2, x^3) \in S\}, \\ & \min\{(x^1, x^2, -x^1 - x^2, 4x^3) : (x^1, x^2, x^3) \in S\}. \end{aligned}$$

**Step 2.** One can obtain the sets  $E_i$ ,  $i = 1, 2$ , using the approach described in [20]:

$$E_1 = \gamma(a_1, a_2, a_5, a_6), \quad E_2 = \gamma(a_1, a_2, a_3).$$

The sets  $E_i$ ,  $i = 1, 2$  are drawn in Figure 6 (a) and (b) by the gray areas, respectively.



**Figure 6.** Constraint region,  $E_1$  and  $E_2$  of Example 3.

**Step 3.** Set  $E = E_1 \cap E_2$ . So, we obtain  $E = \gamma(a_1, a_2) = IR$ . It is shown in Figure 5 by the hatched line.

**Step 4.** Solve the following LP:

$$\min\{-x^1 - 4x^2 : (x^1, x^2, x^3) \in S\}.$$

The extreme optimal solutions are  $x^* = (0, 2, 2)$  and  $y^* = (0, 2, 4)$ . Because  $x^*, y^* \notin E$ , go to Step 5.

**Step 5.** Solve the following problem:

$$\min\{-x^1 - 4x^2 : (x^1, x^2, x^3) \in E\}.$$

Since  $E$  is convex, this problem is the linear programming problem. An optimal solution is  $x^* = (2, 0, 2)$ . Also, one can solve this example using the multi-parametric approach and obtain  $x^* = (2, 0, 2)$ . The obtained optimal solution is equal with the obtained optimal solution using the proposed algorithm.

## 6. Conclusions

In this paper, we have presented a relation between a class of LBLMFP problems and multi-criteria optimization for the first time. We have shown how to construct two MOLP problems so that the extreme points of the set of efficient solutions for both problems are the same as those of the set of feasible solutions for the LBLMFP problem. It is proved that solving the given LBLMFP problem is equivalent to optimizing the leader objective function over a certain efficient set. Based on this result, we proposed an algorithm for solving the LBLMFP problem, and we also showed that it can be simplified in some special cases. Further studies are being conducted to improve the performance of the proposed algorithm and to extend to other classes of LBLMFP problems.

## Conflict of interest

All authors declare no conflicts of interest in this paper.

## References

1. J. F. Bard, *Practical Bilevel Optimization*, Kluwer Academic, Dordrecht, 1998.
2. M. S. Bazaraa, J. J. Jarvis, H. D. Sherali, *Linear Programming and Network Flows*, 2nd ed., Wiley, New York, 1977.
3. M. S. Bazaraa, H. D. Sherali, C. M. Shetty, *Nonlinear Programming, Theory and Algorithms*, 3rd ed., Wiley, New York, 2006.
4. H. P. Benson, *Optimization over the efficient set*, J. Math. Anal. Appl., **98** (1984), 562–580.
5. S. Dempe, *Foundations of Bilevel Programming*, Kluwer Academic, Dordrecht 2003.
6. M. Ehrgott, *Multicriteria Optimization*, 2nd ed., Berlin, Germany: Springer-Verlag, 2005.
7. N. P. Faísca, P. M. Saraiva, B. Rustem, et al. *A multi-parametric programming approach for multilevel hierarchical and decentralized optimisation problems*, Computer Management Sciences, **6** (2009), 377–397.
8. J. Fülöp, *On the equivalency between a linear bilevel programming problem and linear optimization over the efficient set*, Technical Report WP 93-1, Laboratory of Operations Research and Decision Systems, Computer and Automation Institute, Hungarian Academy of Sciences, 1993.

9. J. Glackin, J. G. Ecker, M. Kupferschmid, *Solving bilevel linear programs using multiple objective linear programming*, J. Optimiz. Theory App., **140** (2009), 197–212.
10. J. Han, G. Zhang, Y. Hu, et al. *A solution to bi/tri-level programming problems using particle swarm optimization*, Inform. Sciences, **370** (2016), 519–537.
11. R. Horst, N. V. Thoai, Y. Yamamoto, et al. *On optimization over the efficient set in linear multicriteria programming*, J. Optimiz. Theory App., **134** (2007), 433–443.
12. J. Jorge, *A bilinear algorithm for optimizing a linear function over the efficient set of multiple objective linear programming problem*, J. Global Optim., **31** (2005), 1–16.
13. J. Lu, C. Shi, G. Zhang, et al. *Model and extended Kuhn-Tucker approach for bilevel multi-follower decision making in a referential-uncooperative situation*, J. Global Optim., **38** (2007), 597–608.
14. W. Ma, M. Wang, X. Zhu, *Improved particle swarm optimization based approach for bilevel programming problem-an application on supply chain model*, Int. J. Mach. Learn. Cyb., **5** (2014), 281–292.
15. B. Metev, *Multiobjective optimization methods help to minimize a function over the efficient set*, Cybernetics and Information Technologies, **7** (2007), 22–28.
16. C. Miao, G. Du, R. J. Jiao, et al. *Coordinated optimisation of platform-driven product line planning by bilevel programming*, Int. J. Prod. Res., **55** (2017), 3808–3831.
17. C. O. Pieume, L. P. Fotso, P. Siarry, *An approach for finding efficient points in multiobjective linear programming*, Journal of Information and Optimization Sciences, **29** (2008), 203–216.
18. C. O. Pieume, L. P. Fotso, P. Siarry, *Solving bilevel programming problems with multicriteria optimization techniques*, Opsearch., **46** (2009), 169–183.
19. A. S. Safaei, S. Farsad, M. M. Paydar, *Robust bi-level optimization of relief logistics operations*, Appl. Math. Model., **56** (2018), 359–380.
20. S. Sayin, *An algorithm based on facial decomposition for finding the efficient set in multiple objective linear programming*, Oper. Res. Lett., **19** (1996), 87–94.
21. C. Shi, G. Zhang, J. Lu, *A Kth-best Approach for Linear Bilevel Multi-follower Programming*, J. Global Optim., **33** (2005), 563–578.
22. R. Steuer, *Multiple Criteria Optimization: Theory, Computation, and Application*, Wiley, New York, 1986.
23. G. Zhang, J. Lu, Y. Gao, *Multi-Level Decision Making: Models, Methods and Applications*, Springer-Verlag Berlin AN, 2016.



AIMS Press

©2019 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)