



Research article

Approximation of solutions for nonlinear functional integral equations

Lakshmi Narayan Mishra^{1,*}, Vijai Kumar Pathak¹ and Dumitru Baleanu^{2,3,4}

¹ Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology, Vellore 632014, Tamil Nadu, India

² Department of Mathematics, Faculty of Arts and Sciences, Cankaya University, Ankara, Turkey

³ Institute of Space Sciences, Magurele-Bucharest, Romania

⁴ Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan

* **Correspondence:** Email: lakshminarayanmishra04@gmail.com, l.n.mishra@yahoo.co.in;
Tel: +919838375431; Fax: +919838375431.

Abstract: In this article, we consider a class of nonlinear functional integral equations, motivated by an equation that offers increasing evidence to the extant literature through replication studies. We investigate the existence of solution for nonlinear functional integral equations on Banach space $C[0, 1]$. We use the technique of the generalized Darbo's fixed-point theorem associated with the measure of noncompactness (MNC) to prove our existence result. Also, we have given two examples of the applicability of established existence result in the theory of functional integral equations. Further, we construct an efficient iterative algorithm to compute the solution of the first example, by employing the modified homotopy perturbation (MHP) method associated with Adomian decomposition. Moreover, the condition of convergence and an upper bound of errors are presented.

Keywords: measure of noncompactness; nonlinear functional integral equation; fixed point theorem; modified homotopy perturbation

Mathematics Subject Classification : 45G10, 45M99, 47H08, 47H10

1. Introduction

Integral equations play a notable role in applied mathematics. The practical importance of nonlinear integral equations is increasingly evident from studies that incorporate the same in distinct areas of knowledge that include biology, traffic theory, the theory of optimal control, economics, acoustic scattering, etc. [1,2,9,39]. Precisely, extensive studies on these equations are focused on their solutions by employing the MNC [3, 29, 36, 47]. Besides, other concepts like quasi linearization [35] and

pseudo-spectral methods [52] have also been used in similar studies. In such studies, the existence of solutions is proved with the theory of fixed points. Verifying the existence of solutions, their behavioral properties are also extensively studied. For instance, Hu et al. [20] discussed the global attractivity and asymptotic stability of the solutions, whereas Wang et al. [51] discussed the local attractivity and local stability, and Aghajani et al. [4] and Alvarez et al. [5] gave globally and uniformly locally attractive solutions. Notably, Banaś et al. [6, 8, 10] and Dhage et al. [12, 13, 16, 17] also discussed the attractivity of solutions. Xu et al. [53] renders radially symmetric solutions and their asymptotic estimates. Furthermore, systems of equations have been studied and numerical methods to find solutions have also been proposed [19, 22, 23, 25, 38, 40, 48, 49].

In recent times, the fixed point theory (FPT) is applicable in various scientific fields suggested by Banach [11, 21, 26, 37, 50]. Also, FPT can be applied to seeking solutions of functional integral equations. Functional integral equations of variety of forms chair as a extraordinary and prestigious branch of non-linear analysis and seek various invocations in demonstrating numerous real-life together with real-world problems (cf. [14, 27, 28, 30–34, 41]).

Recently, several research articles have been published in connection:

In 2020, El-Sayed and Ebeada [18] have studied the solvability of self-reference functional and quadratic functional integral equations:

$$x(t) = f\left(t, \int_0^t g(s, x(x(s)))ds\right),$$

$$\text{and } x(t) = f\left(t, \int_0^t f_1(s, x(x(s)))ds \int_0^t f_2(s, x(x(s)))ds\right), \text{ respectively,}$$

where $x \in C[0, T]$, $t \in [0, T]$, and g, f_1, f_2 satisfy Carathéodory condition. To realize the existence of a solution to those integral equations, they used Schauder's fixed point theorem in the Banach space $C[0, T]$.

In 2020, Deep et al. [15] established the existence of solutions of some non-linear functional integral equations in Banach algebra with applications:

$$y(t) = \left(f(t, y(t), y(\theta(t))) + F\left(t, \int_0^t r(t, s, y(\theta(s)))ds, \int_0^t u(t, s, y(a(s)))ds, y(d(t))\right) \right)$$

$$\times L\left(t, \int_0^b p(t, s, y(c(s)))ds, \int_0^b q(t, s, y(\chi(s)))ds, y(\eta(t))\right),$$

for $t \in [0, b]$. Existence result is obtained through the techniques of MNC and Darbo's fixed point theorem in $[0, b]$.

In 2021, Rabbani et al. [46] established some generalized non-linear functional integral equations of two variables via measures of noncompactness and numerical methods to solve it,

$$y(\zeta) = \left(f(\zeta) + A(\zeta, y(\zeta), Y(\alpha(\zeta))) + P\left(\zeta, y(\zeta), \int_0^\zeta F(\zeta, r, y(\beta(r)))dr, y(\gamma(\zeta))\right) \right)$$

$$\times \left(U(\zeta, y(\zeta), Y(\theta(\zeta))) + Q\left(\zeta, y(\zeta), \int_0^b G(\zeta, r, y(\delta(r))), y(\phi(\zeta))\right) \right),$$

where $\zeta \in [0, b]$. To realize the existence of the solution of those integral equations, they have used the concept of MNC and Petryshyn fixed point theorem for the operators in a Banach algebra $C([0, b] \times$

$[0, b], \mathbb{R}$), for $b > 0$ in the form of two operators. They have also discussed an iterative algorithm which was constructed by modified homotopy perturbation method and Adomian polynomials to compute the solution of the example.

In 2022, Karmakar et al. [24] have studied the existence of solutions to non-linear quadratic integral equations via measure of noncompactness:

$$x(t) = g(t, x(t)) + \lambda \int_0^t \mu_1(t, s) \zeta_1(s, x(s)) ds + \int_0^t \mu_2(t, s) \zeta_2(s, x(s)) ds,$$

for $t > 0$, where $g, \mu_1, \mu_2, \zeta_1, \zeta_2$ are real valued continuous functions defined on $\mathbb{R}^+ \times \mathbb{R}$ and λ is a positive constant.

In our work, we study an existence result for the solution of the following nonlinear functional integral equation:

$$\varpi(\varrho) = f\left(\varrho, \varpi(\varrho), g(\varrho, \varpi(\varrho)) \int_0^1 v(\varrho, \eta, \varpi(\eta)) d\eta, \varpi(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta)) d\eta\right), \quad (1.1)$$

where $\varrho \in I = [0, 1]$. In this work, our main work aims to obtain the existence result of Eq (1.1) on Banach space $C[0, 1]$ by applying the technique of the generalized Darbo's fixed-point theorem associated with the MNC, and also, work to obtain the analytic solution of it by applying the semi-analytic method. Now, we describe the importance of why we study Eq (1.1) and what is the perfection of our findings. The first one is that the conditions estimated in several research articles will be analyzed and the second one is that this manuscript affiliate the relevant work in this area. The third one is the bounded condition implies that the "sublinear condition" that has been identified in various literature works does not have a relevant appearance here. Our findings generalized, extended, and complement several results existing in the literature.

The estimate of our work is organized as follows: In Section 2, some notations, definitions and auxiliary facts are given. In Section 3, we prove the existence of solution by using the generalized Darbo fixed point theorem associated with the MNC on $C[0, 1]$. Also, we present two examples to illustrate our theorem. In Section 4, we state an algorithm to find the solution by using MHP and Adomian decomposition method. Correspondingly, we apply the algorithm to one of the examples for finding an approximate solution and tabulate the errors. Also, we show the graphs of $v(\varrho)$ and $\varpi(\varrho)$. In Section 5, we analyze the errors and provide an upper bound of the errors. Our conclusion is presented in Section 6.

2. Preliminaries

In this section, we organize some notations, definitions and auxiliary facts which we need throughout the paper.

Let A be a nonempty subset of a Banach space X . We use \bar{A} and $Conv A$ to denote the closure and convex closure of A respectively. Also, we use P_X to denote the class of nonempty bounded subsets of X and Q_X to denote the subclass of relatively compact sets of P_X . Let $C(I)$ denotes the set of all continuous real valued functions on $[0, 1]$, a classical Banach space.

Definition 2.1. [7] A function $\mu : P_X \rightarrow \mathbb{R}^+$ is called a MNC on X if the following conditions are satisfied:

- (i) The kernel of the function, $\ker \mu = \{A \in P_X : \mu(A) = 0\} \neq \emptyset$ & $\ker \mu \subseteq Q_X$.
- (ii) $A \subseteq B \Rightarrow \mu(A) \leq \mu(B)$, $\forall A, B \in P_X$.
- (iii) $\mu(\bar{A}) = \mu(\text{Conv } A) = \mu(A)$, $\forall A \in P_X$.
- (iv) $\mu(\lambda A + (1 - \lambda)B) \leq \lambda\mu(A) + (1 - \lambda)\mu(B)$, for $\lambda \in [0, 1]$.
- (v) If $\{A_n\}_{n=1}^\infty$ is a sequence of closed sets in P_X such that $A_{n+1} \subseteq A_n$, $\forall n \in \mathbb{N}$ and $\mu(A_n) \rightarrow 0$, then $\bigcap_{n=1}^\infty A_n = A_\infty \neq \emptyset$.

Definition 2.2. Define $\Psi = \{\psi' : \mathbb{R}^+ \rightarrow \mathbb{R}^+\}$ such that each ψ' satisfies the following:

- (i) ψ' is an upper semi-continuous function from the right.
- (ii) $\psi'(\varrho) < \varrho$, $\forall \varrho \in [0, 1]$.

Definition 2.3. We recall the definition of MNC in $C(I)$ as defined in [8]. Let $A \subseteq C(I)$ such that A is nonempty, bounded, and let $\varpi \in A$ together with $\epsilon \geq 0$, then the modulus of continuity of ϖ in I is defined as

$$\omega(\varpi, \epsilon) = \sup_{\varrho, \eta \in I} \{|\varpi(\varrho) - \varpi(\eta)| : |\varrho - \eta| \leq \epsilon\}.$$

$$\text{Let } \omega(A, \epsilon) = \sup_{\varpi \in A} \omega(\varpi, \epsilon) \quad \text{and} \quad \omega_0(A) = \lim_{\epsilon \rightarrow 0} \omega(A, \epsilon).$$

$$\text{Define } i(\varpi) = \sup_{\varrho, \eta \in I} \{|\varpi(\varrho) - \varpi(\eta)| - [|\varpi(\varrho) - \varpi(\eta)|] : \eta \leq \varrho\} \quad \text{and} \quad i(A) = \sup_{\varpi \in A} i(\varpi).$$

Now, MNC is defined as

$$\mu(A) = \omega_0(A) + i(A).$$

Theorem 2.1. [16] Let A be a nonempty, closed, bounded together with convex subset of a Banach space, and let $T : A \rightarrow A$ be a continuous function satisfying

$$\mu(TB) \leq \psi'(\mu(B)), \quad \forall B \subseteq A (B \neq \emptyset), \quad (2.1)$$

for some MNC μ and some $\psi' \in \Psi$. Then, there exist at least one fixed point in T .

This theorem is known as the generalized Darbo fixed point theorem.

3. Main result

In this section, we prove the nonlinear equation (1.1) has a solution in the Banach space $C(I)$, with the supremum norm.

The nonlinear equation (1.1) is studied with the following assumptions:

- (A1) The function $f : I \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous such that $f : I \times \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$. Also, there exists a function $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $\psi(0) = 0$ and $\psi(\varrho) < \frac{\varrho}{2}$ together with $\psi(\varrho) + \psi(\eta) \leq \psi(\varrho + \eta)$ satisfying

$$|f(\varrho, u_1, v_1, w_1) - f(\varrho, u_2, v_2, w_2)| \leq \psi(|u_1 - u_2|) + |v_1 - v_2| + |w_1 - w_2|. \quad (3.1)$$

For $u, v, w \in \mathbb{R}^+$, $\varrho \mapsto f(\varrho, u, v, w)$ is increasing on I and for $\varrho \in I$ and $u \in \mathbb{R}^+$, $v \mapsto f(\varrho, u, v, w)$ and $w \mapsto f(\varrho, u, v, w)$ are increasing on I and for some $N > 0$, $f(\varrho, 0, 0, 0) \leq N$.

(A2) The function $g : I \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and there exists $\kappa', \kappa \geq 0$ such that

$$|g(\varrho, 0)| \leq \kappa' \quad \text{and} \quad |g(\varrho, y_1) - g(\varrho, y_2)| \leq \kappa|y_1 - y_2|. \quad (3.2)$$

(A3) The functions $u : I \times I \times \mathbb{R} \rightarrow \mathbb{R}$ and $v : I \times I \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous such that $u : I \times I \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and $v : I \times I \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$. For arbitrarily fixed $\eta \in I$ ($\Rightarrow \varpi(\eta)$ is also fixed), $\varrho \mapsto u(\varrho, \eta, \varpi(\eta))$ and $\varrho \mapsto v(\varrho, \eta, \varpi(\eta))$ are increasing in I . Also, there exists a constant $l \in [0, \frac{1}{4})$ such that

$$\int_0^1 v(\varrho, \eta, \varpi(\eta)) d\eta \leq l \quad \text{and} \quad \int_0^\varrho u(\varrho, \eta, \varpi(\eta)) d\eta \leq l. \quad (3.3)$$

(A4) There exists $r_0 > 0$ such that

$$\psi(r_0) + (r_0(\kappa + 1) + \kappa')l + N \leq r_0. \quad (3.4)$$

(A5) The constant $\kappa l < 1/4$.

Theorem 3.1. *Under the above assumptions, the nonlinear equation (1.1) has at least one solution in $C(I)$.*

Proof. Let $T : C(I) \rightarrow C(I)$ be an operator defined as

$$(T\varpi)(\varrho) = f\left(\varrho, \varpi(\varrho), g(\varrho, \varpi(\varrho)) \int_0^1 v(\varrho, \eta, \varpi(\eta)) d\eta, \varpi(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta)) d\eta\right).$$

The operator T having a fixed point in $C(I)$ is equivalent to the nonlinear equation (1.1) having a solution in $C(I)$. Hence, we prove T has a fixed point by using Theorem 2.1.

By applying system of Eq (1.1) and imposed postulates (A1–A5), we estimate for every $\varrho \in I$ such that

$$\begin{aligned} & |(T\varpi)(\varrho)| \\ & \leq \left| f\left(\varrho, \varpi(\varrho), g(\varrho, \varpi(\varrho)) \int_0^1 v(\varrho, \eta, \varpi(\eta)) d\eta, \varpi(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta)) d\eta\right) - f(\varrho, 0, 0, 0) \right| + |f(\varrho, 0, 0, 0)| \\ & \leq \psi(|\varpi(\varrho)|) + |g(\varrho, \varpi(\varrho))| \int_0^1 |v(\varrho, \eta, \varpi(\eta))| d\eta + |\varpi(\varrho)| \int_0^\varrho |u(\varrho, \eta, \varpi(\eta))| d\eta + |f(\varrho, 0, 0, 0)| \\ & \leq \psi(|\varpi(\varrho)|) + (|g(\varrho, \varpi(\varrho))| + |\varpi(\varrho)|)l + |f(\varrho, 0, 0, 0)| \\ & \leq \psi(|\varpi(\varrho)|) + (|g(\varrho, \varpi(\varrho)) - g(\varrho, 0)| + |g(\varrho, 0)| + |\varpi(\varrho)|)l + |f(\varrho, 0, 0, 0)| \\ & \leq \psi(|\varpi(\varrho)|) + (\kappa|\varpi(\varrho)| + \kappa' + |\varpi(\varrho)|)l + |f(\varrho, 0, 0, 0)| \\ & \leq \psi(\|\varpi\|) + (\|\varpi\|(\kappa + 1) + \kappa')l + N. \end{aligned}$$

Therefore,

$$\|T\varpi\| \leq \psi(\|\varpi\|) + (\|\varpi\|(\kappa + 1) + \kappa')l + N.$$

For $r_0 > 0$ such that $\|\varpi\| \leq r_0$, by assumption (A4), $\|T\varpi\| \leq r_0$, i.e. T maps B_{r_0} into itself. Now, we prove T is continuous. Let $\{\varpi_n\}_{n=1}^\infty$ be a sequence in B_{r_0} such that $\varpi_n \rightarrow \varpi$, we obtain

$$\begin{aligned}
& |(T\varpi_n)(\varrho) - (T\varpi)(\varrho)| \\
&= \left| f\left(\varrho, \varpi_n(\varrho), g(\varrho, \varpi_n(\varrho)) \int_0^1 v(\varrho, \eta, \varpi_n(\eta))d\eta, \varpi_n(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi_n(\eta))d\eta\right) \right. \\
&\quad \left. - f\left(\varrho, \varpi(\varrho), g(\varrho, \varpi(\varrho)) \int_0^1 v(\varrho, \eta, \varpi(\eta))d\eta, \varpi(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta))d\eta\right) \right| \\
&\leq \psi(|\varpi_n(\varrho) - \varpi(\varrho)|) + \left| g(\varrho, \varpi_n(\varrho)) \int_0^1 v(\varrho, \eta, \varpi_n(\eta))d\eta - g(\varrho, \varpi(\varrho)) \int_0^1 v(\varrho, \eta, \varpi(\eta))d\eta \right| \\
&\quad + \left| \varpi_n(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi_n(\eta))d\eta - \varpi(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta))d\eta \right|.
\end{aligned}$$

Let us now consider

$$\begin{aligned}
& \left| g(\varrho, \varpi_n(\varrho)) \int_0^1 v(\varrho, \eta, \varpi_n(\eta))d\eta - g(\varrho, \varpi(\varrho)) \int_0^1 v(\varrho, \eta, \varpi(\eta))d\eta \right| \\
&\leq |g(\varrho, \varpi_n(\varrho))| \int_0^1 |v(\varrho, \eta, \varpi_n(\eta)) - v(\varrho, \eta, \varpi(\eta))|d\eta \\
&\quad + |g(\varrho, \varpi_n(\varrho)) - g(\varrho, \varpi(\varrho))| \int_0^1 |v(\varrho, \eta, \varpi(\eta))|d\eta \\
&\leq |g(\varrho, \varpi_n(\varrho))|V_{r_0}(\epsilon) + \kappa l|\varpi_n(\varrho) - \varpi(\varrho)| \\
&\leq (\kappa\|\varpi_n\| + \kappa')V_{r_0}(\epsilon) + \kappa l\|\varpi_n - \varpi\|,
\end{aligned}$$

and

$$\begin{aligned}
& \left| \varpi_n(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi_n(\eta))d\eta - \varpi(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta))d\eta \right| \\
&\leq \left| \varpi_n(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi_n(\eta))d\eta - \varpi_n(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta))d\eta \right| \\
&\quad + \left| \varpi_n(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta))d\eta - \varpi(\varrho) \int_0^\varrho u(\varrho, \eta, \varpi(\eta))d\eta \right| \\
&\leq \|\varpi_n\| U_{r_0}(\epsilon) + l\|\varpi_n - \varpi\|,
\end{aligned}$$

where

$$V_{r_0}(\epsilon) = \sup_{\varrho, \eta \in I} \{|v(\varrho, \eta, \varpi(\eta)) - v(\varrho, \eta, y(\eta))| : \varpi, y \in B_{r_0} \& |\varpi - y| \leq \epsilon\},$$

and

$$U_{r_0}(\epsilon) = \sup_{\varrho, \eta \in I} \{|u(\varrho, \eta, \varpi(\eta)) - u(\varrho, \eta, y(\eta))| : \varpi, y \in B_{r_0} \& |\varpi - y| \leq \epsilon\}.$$

As $\epsilon \rightarrow 0$, $V_{r_0} \rightarrow 0$ and $U_{r_0} \rightarrow 0$. We are enabled to obtain

$$\|T\varpi_n - T\varpi\| \leq \psi(\|\varpi_n - \varpi\|) + l(\kappa + 1)\|\varpi_n - \varpi\|,$$

i.e., $T\varpi_n \rightarrow T\varpi$. Hence, we have proved that T is continuous in B_{r_0} . Clearly $T(B_{r_0}^+) \subseteq B_{r_0}^+$, where $B_{r_0}^+ = \{\varpi \in B_{r_0} : \varpi(\varrho) \geq 0, \quad \forall \varrho \in I\}$.

Further, we suppose $A \subseteq B_{r_0}$ such that A is nonempty and $\varpi \in A$. Also, let $\epsilon > 0$ and $\varrho_1, \varrho_2 \in I$ such that $|\varrho_1 - \varrho_2| \leq \epsilon$. Without loss of generality, let us also assume that $\varrho_2 \geq \varrho_1$, we estimate

$$\begin{aligned}
& |(T\varpi)(\varrho_2) - (T\varpi)(\varrho_1)| \\
& \leq \left| f(\varrho_2, \varpi(\varrho_2), g(\varrho_2, \varpi(\varrho_2))) \int_0^1 v(\varrho_2, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_2) \int_0^{\varrho_2} u(\varrho_2, \eta, \varpi(\eta)) d\eta \right. \\
& \quad \left. - f(\varrho_2, \varpi(\varrho_1), g(\varrho_2, \varpi(\varrho_2))) \int_0^1 v(\varrho_2, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_2) \int_0^{\varrho_2} u(\varrho_2, \eta, \varpi(\eta)) d\eta \right| \\
& \quad + \left| f(\varrho_2, \varpi(\varrho_1), g(\varrho_2, \varpi(\varrho_2))) \int_0^1 v(\varrho_2, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_2) \int_0^{\varrho_2} u(\varrho_2, \eta, \varpi(\eta)) d\eta \right. \\
& \quad \left. - f(\varrho_1, \varpi(\varrho_1), g(\varrho_2, \varpi(\varrho_2))) \int_0^1 v(\varrho_2, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_2) \int_0^{\varrho_2} u(\varrho_2, \eta, \varpi(\eta)) d\eta \right| \\
& \quad + \left| f(\varrho_1, \varpi(\varrho_1), g(\varrho_2, \varpi(\varrho_2))) \int_0^1 v(\varrho_2, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_2) \int_0^{\varrho_2} u(\varrho_2, \eta, \varpi(\eta)) d\eta \right. \\
& \quad \left. - f(\varrho_1, \varpi(\varrho_1), g(\varrho_2, \varpi(\varrho_2))) \int_0^1 v(\varrho_1, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_2) \int_0^{\varrho_1} u(\varrho_2, \eta, \varpi(\eta)) d\eta \right| \\
& \quad + \left| f(\varrho_1, \varpi(\varrho_1), g(\varrho_2, \varpi(\varrho_2))) \int_0^1 v(\varrho_1, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_2) \int_0^{\varrho_1} u(\varrho_2, \eta, \varpi(\eta)) d\eta \right. \\
& \quad \left. - f(\varrho_1, \varpi(\varrho_1), g(\varrho_1, \varpi(\varrho_1))) \int_0^1 v(\varrho_1, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_1) \int_0^{\varrho_1} u(\varrho_1, \eta, \varpi(\eta)) d\eta \right| \\
& \leq \psi(|\varpi(\varrho_2) - \varpi(\varrho_1)|) + \omega(f, \epsilon) + |g(\varrho_2, \varpi(\varrho_2))| \int_0^1 |v(\varrho_2, \eta, \varpi(\eta)) - v(\varrho_1, \eta, \varpi(\eta))| d\eta \\
& \quad + |\varpi(\varrho_2)| \int_{\varrho_1}^{\varrho_2} |u(\varrho_2, \eta, \varpi(\eta))| d\eta + |g(\varrho_2, \varpi(\varrho_2)) - g(\varrho_1, \varpi(\varrho_1))| \int_0^1 |v(\varrho_1, \eta, \varpi(\eta))| d\eta \\
& \quad + \left| \varpi(\varrho_2) \int_0^{\varrho_1} u(\varrho_2, \eta, \varpi(\eta)) d\eta - \varpi(\varrho_1) \int_0^{\varrho_1} u(\varrho_1, \eta, \varpi(\eta)) d\eta \right| \\
& \leq \psi(|\varpi(\varrho_2) - \varpi(\varrho_1)|) + \omega(f, \epsilon) + |g(\varrho_2, \varpi(\varrho_2))| \omega(v, \epsilon) + |\varpi(\varrho_2)| l \epsilon \\
& \quad + |g(\varrho_2, \varpi(\varrho_2)) - g(\varrho_1, \varpi(\varrho_1))| \int_0^1 |v(\varrho_1, \eta, \varpi(\eta))| d\eta \\
& \quad + \left| \varpi(\varrho_2) \int_0^{\varrho_1} u(\varrho_2, \eta, \varpi(\eta)) d\eta - \varpi(\varrho_1) \int_0^{\varrho_1} u(\varrho_1, \eta, \varpi(\eta)) d\eta \right|,
\end{aligned}$$

wherein,

$$\begin{aligned}
& |g(\varrho_2, \varpi(\varrho_2)) - g(\varrho_1, \varpi(\varrho_1))| \int_0^1 |v(\varrho_1, \eta, \varpi(\eta))| d\eta \\
& \leq \{|g(\varrho_2, \varpi(\varrho_2)) - g(\varrho_1, \varpi(\varrho_2))| + |g(\varrho_1, \varpi(\varrho_2)) - g(\varrho_1, \varpi(\varrho_1))|\} l \\
& \leq l\omega(g, \epsilon) + \kappa l |\varpi(\varrho_2) - \varpi(\varrho_1)|,
\end{aligned}$$

and

$$\begin{aligned}
& \left| \varpi(\varrho_2) \int_0^{\varrho_1} u(\varrho_2, \eta, \varpi(\eta)) d\eta - \varpi(\varrho_1) \int_0^{\varrho_1} u(\varrho_1, \eta, \varpi(\eta)) d\eta \right| \\
& \leq \left| \varpi(\varrho_2) \int_0^{\varrho_1} u(\varrho_2, \eta, \varpi(\eta)) d\eta - \varpi(\varrho_2) \int_0^{\varrho_1} u(\varrho_1, \eta, \varpi(\eta)) d\eta \right| \\
& \quad + \left| \varpi(\varrho_2) \int_0^{\varrho_1} u(\varrho_1, \eta, \varpi(\eta)) d\eta - \varpi(\varrho_1) \int_0^{\varrho_1} u(\varrho_1, \eta, \varpi(\eta)) d\eta \right| \\
& \leq |\varpi(\varrho_2)| \int_0^{\varrho_1} |u(\varrho_2, \eta, \varpi(\eta)) - u(\varrho_1, \eta, \varpi(\eta))| d\eta + |\varpi(\varrho_2) - \varpi(\varrho_1)| \int_0^{\varrho_1} |u(\varrho_1, \eta, \varpi(\eta))| d\eta \\
& \leq |\varpi(\varrho_2)| \omega(u, \epsilon) + l |\varpi(\varrho_2) - \varpi(\varrho_1)|,
\end{aligned}$$

together with

$$\begin{aligned}
\omega(f, \epsilon) &= \sup_{\varrho_1, \varrho_2 \in I} \{|f(\varrho_1, u, v, w) - f(\varrho_2, u, v, w)| : |\varrho_1 - \varrho_2| \leq \epsilon\}, \\
\omega(u, \epsilon) &= \sup_{\varrho_1, \varrho_2 \in I} \{|u(\varrho_1, \eta, \varpi) - u(\varrho_2, \eta, \varpi)| : |\varrho_1 - \varrho_2| \leq \epsilon\}, \\
\omega(v, \epsilon) &= \sup_{\varrho_1, \varrho_2 \in I} \{|v(\varrho_1, \eta, \varpi) - v(\varrho_2, \eta, \varpi)| : |\varrho_1 - \varrho_2| \leq \epsilon\}, \\
\omega(g, \epsilon) &= \sup_{\varrho_1, \varrho_2 \in I} \{|g(\varrho_2, x) - g(\varrho_1, x)| : |\varrho_2 - \varrho_1| \leq \epsilon\}.
\end{aligned}$$

Then, we can find

$$\begin{aligned}
& |(T\varpi)(\varrho_2) - (T\varpi)(\varrho_1)| \\
& \leq \psi(|\varpi(\varrho_2) - \varpi(\varrho_1)|) + \omega(f, \epsilon) + |g(\varrho_2, \varpi(\varrho_2))| \omega(v, \epsilon) + l\epsilon |\varpi(\varrho_2)| + l\omega(g, \epsilon) \\
& \quad + \kappa l |\varpi(\varrho_2) - \varpi(\varrho_1)| + |\varpi(\varrho_2)| \omega(u, \epsilon) + l |\varpi(\varrho_2) - \varpi(\varrho_1)| \\
& \leq \psi(\omega(\varpi, \epsilon)) + \omega(f, \epsilon) + (\kappa \|\varpi\| + \kappa') \omega(v, \epsilon) + l\epsilon \|\varpi\| + l\omega(g, \epsilon) + (\kappa l + l) \omega(\varpi, \epsilon) + \|\varpi\| \omega(u, \epsilon).
\end{aligned}$$

Now, applying uniform continuity of the function $f(\varrho, u, v, w)$, $u(\varrho, \eta, \varpi)$, $v(\varrho, \eta, \varpi)$ and $g(\varrho, \varpi)$ on the set $I \times [-r_0, r_0] \times [-r_0, r_0] \times [-r_0, r_0]$, $I \times I \times [-r_0, r_0]$, $I \times I \times [-r_0, r_0]$, $I \times [-r_0, r_0]$, respectively. We are enabled to deduce that $\omega(f, \epsilon) \rightarrow 0$, $\omega(u, \epsilon) \rightarrow 0$, $\omega(v, \epsilon) \rightarrow 0$, and $\omega(g, \epsilon) \rightarrow 0$, when $\epsilon \rightarrow 0$. Thus, we can find

$$\begin{aligned}
\omega_0(TA) &\leq \psi(\omega_0(A)) + (\kappa l + l) \omega_0(A) \\
&\leq (\psi + (\kappa + 1)l) (\omega_0(A)).
\end{aligned} \tag{3.5}$$

Suppose $\varpi \in A$, and $\varrho_1, \varrho_2 \in I$, together with $\varrho_1 < \varrho_2$, we estimate that

$$\begin{aligned}
& |(T\varpi)(\varrho_2) - (T\varpi)(\varrho_1)| - |(T\varpi)(\varrho_2) - (T\varpi)(\varrho_1)| \\
& = \left| f\left(\varrho_2, \varpi(\varrho_2), g(\varrho_2, \varpi(\varrho_2)) \int_0^1 v(\varrho_2, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_2) \int_0^{\varrho_2} u(\varrho_2, \eta, \varpi(\eta)) d\eta\right) \right. \\
& \quad \left. - f\left(\varrho_1, \varpi(\varrho_1), g(\varrho_1, \varpi(\varrho_1)) \int_0^1 v(\varrho_1, \eta, \varpi(\eta)) d\eta, \varpi(\varrho_1) \int_0^{\varrho_1} u(\varrho_1, \eta, \varpi(\eta)) d\eta\right) \right|
\end{aligned}$$

Now, we are enabled to write

$$i(T\varpi) \leq (\psi + (\kappa + 1)l)(i(\varpi)).$$

Thus, we can find

$$i(TA) \leq (\psi + (\kappa + 1)l)(i(A)). \quad (3.6)$$

From inequalities (3.5) and (3.6), together with the definition of MNC μ , we can write

$$\mu(TA) = \omega_0(TA) + i(TA) \leq \psi'(\omega_0(A) + i(A)) \leq \psi'(\mu(A)),$$

where $\psi' = \psi + (\kappa + 1)l$, $\psi' \in \Psi$.

Hence, by the generalized Darbo fixed point theorem, T has a fixed point in $C(I)$, i.e., the nonlinear equation (1.1) has at least one solution in $C[0, 1]$. \square

Further, we give two examples with a verification of all the five assumptions of our main theorem.

Example 3.1. In the first example, we consider the following nonlinear integral equation:

$$\varpi(\varrho) = \frac{\varrho}{4(1 + \varpi(\varrho)^2)} \left(1 + \int_0^1 \frac{\varrho\eta}{16(1 + \varpi(\eta)^2)} d\eta \right) + \varpi(\varrho) \int_0^\varrho \frac{\varrho\eta}{16(1 + \varpi(\eta)^2)^2} d\eta. \quad (3.7)$$

Herein, we have $f(\varrho, u, v, w) = \frac{\varrho}{4(1 + u^2)} + v + w$ and $g(\varrho, \varpi(\varrho)) = \frac{\varrho}{4(1 + \varpi^2)}$ satisfying the assumptions (A1) and (A2) respectively such that

$$\begin{aligned} |f(\varrho, u_1, v_1, w_1) - f(\varrho, u_2, v_2, w_2)| &\leq \left| \frac{\varrho}{4(1 + u_1^2)} - \frac{\varrho}{4(1 + u_2^2)} \right| + |v_1 - v_2| + |w_1 - w_2| \\ &\leq \psi(|u_1 - u_2|) + |v_1 - v_2| + |w_1 - w_2|, \end{aligned}$$

where $\psi(\varrho) = \frac{\varrho}{4}$ and

$$\begin{aligned} g(\varrho, 0) &= \frac{\varrho}{4} \leq \frac{1}{4} = \kappa', \\ |g(\varrho, y_1) - g(\varrho, y_2)| &\leq \frac{\varrho}{4} |y_1 - y_2| \leq \frac{1}{4} |y_1 - y_2| = \kappa |y_1 - y_2|. \end{aligned}$$

Also, the functions $u(\varrho, \eta, \varpi) = \frac{\varrho\eta}{16(1 + \varpi^2)}$ and $v(\varrho, \eta, \varpi) = \frac{\varrho\eta}{16(1 + \varpi^2)^2}$ satisfy the assumption (A3) such that $l = \frac{1}{16}$. Thus $\kappa l < \frac{1}{4}$ and for $r_0 = 1$, the assumption (A4) is fulfilled as

$$\psi(r_0) + (r_0(\kappa + 1) + \kappa')l + N = \frac{r_0}{4} + \left(\frac{5r_0}{4} + \frac{1}{4} \right) \frac{1}{16} + \frac{1}{4} \leq r_0.$$

Now, we are enabled to observe that the last inequality admit a positive solution, for $r_0 = 1$. Hence, all assumptions (A1)–(A5) of Theorem 3.1 are fulfilled. Thus, we are enabled to conclude that the nonlinear integral equation (3.7) admits at least one solution in the space $C(I)$.

Example 3.2. Consider the following nonlinear integral equation:

$$\varpi(\varrho) = \frac{\varrho}{1 + \varrho^2} + \frac{\ln \varpi(\varrho)}{1 + \varrho} + \frac{\cos(\varrho \varpi(\varrho))}{2(1 + \varrho)} \int_0^1 \frac{\varrho + \eta}{16(1 + \exp(-\varpi(\eta)))} d\eta + \varpi(\varrho) \int_0^{\varrho} \frac{\arctan \varpi(\eta)}{4(1 + \eta)} d\eta. \quad (3.8)$$

We have $f(\varrho, u, v, w) = \frac{\varrho}{1 + \varrho^2} + \ln u + v + w$ and $g(\varrho, \varpi(\varrho)) = \frac{\cos(\varrho \varpi(\varrho))}{2(1 + \varrho)}$ satisfying the assumptions (A1) and (A2) respectively such that $\psi(\varrho) = \frac{\varrho}{16}$, $\kappa' = \frac{1}{2}$ and $\kappa = \frac{1}{4}$. Also, the functions $u(\varrho, \eta, \varpi) = \frac{\varrho + \eta}{16(1 + \exp(-\varpi(\eta)))}$ and $v(\varrho, \eta, \varpi) = \frac{\arctan \varpi(\eta)}{4(1 + \eta)}$ fulfill the assumption (A3) such that $l = \frac{1}{8}$. Thus, $\kappa l < \frac{1}{4}$ and for $r_0 = 1$, the assumption (A4) is fulfilled. Thus, our second example also fulfills all the five assumptions (A1)–(A5) of Theorem 3.1 and hence by the same theorem, Eq (3.8) has at least one solution in the space $C(I)$.

4. Algorithm to find the solution

In this section, we rephrase and solve Eq (3.7) by using MHP and Adomian decomposition method. The Homotopy perturbation method is a coupling of the idea of homotopy and perturbation methods to eliminate the limitation of traditional perturbation methods. It is also a powerful concept in perturbations theory and topology. In the proposed method, we alter problems of the nonlinear functional equation to some simpler problems and to be free of non-linearity, we apply a linear combination of Adomian polynomials [42]. Thus, we present an iterative algorithm to solve Eq (3.7).

Let us consider the general form of Eq (3.7) as follows:

$$\mathcal{D}(\varrho, \varpi(\varrho)) - f(\varrho, \varpi(\varrho)) = 0, \quad \varrho \in [0, 1]. \quad (4.1)$$

Here, \mathcal{D} is a nonlinear operator and f is some known function. We divide the operator \mathcal{D} according to [43–45] into operators \mathcal{M} and \mathcal{N} to some linear or nonlinear operators. We also divide the function f into f_1 and f_2 . Then Eq (4.1) can be rewritten as $\mathcal{M}(\varrho, \varpi) - f_1(\varrho, \varpi) + \mathcal{N}(\varrho, \varpi) - f_2(\varrho, \varpi) = 0$.

Now, we are enabled to define a MHP as follows:

$$H(p, v) = \mathcal{M}(\varrho, v) - f_1(\varrho, v) + p(\mathcal{N}(\varrho, v) - f_2(\varrho, v)) = 0, \quad (4.2)$$

$$\text{and } v(\varrho) = \sum_{i=0}^{\infty} p^i v_i(\varrho) \text{ together with } \lim_{p \rightarrow 1} v(\varrho) \simeq \varpi(\varrho). \quad (4.3)$$

Herein, $p \in [0, 1]$ and is called an embedding parameter. Letting $p = 0$ to $p = 1$, we are enabled to find $\mathcal{M}(\varrho, v) = f_1(\varrho, v)$ to $\mathcal{N}(\varrho, v) = f_2(\varrho, v)$. Further, in Eq (3.7), we choose the operators as follows:

$$\mathcal{M}(\varrho, \varpi(\varrho)) = \varpi(\varrho),$$

$$\mathcal{N}(\varrho, \varpi(\varrho)) = -\frac{\varrho}{4(1 + \varpi(\varrho)^2)} \int_0^1 \frac{\varrho \eta}{16(1 + \varpi(\eta)^2)} d\eta - \varpi(\varrho) \int_0^{\varrho} \frac{\varrho \eta}{16(1 + \varpi(\eta)^2)^2} d\eta,$$

$$f_1(\varrho, \varpi(\varrho)) = \varrho \text{ and } f_2(\varrho, \varpi(\varrho)) = \frac{\varrho}{4(1 + \varpi(\varrho)^2)} - \varrho.$$

For our ease in calculation, we further divide the operator \mathcal{N} into \mathcal{N}' and \mathcal{N}'' and approximate them by using Adomian decomposition as

$$\begin{aligned}\mathcal{N}'(\varrho, v(\varrho)) &= -\frac{\varrho}{4(1+v(\varrho)^2)} \int_0^1 \frac{\varrho\eta}{16(1+v(\eta)^2)} d\eta = -\sum_{i=0}^{\infty} p^i \mathcal{A}'_i(\varrho), \\ \mathcal{N}''(\varrho, v(\varrho)) &= -v(\varrho) \int_0^{\varrho} \frac{\varrho\eta}{16(1+v(\eta)^2)^2} d\eta = -\sum_{i=0}^{\infty} p^i \mathcal{A}''_i(\varrho), \\ f_2(\varrho, v(\varrho)) &= \sum_{i=0}^{\infty} p^i \mathcal{A}_i(\varrho),\end{aligned}\tag{4.4}$$

wherein the Adomian polynomials are given as follows:

$$\begin{aligned}\mathcal{A}_i(\varrho) &= \frac{1}{i!} \left[\frac{d^i}{dp^i} \left(\frac{\varrho}{4(1+(\sum_{j=0}^i p^j v_j(\varrho))^2)} - \varrho \right) \right]_{p=0}, \\ \mathcal{A}'_i(\varrho) &= \frac{1}{i!} \left[\frac{d^i}{dp^i} \left(\frac{\varrho}{4(1+(\sum_{j=0}^i p^j v_j(\varrho))^2)} \int_0^1 \frac{\varrho\eta}{16(1+(\sum_{j=0}^i p^j v_j(\eta))^2)} d\eta \right) \right]_{p=0}, \\ \mathcal{A}''_i(\varrho) &= \frac{1}{i!} \left[\frac{d^i}{dp^i} \left(\left(\sum_{j=0}^i p^j v_j(\varrho) \right) \int_0^{\varrho} \frac{\varrho\eta}{16(1+(\sum_{j=0}^i p^j v_j(\eta))^2)} d\eta \right) \right]_{p=0}.\end{aligned}$$

Putting Eqs (4.3) and (4.4) in Eq (4.2), we get

$$\mathcal{M} \left(\sum_{i=0}^{\infty} p^i v_i(\varrho) \right) - f_1(\varrho) + p \left(-\sum_{i=0}^{\infty} p^i \mathcal{A}'_i(\varrho) - \sum_{i=0}^{\infty} p^i \mathcal{A}''_i(\varrho) - \sum_{i=0}^{\infty} p^i \mathcal{A}_i(\varrho) \right) = 0.$$

Equating the coefficients of powers of p to zero and solving for v_i , $i = 0, 1, 2, \dots$, we are now enabled to find an iterative algorithm for the numerical solution of Eq (3.7) as follows:

$$\begin{aligned}v_0(\varrho) &= \mathcal{M}^{-1}(f_1(\varrho)), \\ v_j(\varrho) &= \mathcal{M}^{-1}(A_{j-1}(\varrho) + A'_{j-1}(\varrho) + A''_{j-1}(\varrho)) \quad j = 1, 2, \dots\end{aligned}\tag{4.5}$$

Applying the algorithm to the same example, we get $v_0(\varrho) = \varrho$. For the case $j = 1$, we have

$$\begin{aligned}A_0(\varrho) &= \frac{\varrho}{4} \left(\frac{1}{1+\varrho^2} \right) - \varrho, \quad A'_0(\varrho) = \frac{\ln 2}{128} \left(\frac{\varrho^2}{1+\varrho^2} \right), \quad A''_0(\varrho) = \frac{\varrho^4}{32(1+\varrho^2)}, \\ \text{and thus } v_1(\varrho) &= \mathcal{M}^{-1}(A_0(\varrho) + A'_0(\varrho) + A''_0(\varrho)) = \frac{1}{1+\varrho^2} \left(\frac{\ln 2}{128} \varrho^2 + \frac{\varrho}{4} + \frac{\varrho^4}{32} \right) - \varrho.\end{aligned}$$

Now, let us consider the case $j = 2$, we have

$$\begin{aligned}A_1(\varrho) &= \frac{1}{(1+\varrho^2)^3} \left(-\frac{\ln 2}{256} \varrho^4 - \frac{\varrho^3}{8} - \frac{\varrho^6}{64} \right) + \frac{\varrho^3}{2(1+\varrho^2)^2}, \\ A'_1(\varrho) &= \frac{1}{(1+\varrho^2)^3} \left(-\frac{\ln 2}{2048} \varrho^7 - \frac{\ln^2 2}{8192} \varrho^5 - \frac{\ln 2}{256} \varrho^4 \right) + \frac{\ln 2}{64(1+\varrho^2)^2} \varrho^4 + \frac{0.07985190531633895 \varrho^2}{32(1+\varrho^2)},\end{aligned}$$

and

$$A_1''(\varrho) = \frac{\varrho^4}{32(1+\varrho^2)} - \frac{((3\ln 2 + 60)\varrho^8 + (9\ln 2 + 180)\varrho^6 + (9\ln 2 + 180)\varrho^4 + 3\ln 2\varrho^2 + 60\varrho^2) \arctan \varrho}{24576\varrho^6 + 73728\varrho^4 + 73728\varrho^2 + 24576} \\ - \frac{(-1408\varrho^8 + (3\ln 2 - 132)\varrho^7 - 1152\varrho^6 - (8\ln 2 + 160)\varrho^5 - (3\ln 2 + 60)\varrho^3)}{+24576\varrho^6 + 73728\varrho^4 + 73728\varrho^2 + 24576}.$$

Thus, we have

$$v_2(\varrho) = \frac{1}{(1+\varrho^2)^3} \left(-\frac{\ln 2}{2048}\varrho^7 - \frac{\varrho^6}{64} - \frac{\varrho^3}{8} - \frac{\ln 2}{128}\varrho^4 - \frac{\ln^2 2}{8192}\varrho^5 \right) + \frac{1}{(1+\varrho^2)^2} \left(-\frac{\ln 2}{64}\varrho^4 - \frac{\varrho^3}{2} \right) \\ + \frac{1}{1+\varrho^2} \left(0.03125\varrho^4 + 0.0024953720411\varrho^2 \right) \\ - \frac{((3\ln 2 + 60)\varrho^8 + (9\ln 2 + 180)\varrho^6 + (9\ln 2 + 180)\varrho^4 + 3\ln 2\varrho^2 + 60\varrho^2) \arctan \varrho}{24576\varrho^6 + 73728\varrho^4 + 73728\varrho^2 + 24576} \\ - \frac{(-1408\varrho^8 + (3\ln 2 - 132)\varrho^7 - 1152\varrho^6 - (8\ln 2 + 160)\varrho^5 - (3\ln 2 + 60)\varrho^3)}{+24576\varrho^6 + 73728\varrho^4 + 73728\varrho^2 + 24576}.$$


By Eq (4.3), we get an approximated solution,

$$\varpi_1(\varrho) = \frac{1}{1+\varrho^2} \left(\frac{\ln 2}{128}\varrho^2 + \frac{\varrho}{4} + \frac{\varrho^4}{32} \right). \\ \varpi_2(\varrho) = \frac{1}{(1+\varrho^2)^3} \left(-\frac{\ln 2}{2048}\varrho^7 - \frac{\varrho^6}{64} - \frac{\ln^2 2}{8192}\varrho^5 - \frac{\ln 2}{128}\varrho^4 - \frac{\varrho^3}{8} \right) + \frac{1}{(1+\varrho^2)^2} \left(\frac{\ln 2}{64}\varrho^4 + \frac{\varrho^3}{2} \right) \\ + \frac{1}{1+\varrho^2} \left(\frac{\varrho^4}{32} + \frac{\ln 2}{128}\varrho^2 + \frac{\varrho}{4} + 0.03125\varrho^4 + 0.0024953720411\varrho^2 \right) \\ - \frac{((3\ln 2 + 60)\varrho^8 + (9\ln 2 + 180)\varrho^6 + (9\ln 2 + 180)\varrho^4 + 3\ln 2\varrho^2 + 60\varrho^2) \arctan \varrho}{24576\varrho^6 + 73728\varrho^4 + 73728\varrho^2 + 24576} \\ - \frac{(-1408\varrho^8 + (3\ln 2 - 132)\varrho^7 - 1152\varrho^6 - (8\ln 2 + 160)\varrho^5 - (3\ln 2 + 60)\varrho^3)}{+24576\varrho^6 + 73728\varrho^4 + 73728\varrho^2 + 24576}.$$

Thus, by putting $\varpi_1(\varrho)$ together with $\varpi_2(\varrho)$ in Eq (3.7) and equating the both sides, we are enabled to obtain the absolute errors of $\varpi_1(\varrho)$ together with $\varpi_2(\varrho)$ as shown in Table 1. In this way, we are now enabled to observe that increasing the number of cases in the algorithm (4.5), we get better approximations of $\varpi(\varrho)$. The graphs of $v(\varrho)$ and $\varpi(\varrho)$ are shown in Figure 1.

Table 1. Absolute errors.

ϱ	Absolute error of $\varpi_1(\varrho)$	Absolute error of $\varpi_2(\varrho)$
0	0	1.61×10^{-2}
0.1	3×10^{-4}	1.59×10^{-2}
0.2	1.9×10^{-3}	1.42×10^{-2}
0.3	5.9×10^{-3}	1.01×10^{-2}
0.4	1.3×10^{-2}	3.8×10^{-3}
0.5	2.34×10^{-2}	4.2×10^{-3}
0.6	3.68×10^{-2}	1.29×10^{-2}
0.7	5.3×10^{-2}	2.12×10^{-2}
0.8	7.15×10^{-2}	2.82×10^{-2}
0.9	9.18×10^{-2}	3.42×10^{-2}
1.0	1.135×10^{-1}	3.91×10^{-2}

**Figure 1.** Plots of $v(\varrho)$ and $\varpi(\varrho)$.

5. Error analysis

In this section, we give the following two theorems and prove that they also concur with our functional integral equation (1.1).

Theorem 5.1. Let $\varpi_k(\varrho) = \sum_{i=0}^k v_i(\varrho)$, $k \in \mathbb{N}$ and \mathcal{M}^{-1} be a linear operator. Then the following equation is equivalent to the algorithm (4.5):

$$\varpi_{n+1} = \mathcal{M}_1^{-1}(f_1) + \mathcal{M}^{-1}(f_2(\varrho, \varpi_n) - \mathcal{N}(\varpi_n)). \quad (5.1)$$

Proof. Given that

$$\varpi_{n+1}(\varrho) = v_0(\varrho) + v_1(\varrho) + v_2(\varrho) + \dots v_{n+1}(\varrho).$$

By using algorithm (4.5), we are enabled to find

$$\begin{aligned}
 \varpi_{n+1}(\varrho) &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}(A_0(\varrho) + A'_0(\varrho) + A''_0(\varrho)) + \mathcal{M}^{-1}(A_1(\varrho) + A'_1(\varrho) + A''_1(\varrho)) \\
 &\quad + \dots + \mathcal{M}^{-1}(A_n(\varrho) + A'_n(\varrho) + A''_n(\varrho)) \\
 &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(\sum_{i=0}^n \mathcal{A}_i(\varrho) + \sum_{i=0}^n \mathcal{A}'_i(\varrho) + \sum_{i=0}^n \mathcal{A}''_i(\varrho)\right) \\
 &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(f_2\left(\varrho, \sum_{i=0}^n v_i(\varrho)\right) - \mathcal{N}'\left(\varrho, \sum_{i=0}^n v_i(\varrho)\right) - \mathcal{N}''\left(\varrho, \sum_{i=0}^n v_i(\varrho)\right)\right) \\
 &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(f_2\left(\varrho, \sum_{i=0}^n v_i(\varrho)\right) - \mathcal{N}\left(\varrho, \sum_{i=0}^n v_i(\varrho)\right)\right) \\
 &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}(f_2(\varrho, \varpi_n(\varrho)) - \mathcal{N}(\varrho, \varpi_n(\varrho))).
 \end{aligned}$$

Now, we prove the equivalence by using induction. By using assumption, we know that $\varpi_0 = v_0$.

$$\begin{aligned}
 \varpi_1 &= \mathcal{M}^{-1}(f_1) + \mathcal{M}^{-1}(-\mathcal{N}(\varpi_0) + f_2(\varpi_0)) \\
 &= \mathcal{M}^{-1}(f_1) + \mathcal{M}^{-1}(-\mathcal{N}(v_0) + f_2(v_0)) \\
 &= \mathcal{M}^{-1}(f_1) + \mathcal{M}^{-1}(-\mathcal{N}'(v_0) - \mathcal{N}''(v_0) + f_2(v_0)), \\
 \text{i.e., } \varpi_1(\varrho) &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}(A'_0(\varrho) + A''_0(\varrho) + A_0(\varrho)) = v_0(\varrho) + v_1(\varrho).
 \end{aligned}$$

Therefore,

$$v_1(\varrho) = \mathcal{M}^{-1}(A'_0(\varrho) + A''_0(\varrho) + A_0(\varrho)).$$

Hence, the equivalence is satisfied for v_1 . We now verify for v_2 .

$$\begin{aligned}
 \varpi_2 &= \mathcal{M}^{-1}(f_1) + \mathcal{M}^{-1}(-\mathcal{N}(\varpi_1) + f_2(\varpi_1)) \\
 &= \mathcal{M}^{-1}(f_1) + \mathcal{M}^{-1}(-\mathcal{N}'(v_0 + v_1) - \mathcal{N}''(v_0 + v_1) + f_2(v_0 + v_1)), \\
 \text{i.e., } \varpi_2(\varrho) &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}(A'_0(\varrho) + A'_1(\varrho) + A''_0(\varrho) + A''_1(\varrho) + A_0(\varrho) + A_1(\varrho)) \\
 &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}(A'_0(\varrho) + A''_0(\varrho) + A_0(\varrho)) + \mathcal{M}^{-1}(A'_1(\varrho) + A''_1(\varrho) + A_1(\varrho)) \\
 &= v_0(\varrho) + v_1(\varrho) + v_2(\varrho).
 \end{aligned}$$

Therefore,

$$v_2(\varrho) = \mathcal{M}^{-1}(A'_1(\varrho) + A''_1(\varrho) + A_1(\varrho)).$$

By using induction hypothesis, let us assume that the statement is true for v_n and we prove that Eq (5.1) gives rise to v_{n+1} as in the algorithm (4.5).

$$\begin{aligned}
 \varpi_{n+1}(\varrho) &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}(f_2(\varrho, \varpi_n(\varrho)) - \mathcal{N}(\varrho, \varpi_n(\varrho))) \\
 &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(f_2\left(\varrho, \sum_{i=0}^n v_i(\varrho)\right) - \mathcal{N}\left(\varrho, \sum_{i=0}^n v_i(\varrho)\right)\right) \\
 &= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(\sum_{i=0}^n \mathcal{A}_i(\varrho) + \sum_{i=0}^n \mathcal{A}'_i(\varrho) + \sum_{i=0}^n \mathcal{A}''_i(\varrho)\right)
 \end{aligned}$$

$$\begin{aligned}
&= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(\sum_{i=0}^n \mathcal{A}_i(\varrho) + \sum_{i=0}^n \mathcal{A}'_i(\varrho) + \sum_{i=0}^n \mathcal{A}''_i(\varrho)\right) \\
&= \mathcal{M}^{-1}(f_1(\varrho)) + \sum_{i=0}^n \left(\mathcal{M}^{-1}(\mathcal{A}_i(\varrho) + \mathcal{A}'_i(\varrho) + \mathcal{A}''_i(\varrho))\right) \\
&= v_0(\varrho) + \sum_{i=1}^n v_i(\varrho) + \mathcal{M}^{-1}(A_n(\varrho) + A'_n(\varrho) + A''_n(\varrho)) = \sum_{i=0}^{n+1} v_i(\varrho).
\end{aligned}$$

Therefore,

$$v_{n+1}(\varrho) = \mathcal{M}^{-1}(A_n(\varrho) + A'_n(\varrho) + A''_n(\varrho)).$$

Hence, the algorithm is proved. \square

Theorem 5.2. Let $(C[0, 1], \|\varpi\|_\infty)$ be a Banach space and let $\varpi_n(\varrho) = \sum_{i=0}^n v_i(\varrho)$, $n \in \mathbb{N}$ such that $\|v_i\|_\infty \leq \alpha \|v_{i-1}\|_\infty$, $\forall i \in \mathbb{N}$, where $0 \leq \alpha < 1$, then:

- (i) The sequence $\{\varpi_n\}_{n=0}^\infty$ is convergent.
- (ii) The limit of the sequence say $\lim_{n \rightarrow \infty} \varpi_n = \varpi^*$ fulfills the algorithm (4.5) and Eq (4.1).

Proof. (i) It is enough to show the sequence is Cauchy.

Let $m, n \in \mathbb{N}$ with $m > n$ and let $\epsilon > 0$. Then, for $\varrho \in I$, with the above assumptions, we estimate

$$\begin{aligned}
|\varpi_m(\varrho) - \varpi_n(\varrho)| &= \left| \sum_{i=n+1}^m v_i(\varrho) \right| = |v_{n+1}(\varrho) + v_{n+2}(\varrho) + \dots + v_m(\varrho)| \\
&\leq \|v_{n+1}\|_\infty + \dots + \|v_m\|_\infty \\
&\leq \alpha^{n+1} \|v_0\|_\infty + \alpha^{n+2} \|v_0\|_\infty + \dots + \alpha^m \|v_0\|_\infty \\
&= \left(\frac{1 - \alpha^{m-n}}{1 - \alpha} \right) \alpha^{n+1} \|v_0\|_\infty, \\
\text{i.e., } \|\varpi_m - \varpi_n\|_\infty &\leq \left(\frac{\alpha^{n+1}}{1 - \alpha} \right) \|v_0\|_\infty < \epsilon.
\end{aligned}$$

Hence, the sequence $\{\varpi_n\}_{n=0}^\infty$ is Cauchy in $(C[0, 1], \|\varpi\|_\infty)$, i.e., $\lim_{n \rightarrow \infty} \varpi_n(\varrho) = \lim_{n \rightarrow \infty} \sum_{i=0}^n v_i(\varrho) = \sum_{i=0}^\infty v_i(\varrho) = \varpi^*(\varrho)$.

- (ii) We first prove that the limit of the sequence fulfills the algorithm (4.5). Let $\{\varpi_n\} \rightarrow \varpi^*$, we estimate

$$\begin{aligned}
\lim_{n \rightarrow \infty} \varpi_n(\varrho) &= \lim_{n \rightarrow \infty} \left(\mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(-\mathcal{N}(\varrho, \varpi_{n-1}(\varrho)) + f_2(\varrho, \varpi_{n-1}(\varrho))\right) \right) \\
&= \lim_{n \rightarrow \infty} \left(\mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(-\mathcal{N}\left(\varrho, \sum_{i=0}^{n-1} v_i(\varrho)\right) + f_2\left(\varrho, \sum_{i=0}^{n-1} v_i(\varrho)\right)\right) \right) \\
&= \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(-\lim_{n \rightarrow \infty} \mathcal{N}'\left(\varrho, \sum_{i=0}^{n-1} v_i(\varrho)\right) - \lim_{n \rightarrow \infty} \mathcal{N}''\left(\varrho, \sum_{i=0}^{n-1} v_i(\varrho)\right)\right)
\end{aligned}$$

$$\begin{aligned}
& + \lim_{n \rightarrow \infty} f_2\left(\varrho, \sum_{i=0}^{n-1} v_i(\varrho)\right) \\
& = \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(\lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} \mathcal{A}'_i(\varrho) + \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} A''_i(\varrho) + \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} \mathcal{A}_i(\varrho)\right) \\
& = \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(\sum_{i=0}^{\infty} \mathcal{A}'_i(\varrho) + \sum_{i=0}^{\infty} A''_i(\varrho) + \sum_{i=0}^{\infty} \mathcal{A}_i(\varrho)\right), \\
\text{i.e., } \lim_{n \rightarrow \infty} \varpi_n(\varrho) & = \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(-\mathcal{N}'\left(\varrho, \sum_{i=0}^{\infty} v_i(\varrho)\right) - \mathcal{N}''\left(\varrho, \sum_{i=0}^{\infty} v_i(\varrho)\right) + f_2\left(\varrho, \sum_{i=0}^{\infty} v_i(\varrho)\right)\right).
\end{aligned}$$

In other words,

$$\varpi^*(\varrho) = \mathcal{M}^{-1}(f_1(\varrho)) + \mathcal{M}^{-1}\left(-\mathcal{N}(\varrho, \varpi^*(\varrho)) + f_2(\varrho, \varpi^*(\varrho))\right). \quad (5.2)$$

Hence, the algorithm is proved. Now for proving the Eq (4.1), we apply the operator \mathcal{M} to the above equation as

$$\mathcal{M}(\varpi^*) = f_1(\varrho) - \mathcal{N}(\varpi^*) + f_2(\varpi^*) \Rightarrow \mathcal{M}(\varpi^*) + \mathcal{N}(\varpi^*) = f_1(\varrho) + f_2(\varpi^*) \Rightarrow A(\varpi^*) = f(\varpi^*),$$

and the proof is completed. □

The following corollary gives upper bound of the errors.

Corollary 5.1. *Under the assumptions of the above theorem, $\|\varpi^* - \varpi_n\|_{\infty} \leq \left(\frac{\alpha^{n+1}}{1-\alpha}\right) \|f_1\|_{\infty}, \forall n \in \mathbb{N}$.*

6. Conclusions

We have thus verified and proved the existence result of the considered nonlinear functional integral equation on the Banach space $C[0, 1]$. The result is obtained by the applications of the generalized Darbo fixed point theorem associated with the MNC in the Banach space. Our result is demonstrated with the two examples. Also, we have introduced a numerical algorithm by using the MHP approach along with the Adomian decomposition method to find the approximate solution with relevant accuracy. Moreover, an error analysis with the upper bound of errors and the condition of convergence are presented. In this paper, the MATLAB program has been used for computations and programming.

Conflict of interest

The authors declare no conflict of interest.

References

1. M. A. Abdou, On the solution of linear and nonlinear integral equation, *Appl. Math. Comput.*, **146** (2003), 857–871. [https://doi.org/10.1016/S0096-3003\(02\)00643-4](https://doi.org/10.1016/S0096-3003(02)00643-4)

2. I. K. Argyros, Quadratic equations and applications to Chandrasekhar's and related equations, *Bull. Aust. Math. Soc.*, **32** (1985), 275–292. <https://doi.org/10.1017/S0004972700009953>
3. A. Aghajani, J. Banaś, Y. Jalilian, Existence of solutions for a class of nonlinear Volterra singular integral equations, *Comput. Math. Appl.*, **62** (2011), 1215–1227. <https://doi.org/10.1016/j.camwa.2011.03.049>
4. A. Aghajani, Y. Jalilian, Existence and global attractivity of solutions of a nonlinear functional integral equation, *Commun. Nonlinear Sci. Numer. Simul.*, **15** (2010), 3306–3312. <https://doi.org/10.1016/j.cnsns.2009.12.035>
5. E. Alvarez, C. Lizama, Attractivity for functional Volterra integral equations of convolution type, *J. Comput. Appl. Math.*, **301** (2016), 230–240. <https://doi.org/10.1016/j.cam.2016.01.048>
6. J. Banaś, K. Balachandran, D. Julie, Existence and global attractivity of solutions of a nonlinear functional integral equation, *Appl. Math. Comput.*, **216** (2010), 261–268. <https://doi.org/10.1016/j.amc.2010.01.049>
7. J. Banaś, K. Goebel, *Measure of noncompactness in Banach spaces*, Lecture Notes in Pure and Applied Mathematics, Vol. 60, New York: Dekker, 1980.
8. J. Banaś, L. Olszowy, Measures of noncompactness related to monotonicity, In: *Annales Societatis Mathematicae Polonae. Seria I: Commentationes Mathematicae*, 2001, 13–23.
9. J. Banaś, B. Rzepka, On existence and asymptotic stability of solutions of a nonlinear integral equation, *J. Math. Anal. Appl.*, **284** (2003), 165–173. [https://doi.org/10.1016/S0022-247X\(03\)00300-7](https://doi.org/10.1016/S0022-247X(03)00300-7)
10. J. Banaś, B. Rzepka, On local attractivity and asymptotic stability of solutions of a quadratic Volterra integral equation, *Appl. Math. Comput.*, **213** (2009), 102–111. <https://doi.org/10.1016/j.amc.2009.02.048>
11. D. Chalishajar, C. Ravichandran, S. Dhanalakshmi, R. Murugesu, Existence of fractional impulsive functional integro-differential equations in Banach spaces, *Appl. Syst. Innov.*, **2** (2019), 1–17. <https://doi.org/10.3390/asi2020018>
12. B. C. Dhage, Global attractivity results for nonlinear functional integral equations via a Krasnoselskii type fixed point theorem, *Nonlinear Anal.*, **70** (2009), 2485–2493. <https://doi.org/10.1016/j.na.2008.03.033>
13. B. C. Dhage, Local asymptotic attractivity for nonlinear quadratic functional integral equations, *Nonlinear Anal.*, **70** (2009), 1912–1922. <https://doi.org/10.1016/j.na.2008.02.109>
14. A. Deep, Deepmala, J. R. Roshan, Solvability for generalized nonlinear functional integral equations in Banach spaces with applications, *J. Integral Equ. Appl.*, **33** (2021), 19–30. <https://doi.org/10.1216/jie.2021.33.19>
15. A. Deep, Deepmala, C. Tunç, On the existence of solutions of some non-linear functional integral equations in Banach algebra with applications, *Arab J. Basic Appl. Sci.*, **27** (2020), 279–286. <https://doi.org/10.1080/25765299.2020.1796199>
16. B. C. Dhage, S. B. Dhage, H. K. Pathak, A generalization of Darbo's fixed point theorem and local attractivity of generalized nonlinear functional integral equations, *Differ. Equ. Appl.*, **7** (2015), 57–77. <https://doi.org/10.7153/dea-07-05>

17. B. C. Dhage, V. Lakshmikantham, On global existence and attractivity results for nonlinear functional integral equations, *Nonlinear Anal.*, **72** (2010), 2219–2227. <https://doi.org/10.1016/j.na.2009.10.021>
18. A. M. A. El-Sayed, H. R. Ebead, On the solvability of a self-reference functional and quadratic functional integral equations, *Filomat*, **34** (2020), 129–141. <https://doi.org/10.2298/FIL2001129E>
19. R. C. Guerra, On the solution of a class of integral equations using new weighted convolutions, *J. Integral Equ. Appl.*, **34** (2022), 39–58. <https://doi.org/10.1216/jie.2022.34.39>
20. X. L. Hu, J. R. Yan, The global attractivity and asymptotic stability of solution of a nonlinear integral equation, *J. Math. Anal. Appl.*, **321** (2006), 147–156. <https://doi.org/10.1016/j.jmaa.2005.08.010>
21. K. Jothimani, K. Kaliraj, S. K. Panda, K. S. Nisar, C. Ravichandran, Results on controllability of non-densely characterized neutral fractional delay differential system, *Evol. Equ. Control Theory*, **10** (2021), 619–631. <https://doi.org/10.3934/eect.2020083>
22. K. Jangid, S. D. Purohit, R. Agarwal, On Gruss type inequality involving a fractional integral operator with a multi-index Mittag-Leffler function as a kernel, *Appl. Math. Inf. Sci.*, **16** (2022), 269–276. <https://doi.org/10.18576/amis/160214>
23. A. Karimi, K. Maleknejad, R. Ezzati, Numerical solutions of system of two-dimensional Volterra integral equations via Legendre wavelets and convergence, *Appl. Numer. Math.*, **156** (2020), 228–241. <https://doi.org/10.1016/j.apnum.2020.05.003>
24. S. Karmakar, H. Garai, L. K. Dey, A. Chanda, Existence of solutions to non-linear quadratic integral equations via measure of non-compactness, *Filomat*, **36** (2022), 73–87. <https://doi.org/10.2298/FIL2201073K>
25. A. Karoui, A. Jawahdou, Existence and approximate L^p and continuous solutions of nonlinear integral equations of the Hammerstein and Volterra types, *Appl. Math. Comput.*, **216** (2010), 2077–2091. <https://doi.org/10.1016/j.amc.2010.03.042>
26. K. Logeswari, C. Ravichandran, K. S. Nisar, Mathematical model for spreading of COVID-19 virus with the Mittag-Leffler kernel, *Numer. Methods Partial Differ. Equ.*, 2020, 1–16. <https://doi.org/10.1002/num.22652>
27. L. N. Mishra, R. P. Agarwal, On existence theorems for some nonlinear functional-integral equations, *Dyn. Syst. Appl.*, **25** (2016), 303–320.
28. L. N. Mishra, R. P. Agarwal, M. Sen, Solvability and asymptotic behavior for some nonlinear quadratic integral equation involving Erdélyi-Kober fractional integrals on the unbounded interval, *Progr. Fract. Differ. Appl.*, **2** (2016), 153–168. <https://doi.org/10.18576/pfda/020301>
29. K. Maleknejad, K. Nouri, R. Mollapourasl, Investigation on the existence of solutions for some nonlinear functional-integral equations, *Nonlinear Anal.*, **71** (2009), e1575–e1578. <https://doi.org/10.1016/j.na.2009.01.207>
30. B. Matani, J. R. Roshan, Multivariate generalized Meir-Keeler condensing operators and their applications to systems of integral equations, *J. Fixed Point Theory Appl.*, **22** (2020), 1–28. <https://doi.org/10.1007/s11784-020-00820-6>

31. B. Matani, J. R. Roshan, N. Hussain, An extension of Darbo's theorem via measure of non-compactness with its application in the solvability of a system of integral equations, *Filomat*, **33** (2019), 6315–6334. <https://doi.org/10.2298/FIL1919315M>
32. L. N. Mishra, M. Sen, On the concept of existence and local attractivity of solutions for some quadratic Volterra integral equation of fractional order, *Appl. Math. Comput.*, **285** (2016), 174–183. <https://doi.org/10.1016/j.amc.2016.03.002>
33. L. N. Mishra, M. Sen, R. N. Mohapatra, On existence theorems for some generalized nonlinear functional-integral equations with applications, *Filomat*, **31** (2017), 2081–2091. <https://doi.org/10.2298/FIL1707081N>
34. L. N. Mishra, H. M. Srivastava, M. Sen, Existence results for some nonlinear functional-integral equations in Banach algebra with applications, *Int. J. Anal. Appl.*, **11** (2016), 1–10.
35. E. Najafi, Nyström-quasilinearization method and smoothing transformation for the numerical solution of nonlinear weakly singular Fredholm integral equations, *J. Comput. Appl. Math.*, **368** (2020), 112538. <https://doi.org/10.1016/j.cam.2019.112538>
36. H. Nasiri, J. R. Roshan, M. Mursaleen, Solvability of system of Volterra integral equations via measure of noncompactness, *Comput. Appl. Math.*, **40** (2021), 1–25. <https://doi.org/10.1007/s40314-021-01552-0>
37. K. S. Nisar, K. Jothimani, K. Kaliraj, C. Ravichandran, An analysis of controllability results for nonlinear Hilfer neutral fractional derivatives with non-dense domain, *Chaos Solitons Fract.*, **146** (2021), 110915. <https://doi.org/10.1016/j.chaos.2021.110915>
38. S. Noeiaghdam, M. A. F. Araghi, D. Sidorov, Dynamical strategy on homotopy perturbation method for solving second kind integral equations using the CESTAC method, *J. Comput. Appl. Math.*, **411** (2022), 114226. <https://doi.org/10.1016/j.cam.2022.114226>
39. D. O'Regan, Existence results for nonlinear integral equations, *J. Math. Anal. Appl.*, **192** (1995), 705–726. <https://doi.org/10.1006/jmaa.1995.1199>
40. Y. B. Pan, J. Huang, Extrapolation method for solving two-dimensional Volterra integral equations of the second kind, *Appl. Math. Comput.*, **367** (2020), 124784. <https://doi.org/10.1016/j.amc.2019.124784>
41. V. K. Pathak, L. N. Mishra, Application of fixed point theorem to solvability for non-linear fractional Hadamard functional integral equations, *Mathematics*, **10** (2022), 1–16. <https://doi.org/10.3390/math10142400>
42. M. Rabbani, B. Zarali, Solution of Fredholm integro-differential equations system by modified decomposition method, *J. Math. Comput. Sci.*, **5** (2012), 258–264.
43. M. Rabbani, New homotopy perturbation method to solve non-linear problems, *J. Math. Comput. Sci.*, **7** (2013), 272–275.
44. M. Rabbani, Modified homotopy method to solve non-linear integral equations, *Int. J. Nonlinear Anal. Appl.*, **6** (2015), 133–136. <https://doi.org/10.22075/IJNAA.2015.262>
45. M. Rabbani, R. Arab, Extension of some theorems to find solution of nonlinear integral equation and homotopy perturbation method to solve it, *Math. Sci.*, **11** (2017), 87–94. <https://doi.org/10.1007/s40096-017-0206-4>

46. M. Rabbani, A. Deep, Deepmala, On some generalized non-linear functional integral equations of two variables via measures of noncompactness and numerical method to solve it, *Math. Sci.*, **15** (2021), 317–324. <https://doi.org/10.1007/s40096-020-00367-0>
47. J. R. Roshan, Existence of solutions for a class of system of functional integral equation via measure of noncompactness, *J. Comput. Appl. Math.*, **313** (2017), 129–141. <https://doi.org/10.1016/j.cam.2016.09.011>
48. S. K. Sahoo, P. O. Mohammed, B. Kodamasingh, M. Tariq, Y. S. Hamed, New fractional integral inequalities for convex functions pertaining to Caputo-Fabrizio operator, *Fractal Fract.*, **6** (2022), 1–17. <https://doi.org/10.3390/fractalfract6030171>
49. H. M. Srivastava, S. K. Sahoo, P. O. Mohammed, B. Kodamasingh, Y. S. Hamed, New Riemann-Liouville fractional-order inclusions for convex functions via integral-valued setting associated with pseudo-order relations, *Fractal Fract.*, **6** (2022), 1–17. <https://doi.org/10.3390/fractalfract6040212>
50. N. Valliammal, C. Ravichandran, Results on fractional neutral integro-differential systems with state-dependent delay in Banach spaces, *Nonlinear Stud.*, **25** (2018), 159–171.
51. J. R. Wang, X. W. Dong, Y. Zhou, Existence, attractiveness and stability of solutions for quadratic Urysohn fractional integral equations, *Commun. Nonlinear Sci. Numer. Simul.*, **17** (2012), 545–554. <https://doi.org/10.1016/j.cnsns.2011.05.034>
52. X. Y. Zhang, A new strategy for the numerical solution of nonlinear Volterra integral equations with vanishing delays, *Appl. Math. Comput.*, **365** (2020), 124608. <https://doi.org/10.1016/j.amc.2019.124608>
53. J. K. Xu, H. X. Wu, Z. Tan, Radial symmetry and asymptotic behaviors of positive solutions for certain nonlinear integral equations, *J. Math. Anal. Appl.*, **427** (2015), 307–319. <https://doi.org/10.1016/j.jmaa.2015.02.043>



AIMS Press

© 2022 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)