



Research article

Nonlinear fractional differential equations and their existence via fixed point theory concerning to Hilfer generalized proportional fractional derivative

Saima Rashid^{1,*}, Abdulaziz Garba Ahmad², Fahd Jarad^{3,4,5,*} and Ateq Alsaadi⁶

¹ Department of Mathematics, Government College University, Faisalabad, Pakistan

² Department of Mathematics Programme, National Mathematical Centre, Abuja, Nigeria

³ Department of Mathematics, Çankaya University, Ankara, Turkey

⁴ Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan

⁵ Department of Mathematics, King Abdulaziz University, Jeddah, Saudi Arabia

⁶ Department of Mathematics and Statistics, College of Science, Taif University, P.O.Box 11099, Taif 21944, Saudi Arabia

* **Correspondence:** Emails: saimarashid@gcuf.edu.pk, fahd@cankaya.edu.tr.

Abstract: This article adopts a class of nonlinear fractional differential equation associating Hilfer generalized proportional fractional (*GPF*) derivative with having boundary conditions, which amalgamates the Riemann-Liouville (*RL*) and Caputo-*GPF* derivative. Taking into consideration the weighted space continuous mappings, we first derive a corresponding integral for the specified boundary value problem. Also, we investigate the existence consequences for a certain problem with a new unified formulation considering the minimal suppositions on nonlinear mapping. Detailed developments hold in the analysis and are dependent on diverse tools involving Schauder's, Schaefer's and Krasnoselskii's fixed point theorems. Finally, we deliver two examples to check the efficiency of the proposed scheme.

Keywords: existence of solution; Hilfer proportional fractional derivative; boundary value problem; fixed point theory

Mathematics Subject Classification: 46S40, 47H10, 54H25

1. Introduction

Fractional calculus has attempted to be accessed as a promising technique in fluid mechanics [1], nano-material [2], thermal energy [3], epidemics [4] and other scientific disciplines over recent

decades. For example, by provoking interest in both cutting-edge and conventional pure and applied analytical techniques, it has reinforced creative collaboration between different disciplines, existence, and relevant applications in real-world manifestations, see [5–7]. In 1965, the possibility of fractional calculus depended on the conversation between L’Hospital and Leibnitz as letters in [8]. After that, many researchers started experimenting in this field, and a large portion of them concentrated on describing novel fractional formulations [9–11]. Various classifications were raised in this process and were expressed by the advancement of research.

Recently, extensive investigation has been proposed for the qualitative characterization of verification for various fractional differential equations (*FDEs*) with initial and boundary value problems. Several significant approaches regarding the existence, uniqueness, multiplicity and stability have been reported by proposing certain fixed point theorems. Although many of the important problems have been tackled by the classical fractional derivatives (*RL* and Caputo) [12–14], it has a few limitations when used to design physical issues as a result of the necessary assumptions are themselves fractional and may be unsuitable for physical problems. The Caputo derivative has the opportunity of being appropriate for physical problems because it only necessitates classical initial conditions [15, 16].

Khalil et al. [17] invented an interesting definition of fractional derivative, which is known to be a conformable derivative. In actuality, this so-called derivative is not a fractional derivative, however, it is essentially a first derivative duplicated by an extra straightforward factor. Consequently, this novel concept appears to be a regular extension of the classical derivative. More characterizations and the extended form of this derivative have been expounded in [18]. Then authors [19] explored an extension of the conformable derivative by considering proportional derivative. This fact leads to the modified conformable (proportional) differential operator of order λ . The researchers investigated numerous integral inequalities using classical [20], conformable and generalized conformable fractional integrals [21, 22]. Qurashi et al. [23] proposed new fractional derivatives and integrals that have nonsingular kernels. By using the generalized proportional Hadamard fractional integral operator, Zhou et al. [24] investigated some general inequalities and their variant forms.

This recently characterized local derivative approaches to the original function as $\lambda \mapsto 0$. In this manner, they had the option to improve the conformable derivative. Jarad et al. [25] presented another kind of fractional operators created from the extended conformable derivatives. The exponential function appeared as a kernel in their examination with outstanding performance [26–28]. For the interest of readers, we draw in their thoughtfulness regarding some new papers [29, 30].

In parallel with the concentrated exploration of the fractional derivative, the existence-uniqueness of verification fits to the intense prominent qualitative characterizations of *FDEs*, see [31–34].

Inspired by the work, we utilize a novel fractional derivative which is known as Hilfer-*GPF* derivative for finding the existence-uniqueness of solutions for a new class of nonlinear *FDEs* having non-local boundary conditions. For this we consider the subsequent *BVP* for a class of Hilfer-*FDEs*:

$$\begin{cases} \mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} y(\varphi) = \mathcal{G}(\varphi, y(\varphi)), & \lambda \in (0, 1), \zeta \in [0, 1], \varphi \in (\varpi_1, \varpi_2], \\ \mathcal{I}_{\varpi_1^+}^{1-\delta; \vartheta} [m_1 y(\varpi_1^+) + m_2 y(\varpi_2^-)] = e_j, & \delta = \lambda + \zeta(1 - \lambda), e_j \in \mathbb{R}, \end{cases} \quad (1.1)$$

where $\mathcal{G} : (\varpi_1, \varpi_2] \times \mathbb{R} \mapsto \mathbb{R}$ be a continuous mapping, $\mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta}(\cdot)$ is the Hilfer-*GPF* derivative of order $\lambda \in (0, 1)$ and $\mathcal{I}_{\varpi_1^+}^{1-\delta; \vartheta}(\cdot)$ is the *GPF* integral of order $1 - \delta > 0$. We find the existence consequences by

the fixed point techniques of Schauder, Schaefer and Kransnoselskii. Additionally, the investigation of nonlinear *FDEs* as far as their information sources (fractional orders, related boundaries, and suitable function) has fascinated the interest of mathematicians because of its importance in Orlicz space (see [35]). Rely upon this, the subject of coherence of verification of the Hilfer-*FDEs* regarding inputs is significant and worth assuming.

The organization of the paper is as follows. In Section 2, we proceed with some basics concepts and detailed consequences as a review literature. In Section 3, we establish an equivalence criterion of integral equation of *BVP* (1.1) and then proposed the existence consequences for *GPF*-derivative by well-noted fixed point theorem. Also, in Section 4, illustrative examples are presented to check the applicability of the findings developed in Section 3. The conclusion with some open problems is presented in Section 5.

2. Preliminaries

In what follows, we demonstrate some preliminaries, initial results and spaces which are essential for proving further consequences. Throughout this investigation, let $L^p(\varpi_1, \varpi_2)$, $p \geq 1$, is the space of Lebesgue integrable mappings on (ϖ_1, ϖ_2) .

Assume that $\varpi_1, \varpi_2 \in (-\infty, +\infty)$ be a finite and infinite intervals on \mathbb{R} .

Furthermore, we elaborate the subsequent weighted spaces with induced norms defined by (see [8]). Suppose that $\mathbb{C}[\varpi_1, \varpi_2]$ is said to be the space of continuous functions defined on $[\varpi_1, \varpi_2]$ and the norm is defined as follows:

$$\|\mathcal{G}\|_{\mathbb{C}[\varpi_1, \varpi_2]} = \max_{\varphi \in [\varpi_1, \varpi_2]} |\mathcal{G}|,$$

and $\mathcal{AC}^n[\varpi_1, \varpi_2]$ represents the space of n -times absolutely continuous differentiable mappings defined as follows:

$$\mathcal{AC}^n[\varpi_1, \varpi_2] = \{\mathcal{G} : (\varpi_1, \varpi_2) \mapsto \mathbb{R} : \mathcal{G}^{n-1} \in \mathcal{AC}[\varpi_1, \varpi_2]\},$$

$\mathbb{C}_\delta[\varpi_1, \varpi_2]$ denotes the weighted space of \mathcal{G} on $(\varpi_1, \varpi_2]$ is defined as

$$\mathbb{C}_\delta^\delta[\varpi_1, \varpi_2] = \{\mathcal{G} : (\varpi_1, \varpi_2) \mapsto \mathbb{R} : (\varphi - \varpi_1)^\delta \mathcal{G}(\varphi) \in \mathbb{C}[\varpi_1, \varpi_2]\}, \delta \in [0, 1)$$

with the norm

$$\|\mathcal{G}\|_{\mathbb{C}_\delta^\delta[\varpi_1, \varpi_2]} = \|(\varphi - \varpi_1)^\delta \mathcal{G}(\varphi)\|_{\mathbb{C}[\varpi_1, \varpi_2]} = \max_{\varphi \in [\varpi_1, \varpi_2]} |(\varphi - \varpi_1)^\delta \mathcal{G}|.$$

Also, the weighted space of a function \mathcal{G} on $(\varpi_1, \varpi_2]$ is denoted by $\mathbb{C}_\delta^n(\varpi_1, \varpi_2]$ defined as

$$\mathbb{C}_\delta^n[\varpi_1, \varpi_2] = \{\mathcal{G} : (\varpi_1, \varpi_2) \mapsto \mathbb{R} : \mathcal{G}(\varphi) \in \mathbb{C}^{n-1}[\varpi_1, \varpi_2]; \mathcal{G}^n(\varphi) \in \mathbb{C}_\delta[\varpi_1, \varpi_2]\}, \delta \in [0, 1)$$

with the norm

$$\|\mathcal{G}\|_{\mathbb{C}_\delta^n[\varpi_1, \varpi_2]} = \sum_{k=0}^{n-1} \|\mathcal{G}^k\|_{\mathbb{C}[\varpi_1, \varpi_2]} + \|\mathcal{G}^n\|_{\mathbb{C}_\delta[\varpi_1, \varpi_2]}, \forall n \in \mathbb{N}.$$

For $n = 0$, $\mathbb{C}_\delta^n[\varpi_1, \varpi_2]$ coincides with $\mathbb{C}_\delta[\varpi_1, \varpi_2]$.

Definition 2.1. ([8]) Assume that $\mathcal{G} \in L_1([\varpi_1, \varpi_2], \mathbb{R})$, then the RL fractional integral operator of \mathcal{G} of order $\lambda > 0$ is stated as

$$\mathcal{J}_{\varpi_1^+}^\lambda \mathcal{G}(\varphi) = \frac{1}{\Gamma(\lambda)} \int_{\varpi_1}^{\varphi} (\varphi - \ell)^{\lambda-1} \mathcal{G}(\ell) d\ell, \varphi > \varpi_1, \quad (2.1)$$

where $\Gamma(\cdot)$ represents the classical Gamma function.

Definition 2.2. ([8]) Assume that $\mathcal{G} \in \mathbb{C}([\varpi_1, \varpi_2])$, then the RL fractional derivative operator of \mathcal{G} of order $\lambda > 0$ is stated as

$$\mathcal{D}_{\varpi_1^+}^\lambda \mathcal{G}(\varphi) = \frac{1}{\Gamma(n-\lambda)} \frac{d^n}{d\varphi^n} \int_{\varpi_1}^{\varphi} (\varphi - \ell)^{n-\lambda-1} \mathcal{G}(\ell) d\ell, \varphi > \varpi_1, n-1 < \lambda < n, n \in \mathbb{N}, \quad (2.2)$$

where $\Gamma(\cdot)$ represents the Gamma function.

Definition 2.3. ([8]) Assume that $\mathcal{G} \in \mathbb{C}([\varpi_1, \varpi_2])$, then the Caputo fractional derivative operator of \mathcal{G} of order $\lambda > 0$ is stated as

$${}^c \mathcal{D}_{\varpi_1^+}^\lambda \mathcal{G}(\varphi) = \frac{1}{\Gamma(n-\lambda)} \int_{\varpi_1}^{\varphi} (\varphi - \ell)^{n-\lambda-1} \mathcal{G}^n(\ell) d\ell, \varphi > \varpi_1, n-1 < \lambda < n, n \in \mathbb{N}, \quad (2.3)$$

where $\Gamma(\cdot)$ represents the Gamma function.

Definition 2.4. ([25]) For $\vartheta \in (0, 1]$, $\lambda \in \mathbb{C}$, $\Re(\lambda) > 0$, then the left-sided generalized proportional integral of \mathcal{G} of order $\lambda > 0$ is stated as

$$\mathcal{J}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{G}(\varphi) = \frac{1}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} (\varphi - \ell)^{\lambda-1} \mathcal{G}(\ell) d\ell, \varphi > \varpi_1. \quad (2.4)$$

Definition 2.5. ([25]) For $\vartheta \in (0, 1]$, $\lambda \in \mathbb{C}$, $\Re(\lambda) > 0$, then the left-sided generalized proportional derivative of \mathcal{G} of order $\lambda > 0$ is stated as

$$\mathcal{D}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{G}(\varphi) = \frac{\mathcal{D}^{n, \vartheta}}{\vartheta^{n-\lambda} \Gamma(n-\lambda)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} (\varphi - \ell)^{n-\lambda-1} \mathcal{G}(\ell) d\ell, \varphi > \varpi_1, \quad (2.5)$$

where $n = [\lambda] + 1$.

Definition 2.6. ([25]) For $\vartheta \in (0, 1]$, $\lambda \in \mathbb{C}$, $\Re(\lambda) > 0$, then the left-sided generalized proportional integral of \mathcal{G} of order $\lambda > 0$ is stated as

$${}^c \mathcal{D}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{G}(\varphi) = \frac{1}{\vartheta^{n-\lambda} \Gamma(n-\lambda)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} (\varphi-\ell)^{n-\lambda-1} (\mathcal{D}^{n, \vartheta} \mathcal{G})(\ell) d\ell, \quad \varphi > \varpi_1, \quad (2.6)$$

where $n = [\lambda] + 1$.

Remark 2.1. Specifically, if $\vartheta = 1$ Definitions 2.4–2.6 reduces to Definitions 2.1–2.3, respectively.

Definition 2.7. ([25]) For $n \in \mathbb{N}$, $\lambda \in (n-1, n)$, $\vartheta \in (0, 1]$, $\zeta \in [0, 1]$, then the left/right-sided Hilfer-GPF derivative having order λ , type ζ of \mathcal{G} is stated as follows:

$$(\mathcal{D}_{\varpi_1^+} \mathcal{G})(y) = \mathcal{I}_{\varpi_1^+}^{\zeta(n-\lambda), \vartheta} [\mathcal{D}^{\vartheta} (\mathcal{I}_{\varpi_1^+}^{(1-\zeta)(n-\lambda), \vartheta} \mathcal{G})](y), \quad (2.7)$$

where $\mathcal{D}^{\vartheta} \mathcal{G}(y) = (1-\vartheta)\mathcal{G}(y) + \vartheta \mathcal{G}'(y)$ and \mathcal{I} assumed to be GPF-integral stated in 2.4.

Specifically, if $n = 1$, then Definition 2.7 reduces to

$$(\mathcal{D}_{\varpi_1^+} \mathcal{G})(y) = \mathcal{I}_{\varpi_1^+}^{\zeta(1-\lambda), \vartheta} [\mathcal{D}^{\vartheta} (\mathcal{I}_{\varpi_1^+}^{(1-\zeta)(1-\lambda), \vartheta} \mathcal{G})](y). \quad (2.8)$$

In the present investigation, we discuss the case where $n = 1$, $\lambda \in (0, 1)$, $\zeta \in [0, 1]$ and $\delta = \lambda + \zeta - \lambda\zeta$.

Remark 2.2. It is remarkable to mention that:

(a) The Hilfer fractional derivative can be considered as an interpolator between the GPF-derivative and Caputo GPF-derivative, respectively, as

$$\mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} \mathcal{G} = \begin{cases} \mathcal{D}^{\vartheta} \mathcal{I}_{\varpi_1^+}^{(1-\lambda); \vartheta} \mathcal{G}, & \zeta = 0 \text{ (see Definition 2.5),} \\ \mathcal{I}_{\varpi_1^+}^{1-\lambda; \vartheta} \mathcal{D}^{\vartheta} \mathcal{G}, & \zeta = 1 \text{ (see Definition 2.6),} \end{cases} \quad (2.9)$$

(b) The following assumptions holds true:

$$0 < \delta \leq 1 \quad \delta \geq \lambda, \quad \delta > \zeta, \quad 1 - \delta < 1 - \zeta(1 - \lambda).$$

(c) Particularly, if $\lambda \in (0, 1)$, $\zeta \in [0, 1]$ and $\delta = \lambda + \zeta - \lambda\zeta$, then

$$(\mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} \mathcal{G})(\varphi) = \left(\mathcal{I}_{\varpi_1^+}^{\zeta(1-\lambda)} [\mathcal{D}^{\vartheta} (\mathcal{I}_{\varpi_1^+}^{(1-\zeta)(1-\lambda)} \mathcal{G})] \right)(\varphi),$$

therefore, we have

$$(\mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} \mathcal{G})(\varphi) = \left(\mathcal{I}_{\varpi_1^+}^{\zeta(1-\lambda)} (\mathcal{D}_{\varpi_1^+}^{\delta; \vartheta} \mathcal{G}) \right)(\varphi),$$

where $(\mathcal{D}_{\varpi_1^+}^{\delta; \vartheta} \mathcal{G})(\varphi) = \frac{d}{d\varphi} (\mathcal{I}_{\varpi_1^+}^{(1-\zeta)(1-\lambda); \vartheta} \mathcal{G})(\varphi)$.

Now we define the weighted spaces of continuous mappings on $(\varpi_1, \varpi_2]$:

$$\mathbb{C}_{1-\delta}^{\lambda, \zeta; \vartheta}[\varpi_1, \varpi_2] = \{\mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2], \mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} \mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]\}, \quad \delta = \lambda + \zeta(1 - \lambda) \quad (2.10)$$

and

$$\mathbb{C}_{1-\delta}^{\delta}[\varpi_1, \varpi_2] = \{\mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2], \mathcal{D}_{\varpi_1^+}^{\delta; \vartheta} \mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]\}. \quad (2.11)$$

Since $\mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} = \mathcal{I}_{\varpi_1^+}^{\zeta(1-\lambda), \vartheta} \mathcal{D}_{\varpi_1^+}^{\delta; \vartheta}$, therefore, we have $\mathbb{C}_{1-\delta}^{\delta}[\varpi_1, \varpi_2] \subset \mathbb{C}_{1-\delta}^{\lambda, \zeta}[\varpi_1, \varpi_2]$.

Theorem 2.1. ([25]) For $\wp \geq \varpi_1, \vartheta \in (0, 1], \Re(\lambda), \Re(\zeta) > 0$. If $\mathcal{G} \in \mathbb{C}([\varpi_1, \varpi_2], \mathbb{R})$, then

$$\mathcal{I}_{\varpi_1^+}^{\lambda; \vartheta}(\mathcal{I}_{\varpi_1^+}^{\zeta; \vartheta} \mathcal{G})(\wp) = \mathcal{I}_{\varpi_1^+}^{\zeta; \vartheta}(\mathcal{I}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{G})(\wp) = (\mathcal{I}_{\varpi_1^+}^{\lambda+\zeta} \mathcal{G})(\wp).$$

Theorem 2.2. ([25]) For $\wp \geq \varpi_1, \vartheta \in (0, 1]$ and $\Re(\lambda) > 0$ and let $\mathcal{G} \in L_1([\varpi_1, \varpi_2])$, then

$$\mathcal{D}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{I}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{G}(\wp) = \mathcal{G}(\wp), \quad n = [\Re(\lambda)] + 1.$$

Lemma 2.1. ([25]) For $\lambda, \varsigma \in \mathbb{C}$ such that $\Re(\lambda) \geq 0$ and $\Re(\varsigma) > 0$. Then for any $\vartheta \in (0, 1]$ we have

$$(a) \quad (\mathcal{I}_{\varpi_1^+}^{\lambda; \vartheta} e^{\frac{\vartheta-1}{\vartheta} \ell} (\ell - \varpi_1)^{\varsigma-1})(\wp) = \frac{\Gamma(\varsigma)}{\vartheta^{\lambda} \Gamma(\varsigma + \lambda)} e^{\frac{\vartheta-1}{\vartheta} \wp} (\wp - \varpi_1)^{\varsigma + \lambda - 1},$$

$$(b) \quad (\mathcal{D}_{\varpi_1^+}^{\lambda; \vartheta} e^{\frac{\vartheta-1}{\vartheta} \ell} (\ell - \varpi_1)^{\varsigma-1})(\wp) = \frac{\vartheta^{\lambda} \Gamma(\varsigma)}{\Gamma(\varsigma - \lambda)} e^{\frac{\vartheta-1}{\vartheta} \wp} (\wp - \varpi_1)^{\varsigma - \lambda - 1}.$$

Lemma 2.2. ([30]) For $\lambda \in (0, 1), \vartheta \in (0, 1], \zeta \in (0, 1)$ and $\delta = \lambda + \zeta - \lambda\zeta$. If $\mathcal{G} \in \mathbb{C}_{1-\delta}^{\delta}[\varpi_1, \varpi_2]$, then

$$\mathcal{I}_{\varpi_1^+}^{\delta; \vartheta} \mathcal{D}_{\varpi_1^+}^{\delta; \vartheta} \mathcal{G} = \mathcal{I}_{\varpi_1^+}^{\delta; \vartheta} \mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} \mathcal{G}$$

and

$$\mathcal{D}_{\varpi_1^+}^{\delta; \vartheta} \mathcal{I}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{G} = \mathcal{D}_{\varpi_1^+}^{\zeta(1-\lambda); \vartheta} \mathcal{G}.$$

Lemma 2.3. ([30]) For $y \in (\varpi_1, \varpi_2], \lambda \in (0, 1), \vartheta \in (0, 1), \zeta \in [0, 1]$ and $\delta \in (0, 1)$. If $\mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$ and $\mathcal{I}_{\varpi_1^+}^{1-\delta; \vartheta} \mathcal{G}$, then

$$\mathcal{I}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} \mathcal{G}(y) = \mathcal{G}(y) - e^{\frac{\vartheta-1}{\vartheta}(y-\varpi_1)} \frac{(y - \varpi_1)^{\delta-1}}{\vartheta^{\delta-1} \Gamma(\delta)} (\mathcal{I}_{\varpi_1^+}^{1-\delta; \vartheta})(\varpi_1^+).$$

Lemma 2.4. ([30]) For $\delta \in [0, 1), \vartheta \in (0, 1]$ and $g_1 \in \mathbb{C}_{\delta}$. If $\mathcal{G} \in \mathbb{C}_{\delta}[\varpi_1, \varpi_2]$, then

$$\mathcal{I}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{G}(\varpi_1) = \lim_{y \rightarrow \varpi_1^+} \mathcal{I}_{\varpi_1^+}^{\lambda; \vartheta} \mathcal{G}(y) = 0, \quad \delta \in [0, \lambda).$$

Lemma 2.5. ([30]) For $\lambda \in (0, 1), \zeta \in [0, 1]$ and $\delta = \lambda + \zeta - \lambda\zeta$ and let $\mathcal{G} : (\varpi_1, \varpi_2] \times \mathbb{R} \mapsto \mathbb{R}$ be a mapping such that $\mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$ for any $y \in \mathbb{C}_{1-\delta}^{\delta}[\varpi_1, \varpi_2]$, then y satisfies problem (1.1) if and only if y satisfies the Volterra integral equation

$$y(\wp) = \frac{(\wp - \varpi_1)^{\delta-1} e^{\frac{\vartheta-1}{\vartheta}(\wp-\varpi_1)} \mathcal{I}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_1^+)}{\vartheta^{\delta-1} \Gamma(\delta)} + \frac{1}{\vartheta^{\lambda} \Gamma(\lambda)} \int_{\varpi_1}^{\wp} e^{\frac{\vartheta-1}{\vartheta}(\wp-s)} (\wp - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell. \quad (2.12)$$

3. Existence of solution

This section consists of the existence of solution to BVP (1.1) in $\mathbb{C}_{1-\delta}^{\lambda, \zeta; \vartheta}[\varpi_1, \varpi_2]$.

Lemma 3.1. For $\lambda \in (0, 1)$, $\zeta \in [0, 1]$, where $\delta = \lambda + \zeta - \lambda\zeta$ and suppose there be a function $\mathcal{G} : (\varpi_1, \varpi_2] \times \mathbb{R} \rightarrow \mathbb{R}$ such that $\mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$ for any $y \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$. If $y \in \mathbb{C}_{1-\delta}^\delta[\varpi_1, \varpi_2]$, then y fulfills BVP (1.1) if and only if y holds the following identity

$$\begin{aligned} y(\varphi) &= \frac{e_j}{m_1 + m_2} \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi_1)}(\varphi - \varpi_1)^{\delta-1}}{\vartheta^\delta \Gamma(\delta)} - \frac{m_2}{m_1 + m_2} \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi_1)}(\varphi - \varpi_1)^{\delta-1}}{\vartheta^\delta \Gamma(\delta)} \\ &\quad \times \frac{1}{\vartheta^{1-\delta+\lambda} \Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)}(\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell \\ &\quad + \frac{1}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)}(\varphi - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell. \end{aligned} \quad (3.1)$$

Proof. By means of Lemma 2.5 and utilizing the solution of (1.1) can be expressed as

$$y(\varphi) = \frac{\mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_1^+)}{\vartheta^{\delta-1} \Gamma(\delta)} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi_1)}(\varphi - \varpi_1)^{\delta-1} + \frac{1}{\vartheta^\delta \Gamma(\delta)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)}(\varphi - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell, \quad \varphi > \varpi_1. \quad (3.2)$$

Employing $\mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta}$ on (3.2) and applying the limit $\varphi \rightarrow \varpi_2^-$, we find

$$\mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_2^-) = \mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_1^+) + \frac{1}{\vartheta^{1-\delta+\lambda} \Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)}(\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell. \quad (3.3)$$

Again, employing $\mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta}$ on (3.3), we have

$$\begin{aligned} \mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_2^-) &= \mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_1^+) + \frac{1}{\vartheta^{1-\delta+\lambda} \Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)}(\varphi - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell \\ &= \mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_1^+) + \mathcal{J}_{\varpi_1^+}^{1-\zeta(1-\lambda); \vartheta} \mathcal{G}(\varphi, y(\varphi)). \end{aligned} \quad (3.4)$$

Applying limit $\varphi \mapsto \varpi_1^+$ and utilizing Lemma 2.4 having $1-\delta < 1-\zeta(1-\lambda)$, yields

$$\mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_1^+) = \mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_1^+), \quad (3.5)$$

thus

$$\mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_2^-) = \mathcal{J}_{\varpi_1^+}^{1-\delta; \vartheta} y(\varpi_1^+) + \frac{1}{\vartheta^{1-\delta+\lambda} \Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)}(\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell. \quad (3.6)$$

From boundary condition (1.1), we have

$$\mathcal{J}_{\varpi_1^+}^{1-\delta;\vartheta} y(\varpi_2^-) = \frac{e_J}{m_2} - \frac{m_1}{m_2} \mathcal{J}_{\varpi_1^+}^{1-\delta;\vartheta} y(\varpi_1^+). \quad (3.7)$$

From (3.5) and (3.6), and utilizing (3.4), we have

$$\mathcal{J}_{\varpi_1^+}^{1-\delta;\vartheta} y(\varpi_1^+) = \frac{m_2}{m_1 + m_2} \left(\frac{e_J}{m_2} - \frac{1}{\vartheta^{1-\delta+\lambda}\Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell \right). \quad (3.8)$$

Setting (3.2) in (3.8), one can find

$$\begin{aligned} y(\varphi) &= \frac{e_J}{m_1 + m_2} \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi_1)} (\varphi - \varpi_1)^{\delta-1}}{\vartheta^{\delta-1}\Gamma(\delta)} - \frac{m_2}{m_1 + m_2} \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi_1)} (\varphi - \varpi_1)^{\delta-1}}{\vartheta^{\delta-1}\Gamma(\delta)} \\ &\quad \times \frac{1}{\vartheta^{1-\delta+\lambda}\Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell \\ &\quad + \frac{1}{\vartheta^\lambda\Gamma(\lambda)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} (\varphi - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell. \end{aligned} \quad (3.9)$$

Conversely, employing $\mathcal{J}_{\varpi_1^+}^{1-\delta;\vartheta}$ on (3.1), utilizing Lemmas 2.1 and 2.2, and simple computations yields

$$\begin{aligned} &\mathcal{J}_{\varpi_1^+}^{1-\delta;\vartheta} m_1 y(\varpi_1^+) + \mathcal{J}_{\varpi_1^+}^{1-\delta;\vartheta} m_2 y(\varpi_2^-) \\ &= \frac{m_1 m_2}{m_1 + m_2} \left(\frac{e_J}{m_2} - \frac{1}{\vartheta^{1-\delta+\lambda}\Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell \right) \\ &\quad + \frac{m_2^2}{m_1 + m_2} \left(\frac{e_J}{m_2} - \frac{1}{\vartheta^{1-\delta+\lambda}\Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell \right) \\ &\quad + \frac{m_2}{\vartheta^{1-\delta+\lambda}\Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell \\ &= e_J, \end{aligned} \quad (3.10)$$

This shows that $y(\varphi)$ satisfies boundary condition (1.1).

Furthermore, employing $\mathcal{D}_{\varpi_1^+}^{\delta;\vartheta}$ on (3.1) and applying Lemmas 2.1 and 2.2, we have

$$\mathcal{D}_{\varpi_1^+}^{\delta;\vartheta} y(\varphi) = \mathcal{D}_{\varpi_1^+}^{\zeta(1-\lambda);\vartheta} \mathcal{G}(\varphi, y(\varphi)). \quad (3.11)$$

Since $y \in \mathbb{C}_{1-\delta}^{\delta;\vartheta}[\varpi_1, \varpi_2]$ and in view of definition of $\mathbb{C}_{1-\delta}^{\delta;\vartheta}[\varpi_1, \varpi_2]$, we have $\mathcal{D}_{\varpi_1^+}^{\delta} y \in \mathbb{C}_{n-\delta}^{\vartheta}[\varpi_1, \varpi_2]$, thus, $\mathcal{D}_{\varpi_1^+}^{\zeta(1-\lambda);\vartheta} \mathcal{G} = \mathcal{DI}_{\varpi_1^+}^{1-\zeta(1-\lambda)} \mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$.

For $\mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$, it is noting that $\mathcal{I}_{\varpi_1^+}^{1-\zeta(1-\lambda)} \mathcal{G} \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$. So, \mathcal{G} and $\mathcal{I}_{\varpi_1^+}^{1-\zeta(1-\lambda); \vartheta} \mathcal{G}$ holds the assumptions of Lemma 2.3. Now, employing $\mathcal{I}_{\varpi_1^+}^{\zeta(1-\lambda); \vartheta}$ on (3.11), we have

$$\mathcal{I}_{\varpi_1^+}^{\zeta(1-\lambda); \vartheta} \mathcal{D}_{\varpi_1^+}^{\delta; \vartheta} y(\varphi) = \mathcal{I}_{\varpi_1^+}^{\zeta(1-\lambda); \vartheta} \mathcal{D}_{\varpi_1^+}^{\zeta(1-\lambda); \vartheta} \mathcal{G}(\varphi, y(\varphi)). \tag{3.12}$$

Considering (2.8), (3.11) and Lemma 2.3, we have

$$\mathcal{I}_{\varpi_1^+}^{\delta; \vartheta} \mathcal{D}_{\varpi_1^+}^{\delta; \vartheta} y(\varphi) = \mathcal{G}(\varphi, y(\varphi)) - \frac{\mathcal{I}_{\varpi_1^+}^{1-\zeta(1-\lambda); \vartheta} \mathcal{G}(\varpi_1, y(\varpi_1))}{\vartheta^{\zeta(1-\lambda)} \Gamma(\zeta(1-\lambda))} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi_1)} (\varphi - \varpi_1)^{\zeta(1-\lambda)-1}, \forall \varphi \in (\varpi_1, \varpi_2]. \tag{3.13}$$

By Lemma 2.4, we have $\mathcal{I}_{\varpi_1^+}^{1-\zeta(n-\lambda)} \mathcal{G}(\varpi_1, y(\varpi_1)) = 0$. Thus, we have $\mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} y(\varphi) = \mathcal{G}(\varphi, y(\varphi))$. Hence, this completes the proof. \square

Let us evoke some essential assumptions which are required to prove the existence of solutions for the problem mentioned.

(A₁) Let a function $\mathcal{G} : (\varpi_1, \varpi_2) \times \mathbb{R} \mapsto \mathbb{R}$ with $\mathcal{G}(\cdot, y(\cdot)) \in \mathbb{C}_{1-\delta}^{\zeta(1-\lambda); \vartheta}[\varpi_1, \varpi_2]$. For any $y \in \mathbb{C}_{1-\delta}^{\vartheta}[\varpi_1, \varpi_2]$ and there exist two constants \mathcal{M}_1, m such that

$$|\mathcal{G}(\varphi_1, y)| \leq \mathcal{M}_1 (1 + m \|y\|_{\mathbb{C}_{1-\delta}^{\vartheta}}). \tag{3.14}$$

(A₂) The inequality

$$\mathcal{G} := \frac{m \mathcal{M}_1 \Gamma(\delta)}{\vartheta^{\lambda} \Gamma(\lambda + 1)} [(\varpi_2 - \varpi_1)^{\lambda} + (\varpi_2 - \varpi_1)^{\lambda+1-\delta}] < 1 \tag{3.15}$$

holds.

Now we are in a position to show the existence results for the BVP (1.1) by employing Schauder’s fixed point theorem (see [36]).

Theorem 3.1. *Suppose that the assumptions (A₁) and (A₂) fulfills. Then by Hilfer-BVP (1.1) has at least one solution in $\mathbb{C}_{1-\delta}^{\delta; \vartheta}[\varpi_1, \varpi_2] \subset \mathbb{C}_{1-\delta}^{\lambda, \zeta; \vartheta}[\varpi_1, \varpi_2]$.*

Proof. Defining an operator $\mathbf{T} : \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2] \mapsto \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$ by

$$\begin{aligned} (\mathbf{T}y)(\varphi) &= \frac{e_J}{m_1 + m_2} \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi_1)} (\varphi - \varpi_1)^{\delta-1}}{\vartheta^{\delta-1} \Gamma(\delta)} - \frac{m_2}{m_1 + m_2} \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi_1)} (\varphi - \varpi_1)^{\delta-1}}{\vartheta^{\delta-1} \Gamma(\delta)} \\ &\quad \times \frac{1}{\vartheta^{1-\delta+\lambda} \Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{G}(\ell, y(\ell)) d\ell \\ &\quad + \frac{1}{\vartheta^{\lambda} \Gamma(\lambda)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} (\varphi - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell. \end{aligned} \tag{3.16}$$

Assume that $\mathbb{B}_{\varrho} = \{y \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2] : \|y\|_{\mathbb{C}_{1-\delta}} \leq \varrho\}$ having $\varrho \geq \frac{\Omega}{1-\mathcal{G}}$, for $\mathcal{G} < 1$, we have

$$\Omega := \frac{e_J}{(m_1 + m_2) \vartheta^{\delta-1} \Gamma(\delta)} + \left| \frac{m_2}{m_1 + m_2} \right| \frac{1}{\vartheta^{\delta-1} \Gamma(\delta)}$$

$$\times \frac{\mathcal{M}_1}{\vartheta^{1-\delta+\lambda}} \left[\frac{(\varpi_2 - \varpi_1)^{\lambda+1-\delta}}{\Gamma(\lambda - \delta + 2)} + \frac{(\varpi_2 - \varpi_1)^{2\lambda-\delta+1}}{\Gamma(\lambda + 1)} \right]. \quad (3.17)$$

The proof will be demonstrated by the accompanying three steps:

Case 1. We will prove that $\mathbf{T}(\mathbb{B}_\varrho) \subset \mathbb{B}_\varrho$. Utilizing assumption (A_2) , we have

$$\begin{aligned} & \left| (\mathbf{T}y)(\varphi)(\varphi - \varpi_1)^{1-\delta} \right| \\ & \leq \left| \frac{e_j}{(m_1 + m_2)\vartheta^{\delta-1}\Gamma(\delta)} \right| + \left| \frac{m_2}{m_1 + m_2} \frac{1}{\vartheta^{\delta-1}\Gamma(\delta)} \right| \\ & \quad \times \frac{1}{\vartheta^{1-\delta+\lambda}\Gamma(1 - \delta + \lambda)} \int_{\varpi_1}^{\varpi_2} \left| e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} \right| (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{M}_1(1 + m|y|) d\ell \\ & \quad + \frac{(\varphi - \varpi_1)^{1-\delta}}{\vartheta^\lambda\Gamma(\lambda)} \int_{\varpi_1}^{\varphi} \left| e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} \right| (\varphi - \ell)^{\lambda-1} \mathcal{M}_1(1 + m|y|) d\ell \\ & \leq \frac{e_j}{(m_1 + m_2)\vartheta^{\delta-1}\Gamma(\delta)} + \left| \frac{m_2}{m_1 + m_2} \right| \frac{1}{\vartheta^{\delta-1}\Gamma(\delta)} \\ & \quad \times \frac{1}{\vartheta^{1-\delta+\lambda}\Gamma(1 - \delta + \lambda)} \int_{\varpi_1}^{\varpi_2} \left| e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} \right| (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{M}_1(1 + m\|y\|_{\mathbb{C}_{1-\delta}}) d\ell \\ & \quad + \frac{(\varphi - \varpi_1)^{1-\delta}}{\vartheta^\lambda\Gamma(\lambda)} \int_{\varpi_1}^{\varphi} \left| e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} \right| (\varphi - \ell)^{\lambda-1} \mathcal{M}_1(1 + m\|y\|_{\mathbb{C}_{1-\delta}}) d\ell. \end{aligned} \quad (3.18)$$

Since $\left| e^{\frac{\vartheta-1}{\vartheta}\varphi} \right| < 1$. Observe that, for any $y \in \mathbb{B}_\varrho$, and for every $\varphi \in (\varpi_1, \varpi_2]$, we have

$$\begin{aligned} & \frac{1}{\vartheta^{1-\delta+\lambda}\Gamma(1 - \delta + \lambda)} \int_{\varpi_1}^{\varpi_2} \left| e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} \right| (\varpi_2 - \ell)^{\lambda-\delta} \mathcal{M}_1(1 + m\|y\|_{\mathbb{C}_{1-\delta}}) d\ell \\ & \leq \frac{\mathcal{M}_1(\varpi_2 - \varpi_1)^\lambda}{\vartheta^{1-\delta+\lambda}} \left[\frac{(\varpi_2 - \varpi_1)^{1-\delta}}{\Gamma(\lambda - \delta + 2)} + \frac{m\varrho\Gamma(\delta)}{\Gamma(\lambda + 1)} \right], \text{ since } \left| e^{\frac{\vartheta-1}{\vartheta}\varphi} \right| < 1 \end{aligned} \quad (3.19)$$

and

$$\begin{aligned} & \frac{(\varphi - \varpi_1)^{1-\delta}}{\vartheta^\lambda\Gamma(\lambda)} \int_{\varpi_1}^{\varphi} \left| e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} \right| (\varphi - \ell)^{\lambda-1} \mathcal{M}_1(1 + m\|y\|_{\mathbb{C}_{1-\delta}}) d\ell \\ & \leq \frac{\mathcal{M}_1(\varphi - \varpi_1)^{\lambda-\delta+1}}{\vartheta^\lambda} \left[\frac{(\varphi - \varpi_1)^\lambda}{\Gamma(\lambda + 1)} + \frac{m\varrho\Gamma(\delta)}{\Gamma(\lambda + 1)} \right]. \end{aligned} \quad (3.20)$$

Therefore, we get

$$\begin{aligned} & \left| (\mathbf{T}y)(\varphi)(\varphi - \varpi_1)^{1-\delta} \right| \\ & \leq \frac{e_j}{(m_1 + m_2)\vartheta^{\delta-1}\Gamma(\delta)} + \left| \frac{m_2}{m_1 + m_2} \right| \frac{1}{\vartheta^{\delta-1}\Gamma(\delta)} \end{aligned}$$

$$\begin{aligned} & \times \frac{\mathcal{M}_1(\varpi_2 - \varpi_1)^\lambda}{\vartheta^{1-\delta+\lambda}} \left[\frac{(\varpi_2 - \varpi_1)^{1-\delta}}{\Gamma(\lambda - \delta + 2)} + \frac{m_\varrho \Gamma(\delta)}{\Gamma(\lambda + 1)} \right] \\ & + \frac{\mathcal{M}_1(\varphi - \varpi_1)^{\lambda-\delta+1}}{\vartheta^\lambda} \left[\frac{(\varphi - \varpi_1)^\lambda}{\Gamma(\lambda + 1)} + \frac{m_\varrho \Gamma(\delta)}{\Gamma(\lambda + 1)} \right], \end{aligned} \quad (3.21)$$

which leads to

$$\begin{aligned} \|\mathbf{T}y\|_{\mathbb{C}_{1-\delta}} & \leq \frac{e_j}{(m_1 + m_2)\vartheta^{\delta-1}\Gamma(\delta)} + \left| \frac{m_2}{m_1 + m_2} \right| \frac{1}{\vartheta^{\delta-1}\Gamma(\delta)} \\ & \times \frac{\mathcal{M}_1}{\vartheta^{1-\delta+\lambda}} \left[\frac{(\varpi_2 - \varpi_1)^{\lambda+1-\delta}}{\Gamma(\lambda - \delta + 2)} + \frac{(\varpi_2 - \varpi_1)^{2\lambda-\delta+1}}{\Gamma(\lambda + 1)} \right] \\ & + \frac{\mathcal{M}_1 m_\varrho \Gamma(\delta)}{\vartheta^\lambda \Gamma(\lambda + 1)} \left[(\varphi - \varpi_1)^\lambda + (\varpi_2 - \varpi_1)^{\lambda+1-\delta} \right]. \end{aligned} \quad (3.22)$$

of assumption (A_2) , we conclude that $\|\mathbf{T}y\|_{\mathbb{C}_{1-\delta}} \leq \mathcal{G}\varrho + (1 - \mathcal{G})\varrho = \varrho$. Therefore, $\mathbf{T}(\mathbb{B}_\varrho) \subset \mathbb{B}_\varrho$.

Next we will prove that \mathbf{T} is completely continuous.

Case 2. We prove that the operator \mathbf{T} is completely continuous.

Assume that $\{\bar{z}_n\}$ is a sequence such that $\bar{z}_n \mapsto \bar{z}$ in \mathbb{B}_ϱ as $n \mapsto \infty$. Then for every $\varphi \in (\varpi_1, \varpi_2]$, we have

$$\begin{aligned} & \left| \left((\mathbf{T}\bar{z}_n)(\varphi) - (\mathbf{T}\bar{z})(\varphi) \right) (\varphi - \varpi_1)^{1-\delta} \right| \\ & = \left| \frac{m_2}{m_1 + m_2} \right| \frac{1}{\vartheta^{\delta-1}\Gamma(\delta)} \\ & \times \frac{1}{\vartheta^{1-\delta+\lambda}\Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} \left| e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{\lambda-\delta} \left| \mathcal{G}(\ell, \bar{z}_n(\ell)) - \mathcal{G}(\ell, \bar{z}(\ell)) \right| d\ell \right. \\ & \left. + \frac{(\varphi - \varpi_1)^{1-\delta}}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\varphi} \left| e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)} (\varphi - \ell)^{\lambda-1} \left| \mathcal{G}(\ell, \bar{z}_n(\ell)) - \mathcal{G}(\ell, \bar{z}(\ell)) \right| d\ell \right. \right. \\ & \leq \left| \frac{m_2}{m_1 + m_2} \right| \frac{1}{\vartheta^\lambda \Gamma(\lambda + 1)} (\varpi_2 - \varpi_1)^\lambda \left\| \mathcal{G}(\cdot, \bar{z}_n(\cdot)) - \mathcal{G}(\cdot, \bar{z}(\cdot)) \right\|_{\mathbb{C}_{1-\delta}} \\ & \quad + \frac{\Gamma(\delta)(\varphi - \varpi_1)^{1-\delta+\lambda}}{\vartheta^{\lambda-\delta}\Gamma(\lambda - \delta)} (\varphi - \varpi_1)^{1-\delta+\lambda} \left\| \mathcal{G}(\cdot, \bar{z}_n(\cdot)) - \mathcal{G}(\cdot, \bar{z}(\cdot)) \right\|_{\mathbb{C}_{1-\delta}}. \end{aligned} \quad (3.23)$$

Since $\left| e^{\frac{\vartheta-1}{\vartheta}\varphi} \right| < 1$ and \mathcal{G} is continuous on $(\varpi_1, \varpi_2]$ and $\bar{z}_n \mapsto \bar{z}$, then

$$\left\| (\mathbf{T}\bar{z}_n - \mathbf{T}\bar{z}) \right\|_{\mathbb{C}_{1-\delta}} \rightarrow 0 \text{ as } n \rightarrow \infty, \quad (3.24)$$

which shows that operator \mathbf{T} is continuous on \mathbb{B}_ϱ .

Case 3. We show that $\mathbf{T}(\mathbb{B}_\varrho)$ is relatively compact. In case 1, we have $\mathbf{T}(\mathbb{B}_\varrho) \subset \mathbb{B}_\varrho$. It is observed that $\mathbf{T}(\mathbb{B}_\varrho)$ is uniformly bounded. To show operator \mathbf{T} is equi-continuous on \mathbb{B}_ϱ . In fact, for any $\varpi_1 < \varphi_1 < \varphi_2 < \varpi_2$ and $\bar{z} \in \mathbb{B}_\varrho$, we have

$$\left| (\varphi_2 - \varpi_1)^{1-\delta} (\mathbf{T}y)(\varphi_2) - (\varphi_1 - \varpi_1)^{1-\delta} (\mathbf{T}y)(\varphi) \right|$$

$$\begin{aligned}
&\leq \frac{|(\varphi_2 - \varpi_1)^{n-\kappa} - (\varphi_1 - \varpi_1)^{n-\kappa}|}{\vartheta^{\delta-1}\Gamma(\delta)} \frac{e_J}{m_1 + m_2} + \left| \frac{m_2}{m_1 + m_2} \right| \frac{|(\varphi_2 - \varpi_1)^{n-\kappa} - (\varphi_1 - \varpi_1)^{n-\kappa}|}{\vartheta^{\delta-1}\Gamma(\delta)} \\
&\times \frac{1}{\vartheta^{\lambda-\delta+1}\Gamma(\lambda - \delta + 1)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{\lambda-\delta} |\mathcal{G}(\ell, y(\ell))| d\ell \\
&\quad + \frac{1}{\vartheta^\lambda \Gamma(\lambda)} |(\varphi_2 - \varpi_1)^{1-\delta} \int_{\varpi_1}^{\varphi_2} e^{\frac{\vartheta-1}{\vartheta}(\varphi_2-\ell)} (\varphi_2 - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell \\
&\quad - (\varphi_1 - \varpi_1)^{1-\delta} \int_{\varpi_1}^{\varphi_1} e^{\frac{\vartheta-1}{\vartheta}(\varphi_1-\ell)} (\varphi_1 - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell| \\
&\leq \frac{|(\varphi_2 - \varpi_1)^{n-\kappa} - (\varphi_1 - \varpi_1)^{n-\kappa}|}{\vartheta^{\delta-1}\Gamma(\delta)} \left[\frac{e_J}{m_1 + m_2} + \left| \frac{m_2}{m_1 + m_2} \frac{\|\mathcal{G}\|_{\mathbb{C}_{1-\delta}}}{\vartheta^{1-\delta+\lambda}\Gamma(1-\delta+\lambda)} \int_{\varpi_1}^{\varpi_2} (\varpi_2 - \ell)^{\lambda-\delta} (\ell - \varpi_1)^{\delta-1} d\ell \right] \right. \\
&\quad + \frac{\|\mathcal{G}\|_{\mathbb{C}_{1-\delta}}}{\vartheta^\lambda \Gamma(\lambda)} |(\varphi_2 - \varpi_1)^{1-\delta} \int_{\varpi_1}^{\varphi_2} (\varphi_2 - \ell)^{\lambda-1} (\ell - \varpi_1)^{\delta-1} d\ell \\
&\quad - (\varphi_1 - \varpi_1)^{1-\delta} \int_{\varpi_1}^{\varphi_1} (\varphi_1 - \ell)^{\lambda-1} (\ell - \varpi_1)^{\delta-1} d\ell| \quad (\text{since } |e^{\frac{\vartheta-1}{\vartheta}\varphi}| < 1) \\
&\leq \frac{|(\varphi_2 - \varpi_1)^{n-\kappa} - (\varphi_1 - \varpi_1)^{n-\kappa}|}{\vartheta^{\delta-1}\Gamma(\delta)} \left[\frac{e_J}{m_1 + m_2} + \left| \frac{m_2}{m_1 + m_2} \frac{\Gamma(\delta)}{\vartheta^\lambda \Gamma(\lambda + 1)} (\varpi_2 - \varpi_1)^\lambda \|\mathcal{G}\|_{\mathbb{C}_{1-\delta}} \right] \right. \\
&\quad \left. + \frac{\|\mathcal{G}\|_{\mathbb{C}_{1-\delta}}}{\vartheta^\lambda \Gamma(\lambda)} \mathfrak{B}(\delta - n + 1, \lambda) |(\varphi_2 - \varpi_1)^\lambda - (\varphi_1 - \varpi_1)^\lambda|, \tag{3.25}
\end{aligned}$$

which approaches to zero as $\varphi_2 \rightarrow \varphi_1$, independent of $y \in \varrho$, where $\mathfrak{B}(\cdot, \cdot)$ denotes the Euler Beta function.

Therefore, we deduce that $\mathbf{T}(\mathbb{B}_\varrho)$ is equicontinuous on \mathbb{B}_ϱ , that leads to the relatively compactness. As a result, we conclude that by Arzela-Ascoli theorem, the defined operator $\mathbf{T} : \mathbb{B}_\varrho \mapsto \mathbb{B}_\varrho$ is completely continuous operator.

By Schauder's fixed point theorem, there exists at least one fixed point y of \mathbf{T} in $\mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$. This fixed point y is the solution of (1.1) in $\mathbb{C}_{1-\delta}^{\delta, \vartheta}$, and this completes the proof. \square

Now we present another existence result via Schaefer fixed point theorem. For this, we need the following assumption.

(A₃) Suppose a function $\mathcal{G} : (\varpi_1, \varpi_2] \times \mathbb{R} \mapsto \mathbb{R}$ such that $\mathcal{G}(\cdot, y(\cdot)) \in \mathbb{C}_{1-\delta}^{\zeta(1-\lambda); \vartheta}[\varpi_1, \varpi_2]$ for any $y \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$ and there exist a mapping $\eta(\varphi) \in \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$ such that

$$|\mathcal{G}(\varphi, y)| \leq \eta(\varphi), \quad \forall \varphi \in (\varpi_1, \varpi_2], y \in \mathbb{R}. \tag{3.26}$$

Theorem 3.2. *Suppose that assumption (A₃) satisfies. then Hilfer-BVP (1.1) has at least one solution in $\mathbb{C}_{1-\delta}^\delta \subset \mathbb{C}_{1-\delta}^{\lambda, \zeta; \vartheta}[\varpi_1, \varpi_2]$.*

Proof. For the proof of Theorem 3.2, one can adopt the same technique as we did in Theorem 3.1 and easily prove that the operator $\mathbf{T} : \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2] \mapsto \mathbb{C}_{1-\delta}[\varpi_1, \varpi_2]$ stated in (3.16) is completely continuous. Now we show that

$$\Delta = \{y \in \mathbb{C}_{n-\delta}[\varpi_1, \varpi_2] : y = \sigma \mathbf{T}y, \text{ for some } \sigma \in (0, 1)\} \quad (3.27)$$

is bounded set. Assume that $y \in \Delta$ and $\sigma \in (0, 1)$ be such that $y = \sigma \mathbf{T}y$. By assumption (A₃) and (3.16), then for all $\wp \in [\varpi_1, \varpi_2]$, we have

$$\begin{aligned} & |\mathbf{T}y(\wp)(\wp - \varpi_1)^{n-\delta}| \\ & \leq \sum_{\kappa=1}^n \frac{e^{\frac{\wp-1}{\vartheta}(\wp-\varpi_1)}(\wp - \varpi_1)^{n-\kappa}}{\vartheta^{\delta-\kappa+1}\Gamma(\delta - \kappa + 1)} \frac{e_J}{m_1 + m_2} \\ & \quad + \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{e^{\frac{\wp-1}{\vartheta}(\wp-\varpi_1)}(\wp - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)\Gamma(n - \delta + \lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\wp-1}{\vartheta}(\varpi_2-\ell)}(\varpi_2 - \ell)^{n+\lambda-\delta-1} \eta(\ell) d\ell \\ & \quad + \frac{|(\wp - \varpi_1)^{n-\delta}|}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\wp} e^{\frac{\wp-1}{\vartheta}(\wp-\ell)}(\wp - \ell)^{\lambda-1} \eta(\ell) d\ell \\ & \leq \sum_{\kappa=1}^n \frac{(\wp - \varpi_1)^{n-\kappa}}{\vartheta^{\delta-\kappa+1}\Gamma(\delta - \kappa + 1)} \frac{e_J}{m_1 + m_2} \\ & \quad + \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\wp - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)\Gamma(n - \delta + \lambda)} \int_{\varpi_1}^{\varpi_2} (\varpi_2 - \ell)^{n+\lambda-\delta-1} (\ell - \varpi_1)^{\delta-n} \|\eta\|_{\mathbb{C}_{n-\delta}} d\ell \\ & \quad + \frac{|(\wp - \varpi_1)^{n-\delta}|}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\wp} (\wp - \ell)^{\lambda-1} (\ell - \varpi_1)^{\delta-n} \|\eta\|_{\mathbb{C}_{n-\delta}} d\ell. \end{aligned} \quad (3.28)$$

Since $|e^{\frac{\wp-1}{\vartheta}\wp}| < 1$, we have

$$\begin{aligned} & \|\mathbf{T}y\|_{\mathbb{C}_{n-\delta}} \\ & \leq \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{\delta-\kappa+1}\Gamma(\delta - \kappa + 1)} \frac{e_J}{m_1 + m_2} \\ & \quad + \left[\left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{-\kappa}}{\vartheta^{\delta-\kappa+1}} \frac{\Gamma(\lambda)}{\mathfrak{B}(\lambda, 1)} + \frac{\mathfrak{B}(\delta - n + 1, 1)}{\vartheta^\lambda (\varpi_2 - \varpi_1)^\delta} \right] (\varpi_2 - \varpi_1)^{n+\lambda} \|\eta\|_{\mathbb{C}_{n-\delta}} \\ & := \tau. \end{aligned} \quad (3.29)$$

Since $\sigma \in (0, 1)$, then $y < \mathbf{T}y$. The last inequality with (3.29) leads us to the conclusion that

$$\|y\|_{\mathbb{C}_{n-\delta}} < \|\mathbf{T}y\|_{\mathbb{C}_{n-\delta}} \leq \tau, \quad (3.30)$$

which proves that Δ is bounded. Utilizing Schaefer fixed point postulate, this completes the proof. \square

Our last result is the existence result for the problem (1.1) by using the Kransnoselskii’s fixed point theorem (see [37]), the following assumption is needed:

(A₄) Suppose that $\mathcal{G} : (\varpi_1, \varpi_2] \times \mathbb{R} \mapsto \mathbb{R}$ is a function such that $\mathcal{G}(\cdot, y(\cdot)) \in \mathbb{C}_{n-\delta}^{\zeta(n-\lambda); \vartheta}[\varpi_1, \varpi_2]$ for any $y \in \mathbb{C}_{n-\delta}[\varpi_1, \varpi_2]$ and there exists a constant $L > 0$ such that

$$|\mathcal{G}(\varphi, y) - \mathcal{G}(\varphi, \omega)| \leq L|y - \omega|, \quad \forall \varphi \in (\varpi_1, \varpi_2], y, \omega \in \mathbb{R}. \tag{3.31}$$

Also, we note the following assumption as follows: (A₅) the inequality

$$\begin{aligned} Q &:= \left[\left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)} + \frac{\mathfrak{B}(\delta - n, \lambda + 1)}{\vartheta^\delta\Gamma(\delta - n)} \right] \\ &\times \frac{\Gamma(\delta - n)(\varpi_2 - \varpi_1)^\lambda}{\vartheta^\lambda\mathfrak{B}(\delta - n, 1)\Gamma(\lambda + 1)} \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}} + \frac{e_J}{m_1 + m_2} \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)} L < 1 \end{aligned} \tag{3.32}$$

is hold.

Theorem 3.3. *Suppose that the assumptions (A₄) and (A₅) are satisfied. If*

$$\left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa+\lambda}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)} \frac{\Gamma(\delta - n + 1)}{\vartheta^\lambda\Gamma(\lambda + 1)} L < 1. \tag{3.33}$$

Then the Hilfer-BVP (1.1) has at least one solution in $\mathbb{C}_{n-\delta}^\delta[\varpi_1, \varpi_2] \subset \mathbb{C}_{n-\delta}^{\lambda, \zeta; \vartheta}$.

Proof. Considering the operator **T** stated in Theorem 3.1.

First, surmise the operator **T** into sum of two operators **T**₁ + **T**₂ as follows

$$\mathbf{T}_1 y(\varphi) = \frac{-m_2}{m_1 + m_2} \sum_{\kappa=1}^n \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi)}(\varphi - \varpi_1)^{\delta-\kappa}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)} \frac{1}{\Gamma(n - \delta + \lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)}(\varpi_2 - \ell)^{n-\delta+\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell \tag{3.34}$$

and

$$\mathbf{T}_2 y(\varphi) = \frac{e_J}{m_1 + m_2} \sum_{\kappa=1}^n \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi)}(\varphi - \varpi_1)^{\delta-\kappa}}{\vartheta^{\delta-\kappa+1}\Gamma(\delta - \kappa + 1)} + \frac{1}{\vartheta^\lambda\Gamma(\lambda)} \int_{\varpi_1}^{\varphi} e^{\frac{\vartheta-1}{\vartheta}(\varphi-\ell)}(\varphi - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell. \tag{3.35}$$

Setting $\tilde{\mathcal{G}} = \mathcal{G}(\ell, 0)$ and suppose the ball $\mathbb{B}_\epsilon = \{y \in \mathbb{C}_{n-\delta; \psi}[\varpi_1, \varpi_2] : \|y\|_{\mathbb{C}_{n-\delta; \psi}} \leq \epsilon\}$ having $\epsilon \geq \frac{\sigma}{1-Q}$, $Q < 1$, where

$$\begin{aligned} \sigma &= \left[\left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)} + \frac{\mathfrak{B}(\delta - n, \lambda + 1)}{\vartheta^\delta\Gamma(\delta - n)} \right] \\ &\times \frac{\Gamma(\delta - n)(\varpi_2 - \varpi_1)^\lambda}{\vartheta^\lambda\mathfrak{B}(\delta - n, 1)\Gamma(\lambda + 1)} L < 1 \end{aligned} \tag{3.36}$$

The proof will be done in three cases.

Case 1. We show that $\mathbf{T}_1 y + \mathbf{T}_1 \omega \in \mathbb{B}_\epsilon$ for every $y, \omega \in \mathbb{B}_\epsilon$.

Utilizing assumption (A_4) , then for every $y \in \mathbb{B}_\epsilon$ and $\wp \in (\varpi_1, \varpi_2]$, we have

$$\begin{aligned}
& |(\wp - \varpi_1)^{n-\delta} \mathbf{T}_1(y)| \\
& \leq \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{e^{\frac{\theta-1}{\theta}(\wp-\varpi)} (\wp - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \frac{1}{\Gamma(n - \delta + \lambda)} \\
& \quad \times \int_{\varpi_1}^{\varpi_2} e^{\frac{\theta-1}{\theta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{n-\delta+\lambda-1} \left[|\mathcal{G}(\ell, y(\ell)) - \mathcal{G}(\ell, 0)| + |\mathcal{G}(\ell, 0)| \right] d\ell \\
& \leq \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{e^{\frac{\theta-1}{\theta}(\wp-\varpi)} (\wp - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \frac{1}{\Gamma(n - \delta + \lambda)} \\
& \quad \times \int_{\varpi_1}^{\varpi_2} e^{\frac{\theta-1}{\theta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{n-\delta+\lambda-1} (\ell - \varpi_1)^{\delta-n} \left[L \|y\|_{\mathbb{C}_{n-\delta}} + \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}} \right] d\ell \\
& \leq \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\wp - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \frac{\Gamma(\delta - n + 1)}{\vartheta^\lambda \Gamma(\lambda + 1)} \left[L\epsilon + \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}} \right]. \tag{3.37}
\end{aligned}$$

Since $\left| e^{\frac{\theta-1}{\theta} \wp} \right| < 1$. Therefore, we get

$$\begin{aligned}
& \|\mathbf{T}_1 y\|_{\mathbb{C}_{n-\delta}} \\
& \leq \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa+\lambda}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \frac{\Gamma(\delta - n + 1)}{\vartheta^\lambda \Gamma(\lambda + 1)} \left[L\epsilon + \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}} \right]. \tag{3.38}
\end{aligned}$$

For operator \mathbf{T}_2 , we have

$$\begin{aligned}
& |(\wp - \varpi_1)^{n-\delta} \mathbf{T}_2 \omega(\wp)| \\
& \leq \left| \frac{e_j}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{e^{\frac{\theta-1}{\theta}(\wp-\varpi)} (\wp - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \\
& \quad \times \frac{(\wp - \varpi_1)^{n-\delta}}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\wp} e^{\frac{\theta-1}{\theta}(\wp-\ell)} (\wp - \ell)^{\lambda-1} \left[|\mathcal{G}(\ell, \omega(\ell)) - \mathcal{G}(\ell, 0)| + |\mathcal{G}(\ell, 0)| \right] d\ell \\
& \leq \left| \frac{e_j}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\wp - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \\
& \quad \times \frac{(\wp - \varpi_1)^{n-\delta}}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\wp} e^{\frac{\theta-1}{\theta}(\wp-\ell)} (\wp - \ell)^{\lambda-1} (\ell - \varpi_1)^{\delta-n} \left[L \|\omega\|_{\mathbb{C}_{n-\delta}} + \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}} \right] d\ell. \tag{3.39}
\end{aligned}$$

For every $\omega \in \mathbb{B}_\epsilon$ and $\wp \in (\varpi_1, \varpi_2]$, this shows

$$\begin{aligned}
& \|\mathbf{T}_1 \omega\|_{\mathbb{C}_{n-\delta}} \\
& \leq \left| \frac{e_j}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)}
\end{aligned}$$

$$\times \frac{(\varpi_2 - \varpi_1)^\lambda}{\vartheta^\lambda \Gamma(\delta - n + \lambda + 1)} [L\epsilon + \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}}]. \tag{3.40}$$

From (3.38), (3.40) and utilizing assumption (A₅) with (3.36), we find

$$\begin{aligned} & \| \mathbf{T}_1 y + \mathbf{T}_2 \omega \|_{\mathbb{C}_{n-\delta}} \\ & \leq \| \mathbf{T}_1 y \|_{\mathbb{C}_{n-\delta}} + \| \mathbf{T}_1 \omega \|_{\mathbb{C}_{n-\delta}} \\ & \leq \frac{(\varpi_2 - \varpi_1)^\lambda \Gamma(\delta - n + 1)}{\vartheta^\lambda \Gamma(\lambda + 1)} [L\epsilon + \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}}] \left[\left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} + \frac{\Gamma(\lambda + 1)}{\Gamma(\delta - n + \lambda + 1)} \right] \\ & \quad + \frac{e_j}{m_1 + m_2} \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \\ & \leq Q\epsilon + (1 - Q)\epsilon = \epsilon. \end{aligned} \tag{3.41}$$

Case 2. We prove that the operator \mathbf{T} is a contraction mapping on \mathbb{B}_ϱ .

For any $y, \omega \in \mathbb{B}_\varrho$, and for any $\varphi \in (\varpi_1, \varpi_2]$, then by supposition (A₄), we have

$$\begin{aligned} & |(\varphi - \varpi_1)^{n-\delta} \mathbf{T}_1 y(\varphi) - (\varphi - \varpi_1)^{n-\delta} \mathbf{T}_1 \omega(\varphi)| \\ & \leq \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi)} (\varphi - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \frac{1}{\Gamma(n - \delta + \lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{n-\delta+\lambda-1} [\mathcal{G}(\ell, y(\ell)) - \mathcal{G}(\ell, \omega(\ell))] d\ell \\ & \leq \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{e^{\frac{\vartheta-1}{\vartheta}(\varphi-\varpi)} (\varphi - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \frac{1}{\Gamma(n - \delta + \lambda)} \int_{\varpi_1}^{\varpi_2} e^{\frac{\vartheta-1}{\vartheta}(\varpi_2-\ell)} (\varpi_2 - \ell)^{n-\delta+\lambda-1} L |y(\ell) - \omega(\ell)| d\ell \\ & \leq \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varphi - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \frac{\Gamma(\delta - n + 1)}{\vartheta^\lambda \Gamma(\lambda + 1)} (\varpi_2 - \ell)^\lambda L \|y - \omega\|_{\mathbb{C}_{n-\delta}}. \end{aligned} \tag{3.42}$$

Since $|e^{\frac{\vartheta-1}{\vartheta}\varphi}| < 1$, this yields

$$\begin{aligned} & \| \mathbf{T}y - \mathbf{T}\omega \|_{\mathbb{C}_{n-\delta}} \\ & \leq \left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa+\lambda}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} \frac{\Gamma(\delta - n + 1)}{\vartheta^\lambda \Gamma(\lambda + 1)} L \|y - \omega\|_{\mathbb{C}_{n-\delta}}. \end{aligned} \tag{3.43}$$

Due to assumption (3.33), which shows that the operator \mathbf{T} is a contraction mapping.

Case 3: Now we show that the operator \mathbf{T}_2 is completely continuous on \mathbb{B}_ϵ .

From the continuity of \mathcal{G} , we deduce that the operator $\mathbf{T}_2 : \mathbb{B}_\epsilon \mapsto \mathbb{B}_\epsilon$ is continuous on \mathbb{B}_ϵ . Furthermore, we prove that for all $\epsilon > 0$ there exists some $\epsilon' > 0$ such that $\| \mathbf{T}_2 y \|_{\mathbb{C}_{n-\delta}} < \epsilon'$. In view of case 1, for $y \in \mathbb{B}_\epsilon$, we have that

$$\| \mathbf{T}_2 y \|_{\mathbb{C}_{n-\delta}} \leq \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{\delta-\kappa+1} \Gamma(\delta - \kappa + 1)} \frac{e_j}{m_1 + m_2} + \frac{\mathfrak{B}(\delta - n + 1, \lambda) (\varpi_2 - \varpi_1)^\lambda}{\vartheta^\lambda \Gamma(\lambda)} [L\epsilon + \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}}], \tag{3.44}$$

which is free of φ and y , so there exists

$$\epsilon' = \frac{e_j}{m_1 + m_2} \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda} \Gamma(\delta - \kappa + 1)} + \frac{\mathfrak{B}(\delta - n + 1, \lambda) (\varpi_2 - \varpi_1)^\lambda}{\vartheta^\lambda \Gamma(\lambda)} [L\epsilon + \|\tilde{\mathcal{G}}\|_{\mathbb{C}_{n-\delta}}] \tag{3.45}$$

such that $\|\mathbf{T}_2(y)\|_{\mathbb{C}_{n-\delta}} \leq \epsilon'$. Therefore, \mathbf{T}_2 is uniformly bounded set on \mathbb{B}_ϵ . Finally, to show that \mathbf{T}_2 is equicontinuous in \mathbb{B}_ϵ , for any $z \in \mathbb{B}_\epsilon$ and $\wp_1, \wp_2 \in (\varpi_1, \varpi_2]$ having $\wp_1 < \wp_2$, we have

$$\begin{aligned} & |(\wp_2 - \varpi_1)^{n-\delta} \mathbf{T}_2 y(\wp_2) - (\wp_1 - \varpi_1)^{n-\delta} \mathbf{T}_2 y(\wp_1)| \\ &= \frac{e_j}{m_1 + m_2} \left| \sum_{\kappa=1}^n \frac{(\wp_2 - \varpi_1)^{n-\kappa} - (\wp_1 - \varpi_1)^{n-\delta}}{\vartheta^{\delta-\kappa} \Gamma(\delta - \kappa + 1)} + \frac{(\wp_2 - \varpi_1)^{n-\delta}}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\wp_2} e^{\frac{\vartheta-1}{\vartheta}(\wp_2-\ell)} (\wp_2 - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell \right. \\ &\quad \left. - \frac{(\wp_1 - \varpi_1)^{n-\delta}}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\wp_1} e^{\frac{\vartheta-1}{\vartheta}(\wp_1-\ell)} (\wp_1 - \ell)^{\lambda-1} \mathcal{G}(\ell, y(\ell)) d\ell \right| \\ &\leq \frac{e_j}{m_1 + m_2} \sum_{\kappa=1}^n \frac{|(\wp_2 - \varpi_1)^{n-\kappa} - (\wp_1 - \varpi_1)^{n-\delta}|}{\vartheta^{\delta-\kappa} \Gamma(\delta - \kappa + 1)} \\ &\quad + \left| \frac{(\wp_2 - \varpi_1)^{n-\delta}}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\wp_2} e^{\frac{\vartheta-1}{\vartheta}(\wp_2-\ell)} (\wp_2 - \ell)^{\lambda-1} (\ell - \varpi_1)^{\delta-n} \|\mathcal{G}\|_{\mathbb{C}_{n-\delta, \psi[\varpi_1, \varpi_2]}} d\ell \right. \\ &\quad \left. - \frac{(\wp_1 - \varpi_1)^{n-\delta}}{\vartheta^\lambda \Gamma(\lambda)} \int_{\varpi_1}^{\wp_1} e^{\frac{\vartheta-1}{\vartheta}(\wp_1-\ell)} (\wp_1 - \ell)^{\lambda-1} (\ell - \varpi_1)^{\delta-n} \|\mathcal{G}\|_{\mathbb{C}_{n-\delta, \psi[\varpi_1, \varpi_2]}} d\ell \right| \\ &= \sum_{\kappa=1}^n \frac{e_j}{m_1 + m_2} \frac{|(\wp_2 - \varpi_1)^{n-\kappa} - (\wp_1 - \varpi_1)^{n-\delta}|}{\vartheta^{\delta-\kappa} \Gamma(\delta - \kappa + 1)} + \frac{\mathfrak{B}(\delta - n + 1)}{\vartheta^\lambda} \|\mathcal{G}\|_{\mathbb{C}_{n-\delta, \psi[\varpi_1, \varpi_2]}} |(\wp_2 - \varpi_1)^\lambda - (\wp_1 - \varpi_1)^\lambda|. \end{aligned} \tag{3.46}$$

Since $|e^{\frac{\vartheta-1}{\vartheta}\varphi}| < 1$. It is noting that the right hand side of the aforesaid variant is free of y . So,

$$|(\wp_2 - \varpi_1)^{n-\delta} \mathbf{T}_2 y(\wp_2) - (\wp_1 - \varpi_1)^{n-\delta} \mathbf{T}_2 y(\wp_1)| \mapsto 0, \text{ as } |\wp_2 - \wp_1| \mapsto 0. \tag{3.47}$$

This shows that \mathbf{T}_2 is equicontinuous on \mathbb{B}_ϵ . According to Arzela-Ascoli Theorem, observed that $(\mathbf{T}_2 \mathbb{B}_\epsilon)$ is relatively compact. By Kransnoselskii’s fixed point theorem, the problem (1.1) has at least one solution. \square

4. Examples

Consider the fractional differential equation with boundary condition which encompasses the Hilfer-GPF derivative of the form

$$\begin{cases} \mathcal{D}_{\varpi_1^+}^{\lambda, \zeta; \vartheta} y(\varphi) = \varphi^{-\frac{1}{6}} + \frac{\varphi^{5/6}}{16} \sin y(\varphi), \varphi \in \mathbb{J} = [0, 2], \lambda \in (0, 1), \zeta \in [0, 1] \\ \mathcal{I}_{\varpi_1^+}^{1-\delta; \vartheta} \left[\frac{1}{3} y(0^+) + \frac{2}{3} y(2^-) \right] = \frac{2}{5}, \lambda \leq \delta = \lambda + \zeta - \lambda \zeta, \end{cases} \tag{4.1}$$

By comparison (1.1) with (4.1), we have $\lambda = \frac{1}{2}, \zeta = \frac{1}{3}, \delta = \frac{2}{3}, m_1 = \frac{1}{3}, m_2 = \frac{2}{3}, \vartheta = 1$ and $e_1 = \frac{2}{5}$. It is clear that $\varphi^{\frac{1}{3}} \mathcal{G}(\varphi, y(\varphi)) = \varphi^{\frac{1}{6}} + \frac{\varphi^{7/6}}{16} \sin y(\varphi) \in \mathbb{C}([0, 2])$, So $\mathcal{G}(\varphi, y(\varphi)) \in \mathbb{C}_{\frac{1}{3}}$. Thus, it follows that, for any $y \in \mathbb{R}^+$ and $\varphi \in \mathbb{J}$,

$$|\mathcal{G}(\varphi, y(\varphi))| \leq \varphi^{\frac{1}{6}} \left(1 + \frac{\varphi^{2/3}}{16} |\varphi^{1/3} y(\varphi)| \right)$$

$$\leq \left(1 + \frac{1}{16} \|y\|_{\mathbb{C}_{\frac{1}{3}}}\right). \quad (4.2)$$

Hence, the assumption (A_1) is fulfilled having $\mathcal{M} = 1$ and $m = \frac{1}{16}$. It is easy to verify that the assumption (A_2) is hold too. In fact, by simple computations, we obtain

$$\mathcal{G} := \frac{m\mathcal{M}_1\Gamma(\delta)}{\vartheta^\lambda\Gamma(\lambda+1)} [(\varpi_2 - \varpi_1)^\lambda + (\varpi_2 - \varpi_1)^{\lambda+1-\delta}] \approx -0.03510 < 1. \quad (4.3)$$

Hence, all suppositions of Theorem 3.1 implies that the problem (1.1) has a unique solution in $\mathbb{C}_{\frac{2}{3}}^{\frac{2}{3}}([0, 2])$.

Also, assume that $\mathcal{G}(\varphi, y(\varphi)) = \varphi^{-\frac{1}{6}} + \frac{\varphi^{5/6}}{16} \sin y(\varphi)$. Thus $|\mathcal{G}(\varphi_1, y(\varphi))| \leq \varphi^{-\frac{1}{6}} + \frac{\varphi^{5/6}}{16} = \eta(\varphi) \in \mathbb{C}_{1-\delta}([0, 2])$. So, (A_3) is satisfied. Therefore, in view of Theorem 3.2, we conclude that problem (1.1) has a solution in $\mathbb{C}_{1/3}^{2/3}([0, 2])$.

Finally, if $\mathcal{G}(\varphi, y(\varphi)) = \varphi^{-\frac{1}{6}} + \frac{\varphi^{5/6}}{16} \sin y(\varphi)$, then for $y, \omega \in \mathbb{R}^+$ and $\varphi \in \mathbb{J}$, we have

$$|\mathcal{G}(\varphi, y(\varphi)) - \mathcal{G}(\varphi, \omega(\varphi))| \leq \frac{1}{16} |y - \omega|.$$

Therefore, the assumption (A_4) is fulfilled having $L = \frac{1}{16}$. Clearly, assumption (A_5) and inequality (3.33) are holds. In fact, simple computations yields

$$\begin{aligned} Q &:= \left[\left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)} + \frac{\mathfrak{B}(\delta - n, \lambda + 1)}{\vartheta^\delta\Gamma(\delta - n)} \right] \\ &\times \frac{\Gamma(\delta - n)(\varpi_2 - \varpi_1)^\lambda}{\vartheta^\lambda\mathfrak{B}(\delta - n, 1)\Gamma(\lambda + 1)} L \approx 0.1456 < 1, \end{aligned} \quad (4.4)$$

and

$$\left| \frac{m_2}{m_1 + m_2} \right| \sum_{\kappa=1}^n \frac{(\varpi_2 - \varpi_1)^{n-\kappa+\lambda}}{\vartheta^{n-\delta+\lambda}\Gamma(\delta - \kappa + 1)} \frac{\Gamma(\delta - n + 1)}{\vartheta^\lambda\Gamma(\lambda + 1)} L \approx 0.0495 < 1 \quad (4.5)$$

of Theorem 3.3, shows that problem (1.1) has a solution in $\mathbb{C}_{\frac{1}{3}}^{\frac{2}{3}}([0, 2])$.

5. Conclusions

In this approach, we have established certain existence consequences for the solution of *BVP* for Hilfer-*FDEs* depend on the lessening of *FDEs* to integral equations. The proposed scheme with the fixed point assertions unifies the existing results in the frame of *RL* and Caputo *GPF* sense, respectively. Besides that, the analysis's comprehensive improvements are dependent on various techniques such as Schauders, Schaefer's and Kransnoselskiis fixed point theorems. Also, the Hilfer *GPF*-derivative comprise two parameters and a proportionality index ϑ .

- If $\vartheta \mapsto 1$ and $\lambda = [0, 1]$, then the contemplated problem converted to *RL* and Caputo fractional derivative [8]. If $\vartheta \in (0, 1)$ and $\zeta = 0, 1$ we recaptures the *RL* and Caputo *GPF*-derivative [25], respectively (see Figure 1).

- Clearly, if $\vartheta, \zeta \in (0, 1)$, then the newly employed derivatives amalgamate the existing ones in the adjustment of Hilfer, *RL* and *GPF*-derivative, (see Figure 2).
- If $\vartheta \mapsto 1$ and $\zeta, \lambda \in [0, 1]$, then the formulation for this problem enjoys Hilfer fractional derivative [8], (see Figure 3).




Figure 1. Plot of $y(\varphi)$, for the *RL* fractional derivatives ($\zeta = 0, \vartheta = 1$), and *GPF*-derivatives ($\zeta = 0, \vartheta \in (0, 1)$).




Figure 2. Graph of $y(\varphi)$, for the *RL* fractional derivatives ($\zeta = 0, \vartheta = 1$), *GPF*-derivatives ($\zeta = 0, \vartheta = 0.8$) and Hilfer *GPF*-derivatives ($\zeta \in (0, 1), \vartheta \in (0, 1)$).




Figure 3. Plot of $y(\varphi)$, for the Hilfer fractional derivatives ($\vartheta = 1$) and Hilfer *GPF*-derivatives ($\vartheta \in (0, 1)$).

Moreover, a stimulative example is presented to show the efficacy of the established outcomes. We hope that the testified outcomes here will have a considerable impact for more parameters on the stability and other qualitative features of differential equations in the areas of interest of applied sciences.

Conflict of interest

The authors declare that there is no conflict of interests.

References

1. M. Nazeer, F. Hussain, M. Ijaz Khan, Asad-ur-Rehman, E. R. El-Zahar, Y. M. Chu, et al., Theoretical study of MHD electro-osmotically flow of third-grade fluid in micro channel, *Appl. Math. Comput.*, **420** (2022), 126868. <https://doi.org/10.1016/j.amc.2021.126868>
2. Y. M. Chu, B. M. Shankaralingappa, B. J. Giresha, F. Alzahrani, M. Ijaz Khan, S. U. Khan, Combined impact of Cattaneo-Christov double diffusion and radiative heat flux on bio-convective flow of Maxwell liquid configured by a stretched nano-material surface, *Appl. Math. Comput.*, **419** (2022), 126883. <https://doi.org/10.1016/j.amc.2021.126883>
3. Y. M. Chu, U. Nazir, M. Sohail, M. M. Selim, J. R. Lee, Enhancement in thermal energy and solute particles using hybrid nanoparticles by engaging activation energy and chemical reaction over a parabolic surface via finite element approach, *Fractal Fract.*, **5** (2021), 119. <https://doi.org/10.3390/fractalfract5030119>
4. T. H. Zhao, O. Castillo, H. Jahanshahi, A. Yusuf, M. O. Alassafi, F. E. Alsaadi, et al., A fuzzy-based strategy to suppress the novel coronavirus (2019-NCOV) massive outbreak, *Appl. Comput. Math.*, **20** (2021), 160–176.
5. Z. Denton, A. S. Vatsala, Fractional integral inequalities and applications, *Comput. Math. Appl.*, **59** (2010), 1087–1094. <https://doi.org/10.1016/j.camwa.2009.05.012>
6. R. Khalil, M. Al Horani, A. Yousef, M. Sababheh, A new definition of fractional derivative, *J. Comput. Appl. Math.*, **264** (2014), 65–70. <https://doi.org/10.1016/j.cam.2014.01.002>
7. R. Almeida, A Caputo fractional derivative of a function with respect to another function, *Commun. Nonlinear Sci.*, **44** (2017), 460–481. <https://doi.org/10.1016/j.cnsns.2016.09.006>
8. A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and applications of fractional differential equations*, Elsevier, 2006.
9. R. Hilfer, *Applications of fractional calculus in physics*, World Scientific, 2000.
10. A. Atangana, On the new fractional derivative and application to nonlinear Fisher's reaction-diffusion equation, *Appl. Math. Comput.*, **273** (2016), 948–956. <https://doi.org/10.1016/j.amc.2015.10.021>
11. S. Rashid, F. Jarad, M. A. Noor, H. Kalsoom, Inequalities by means of generalized proportional fractional integral operators with respect to another function, *Mathematics*, **7** (2020), 1225. <https://doi.org/10.3390/math7121225>

12. R. A. Yan, S. R. Sun, Z. L. Han, Existence of solutions of boundary value problems for Caputo fractional differential equations on time scales, *Bull. Iranian Math. soc.*, **42** (2016), 247–262.
13. A. Atangana, D. Baleanu, Application of fixed point theorem for stability analysis of a nonlinear Schrödinger with Caputo-Liouville derivative, *Filomat*, **31** (2017), 2243–2248. <https://doi.org/10.2298/FIL1708243A>
14. F. Jarad, S. Harikrishnan, K. Shah, K. Kanagarajan, Existence and stability results to a class of fractional random implicit differential equations involving a generalized Hilfer fractional derivative, *Discrete Cont. Dyn. S*, **13** (2020), 723–739. <https://doi.org/10.3934/dcdss.2020040>
15. O. A. Arqub, Numerical simulation of time-fractional partial differential equations arising in fluid flows via reproducing Kernel method, *Int. J. Numer. Method. H.*, **30** (2020), 4711–4733. <https://doi.org/10.1108/HFF-10-2017-0394>
16. S. Djennadi, N. Shawagfeh, M. Inc, M. S. Osman, J. F. Gómez-Aguilar, O. A. Arqub, The Tikhonov regularization method for the inverse source problem of time fractional heat equation in the view of ABC-fractional technique, *Phys. Scr.*, **96** (2021), 094006. <https://doi.org/10.1088/1402-4896/ac0867>
17. R. Khalil, M. A. Horani, A. Yousef, M. Sababheh, A new definition of fractional derivative, *J. Comput. Appl. Math.*, **264** (2014), 65–70. <https://doi.org/10.1016/j.cam.2014.01.002>
18. T. Abdeljawad, On conformable fractional calculus, *J. Comput. Appl. Math.*, **279** (2015), 57–66. <https://doi.org/10.1016/j.cam.2014.10.016>
19. D. R. Anderson, D. J. Ulness, Newly defined conformable derivatives, *Adv. Dyn. Syst. Appl.* **10** (2015), 109–137. <https://doi.org/10.13140/RG.2.1.1744.9444>
20. T. H. Zhao, W. M. Qian, Y. M. Chu, Sharp power mean bounds for the tangent and hyperbolic sine means, *J. Math. Inequal.*, **15** (2021), 1459–1472. <https://doi.org/10.7153/jmi-2021-15-100>
21. S. Rashid, E. I. Abouelmagd, S. Sultana, Y. M. Chu, New developments in weighted n -fold type inequalities via discrete generalized \hat{h} -proportional fractional operators, *Fractals*, **30** (2022), 2240056. <https://doi.org/10.1142/S0218348X22400564>
22. S. Rashid, E. I. Abouelmagd, A. Khalid, F. B. Farooq, Y. M. Chu, Some recent developments on dynamical \hat{h} -discrete fractional type inequalities in the frame of nonsingular and nonlocal kernels, *Fractals*, **30** (2022), 2240110. <https://doi.org/10.1142/S0218348X22401107>
23. M. Al Qurashi, S. Rashid, S. Sultana, H. Ahmad, K. A. Gepreel, New formulation for discrete dynamical type inequalities via h -discrete fractional operator pertaining to nonsingular kernel, *Math. Biosci. Eng.*, **18** (2021), 1794–1812. <https://doi.org/10.3934/mbe.2021093>
24. S. S. Zhou, S. Rashid, S. Parveen, A. O. Akdemir, Z. Hammouch, New computations for extended weighted functionals within the Hilfer generalized proportional fractional integral operators, *AIMS Math.*, **6** (2021), 4507–4525. <https://doi.org/10.3934/math.2021267>
25. F. Jarad, T. Abdeljawad, J. Alzabut, Generalized fractional derivatives generated by a class of local proportional derivatives. *Eur. Phys. J. Spec. Top.*, **226** (2017), 3457–3471. <https://doi.org/10.1140/epjst/e2018-00021-7>

26. S. Rashid, S. Sultana, Y. Karaca, A. Khalid, Y. M. Chu, Some further extensions considering discrete proportional fractional operators, *Fractals*, **30** (2022), 2240026. <https://doi.org/10.1142/S0218348X22400266>
27. K. Karthikeyan, P. Karthikeyan, H. M. Baskonus, K. Venkatachalam, Y. M. Chu, Almost sectorial operators on Ψ -Hilfer derivative fractional impulsive integro-differential equations, *Math. Method. Appl. Sci.*, **45** (2022), 8045–8059. <https://doi.org/10.1002/mma.7954>
28. S. Rashid, S. Sultana, Y. Karaca, A. Khalid, Y. M. Chu, Some further extensions considering discrete proportional fractional operators, *Fractals*, **30** (2022), 2240026. <https://doi.org/10.1142/S0218348X22400266>
29. S. Rashid, F. Jarad, M. A. Noor, Grüss-type integrals inequalities via generalized proportional fractional operators, *RACSAM*, **114** (2020), 93. <https://doi.org/10.1007/s13398-020-00823-5>
30. I. Ahmad, P. Kumam, F. Jarad, P. Borisut, Jirakitpuwapat, On Hilfer generalized proportional fractional derivative, *Adv. Differ. Equ.*, **2020** (2020), 329. <https://doi.org/10.1186/s13662-020-02792-w>
31. K. Shah, D. Vivek, K. Kanagarajan, Dynamics and stability of α -fractional pantograph equations with boundary conditions, *Bol. Soc. Paran. Mat.*, **39** (2021), 43–55. <https://doi.org/10.5269/bspm.41154>
32. D. Vivek, K. Kanagarajan, E. Elsayed, Some existence and stability results for Hilfer-fractional implicit differential equations with nonlocal conditions, *Mediterr. J. Math.*, **15** (2018), 15. <https://doi.org/10.1007/s00009-017-1061-0>
33. O. A. Arqub, Reproducing Kernel algorithm for the analytical-numerical solutions of nonlinear systems of singular periodic boundary value problems, *Math. Probl. Eng.*, **2015** (2015), 518406. <https://doi.org/10.1155/2015/518406>
34. S. Djennadi, N. Shawagfeh, O. A. Arqub, A fractional Tikhonov regularization method for an inverse backward and source problems in the time-space fractional diffusion equations, *Chaos Soliton. Fract.*, **150** (2021), 111127. <https://doi.org/10.1016/j.chaos.2021.111127>
35. W. Shammakh, H. Z. Alzum, Existence results for nonlinear fractional boundary value problem involving generalized proportional derivative, *Adv. Differ. Equ.*, **2019** (2019), 94. <https://doi.org/10.1186/s13662-019-2038-z>
36. A. Granas, J. Dugundi, *Fixed point theory*, New York: Springer, 2003.
37. M. Krasnoselskii, Two remarks about the method of successive approximations, *Uspekhi Mat. Nauk*, **10** (1955), 123–127.



AIMS Press

©2023 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)