



Research article

Solving an integral equation via orthogonal generalized α - ψ -Geraghty contractions

Senthil Kumar Prakasam¹, Arul Joseph Gnanaprakasam^{1,*}, Gunaseelan Mani² and Fahd Jarad^{3,4,*}

¹ Department of Mathematics, College of Engineering and Technology, Faculty of Engineering and Technology, SRM Institute of Science and Technology, SRM Nagar, Kattankulathur 603203, Kanchipuram, Chennai, Tamil Nadu, India

² Department of Mathematics, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences, Chennai 602105, Tamil Nadu, India

³ Department of Mathematics, Çankaya University, Etimesgut 06790, Ankara, Turkey

⁴ Department of Medical Research, China Medical University, Taichung 40402, Taiwan

* **Correspondence:** Email: aruljoseph.alex@gmail.com, fahd@cankaya.edu.tr.

Abstract: In this paper, we introduce orthogonal generalized \mathbf{O} - α - ψ -Geraghty contractive type mappings and prove some fixed point theorems in \mathbf{O} -complete \mathbf{O} -b-metric spaces. We also provide an illustrative example to support our theorem. The results proved here will be utilized to show the existence of a solution to an integral equation as an application.

Keywords: \mathbf{O} -b-metric space; fixed point; \mathbf{O} - α -admissible; orthogonal generalized \mathbf{O} - α - ψ -Geraghty contraction

Mathematics Subject Classification: 47H10, 54H25

1. Introduction and preliminaries

Banach [1] introduced one of the most essential Banach Contraction Principles. In 1993, Czerwik [2] initiated the notion of a b-metric space and proved the fixed point theorem (FPT) in this space. Aydi et al. [3] proved the common FPT for weak ϕ -contractions in b-metric space. The existence and uniqueness of a fixed point of ϕ -contractions was proved by Pacurar [4]. In 2018, Zada et al. [5] elaborated a FPT of a rational contraction. Geraghty [6] expanded the Banach contraction principle in 1973 by factoring an auxiliary function of complete metric space.

One of the interesting results was given by Samet et al. [7] by defining α - ψ -contractive maps via α -admissible mappings. After that, Cho et al. [8] introduced the α -GC (Geraghty contraction) type

maps in metric space and proved some FPT's of these functions. Popescu [9] developed α -GC maps and proved the fixed point theorem in complete metric space. From the above work, Karapınar [11] introduced generalized α - ψ -GC type maps, and see also [10, 12]. Furthermore, several authors proved the common fixed point theorem in many metric spaces, see [13–19].

An orthogonality notion in metric spaces was introduced by Gordji et al. [20]. The theory of an orthogonal set has general application in a number of mathematical areas, and there are many types of orthogonality, see [21–29].

In this paper, we prove a FPT in an \mathbf{O} -complete \mathbf{O} -b-metric space (\mathbf{O}_b -MS) with \mathbf{O} -generalized α - ψ -GC type maps. Moreover, an example and application to an integral equation are given to strengthen our main results.

2. Preliminaries

Throughout this paper, the standard letter \mathbb{R} is the set of all real numbers, \mathbb{R}^+ represents the set of all positive real numbers, \mathbb{N} denotes the set of all natural numbers, \mathbb{N}_0 denotes the set of all positive natural numbers, and \mathbb{Z} denotes the set of all integers. First, we recall some standard definitions and other results that will be needed in the sequel.

In 1993, Czerwik [2] introduced a b-metric space as follows:

Definition 2.1. [2] Let Q be a non-void set. Let $\tau: Q \times Q \rightarrow \mathbb{R}^+$ be called b-metric on Q if for all $\theta, \mu, \nu \in Q$ the below conditions hold:

- (i) $0 \leq \tau(\theta, \mu)$ and $\tau(\theta, \mu) = 0$ iff $\theta = \mu$;
- (ii) $\tau(\theta, \mu) = \tau(\mu, \theta)$;
- (iii) $\tau(\theta, \mu) \leq g[\tau(\theta, \nu) + \tau(\nu, \mu)]$.

Then, (Q, τ) is said to be a b-metric space (with constant $g \geq 1$). In 2014, Karapınar [11] introduced the concept of α -regularity as follows:

Definition 2.2. [11] Let (Q, τ) be a b-metric space and $\alpha: Q \times Q \rightarrow Q$ be a map. Q is called α -regular if for each sequence $\{\theta_\eta\} \in Q$ such that $\alpha(\theta_\eta, \theta_{\eta+1}) \geq 1$, $\forall \eta \in \mathbb{N}$ and $\theta_\eta \rightarrow \theta \in Q$ as $\eta \rightarrow \infty$, \exists a sub-sequence $\{\theta_{\eta(\gamma)}\}$ of $\{\theta_\eta\}$ with $\alpha(\theta_{\eta(\gamma)}, \theta) \geq 1$, for all $\gamma \in \mathbb{N}$.

The notion of an orthogonal set was presented by Gordji et al. [21].

Definition 2.3. [21] Let Q be a non empty set and $\perp \subseteq Q \times Q$ be a binary relation. If \perp holds with the constraint

$$\exists \theta_0 \in Q : (\forall \theta \in Q, \theta \perp \theta_0) \quad \text{or} \quad (\forall \theta \in Q, \theta_0 \perp \theta),$$

then (Q, \perp) is said to be an orthogonal set.

Gordji et al. [21] presented the definition of an orthogonal sequence in 2017 as follows.

Definition 2.4. [21] Let (Q, \perp) be an orthogonal set. A sequence $\{\theta_\eta\}_{\eta \in \mathbb{N}}$ is called an orthogonal sequence (\mathbf{O} -sequence) if

$$(\forall \eta, \theta_\eta \perp \theta_{\eta+1}) \quad \text{or} \quad (\forall \eta, \theta_{\eta+1} \perp \theta_\eta).$$

Definition 2.5. A tripled (Q, \perp, τ) is called an \mathbf{O}_b -MS if (Q, \perp) is an orthogonal set and (Q, τ) is a b-metric space.

Definition 2.6. Let (Q, \perp, τ) be an \mathbf{O}_b -MS.

(1) $\{\mu_\eta\}$, an orthogonal sequence in Q , converges at a point μ if

$$\lim_{\eta \rightarrow \infty} (E(\mu_\eta, \mu)) = 0.$$

(2) $\{\mu_\eta\}, \{\mu_m\}$ are orthogonal sequences in Q and are said to be orthogonal-Cauchy sequences if

$$\lim_{\eta, m \rightarrow \infty} (E(\mu_\eta, \mu_m)) < \infty.$$

Definition 2.7. [20] Let (Q, \perp, τ) be an \mathbf{O}_b -MS. Then, $E: Q \rightarrow Q$ is said to be orthogonally continuous in $\mu \in Q$ if, for each \mathbf{O} -sequence $\{\mu_\eta\}_{\eta \in \mathbb{N}}$ in Q with $\mu_\eta \rightarrow \mu$, we have $E(\mu_\eta) \rightarrow E(\mu)$. Also, E is said to be orthogonal continuous on Q if E is orthogonal continuous in each $\mu \in Q$.

Example 2.8. [20] Let $Q = \mathbb{R}$ and suppose $\mu \perp \theta$ if

$$\theta, \mu \in \left(\eta + \frac{1}{3}, \eta + \frac{2}{3} \right)$$

for some $\eta \in \mathbb{Z}$ or $\theta = 0$.

It is clear that (Q, \perp) is an orthogonal set. A map $E: Q \rightarrow Q$ is defined as $E(\theta) = [\theta]$. Then, E is orthogonal-continuous on Q , because if $\{\theta_r\}$ is an arbitrary \mathbf{O} -sequence in Q such that $\{\theta_r\}$ converges to $\theta \in Q$, then the below cases hold:

Case 1: If $\theta_r = 0 \forall r$, then $\theta = 0$ and $E(\theta_r) = 0 = E(\theta)$.

Case 2: If $\theta_{r_0} \neq 0$ for some r_0 , then there exists $m \in \mathbb{Z}$ such that $\theta_r \in (m + \frac{1}{3}, m + \frac{2}{3})$, for all $r \geq r_0$. Thus, $\theta \in [m + \frac{1}{3}, m + \frac{2}{3}]$, and $E(\theta_r) = m = E(\theta)$.

This means that E is orthogonal-continuous on Q , but it is not continuous on Q .

The concept of orthogonal completeness in metric spaces is defined by Gordji et al. [21] as follows.

Definition 2.9. [21] Let (Q, \perp, τ) be an orthogonal metric space. Then, Q is said to be \mathbf{O} -complete if every orthogonal Cauchy sequence is convergent.

Definition 2.10. [21] Let (Q, \perp) be an orthogonal set. A function $E: Q \rightarrow Q$ is called orthogonal-preserving if $E\theta \perp E\mu$ whenever $\theta \perp \mu$.

Ramezani [26] introduced the notion of being orthogonal α -admissible as follows.

Definition 2.11. [26] Let $E: Q \rightarrow Q$ be a map and let $\alpha: Q \times Q \rightarrow \mathbb{R}^+$ be a function. Then, E is said to be \mathbf{O} - α -admissible if $\forall \theta, \mu \in Q$ with $\theta \perp \mu$

$$\alpha(\theta, \mu) \geq 1 \implies \alpha(E\theta, E\mu) \geq 1.$$

In 2021, Gnanaprakasam et al. [25] introduced the notion of being orthogonal triangular α -admissible, defined as below:

Definition 2.12. [25] A self-map $E : Q \rightarrow Q$ is called **O**-triangular α -admissible if the following holds:

- (E1) E is **O**- α -admissible,
 (E2) $\alpha(\theta, \nu) \geq 1$, $\alpha(\nu, \mu) \geq 1$ with $\theta \perp \nu$ and $\nu \perp \mu \implies \alpha(\theta, \mu) \geq 1, \forall \theta, \mu, \nu \in Q$.

Recently, Popescu [9] has developed the notion of a triangular α -orbital admissible mappings, and we extend it to orthogonal in **O**_b-MS. The following definitions will be needed in the main result.

Definition 2.13. Let $E : Q \rightarrow Q$ be a map, and let there be a function $\alpha : Q \times Q \rightarrow \mathbb{R}^+$. Then, E is called **O**- α -orbital admissible if the below constraint holds:

- (E3) $\alpha(\theta, E\theta) \geq 1 \implies \alpha(E\theta, E^2\theta) \geq 1, \forall \theta \in Q$.

Example 2.14. Let $Q = \{0, 1, 2, 3\}$, $\tau : Q \times Q \rightarrow \mathbb{R}, \tau(\theta, \mu) = |\theta - \mu|^2, E : Q \rightarrow Q$ such that $E0 = 0, E1 = 2, E2 = 1, E3 = 3$. $\alpha : Q \times Q \rightarrow \mathbb{R}, \alpha(\theta, \mu) = 1$ if $(\theta, \mu) \in \{(0, 1), (0, 2), (1, 1), (2, 2), (1, 2), (2, 1), (1, 3), (2, 3)\}$ with $\theta \perp \mu$, and $\alpha(\theta, \mu) = 0$ otherwise.

Since $\alpha(1, E1) = \alpha(1, 2) = 1$ and $\alpha(2, E2) = \alpha(2, 1) = 1$, we have that E is orthogonal α -orbital admissible.

Definition 2.15. Let $E : Q \rightarrow Q$ and $\alpha : Q \times Q \rightarrow \mathbb{R}^+$ be maps. Then, E is called **O**-triangular- α -orbital admissible if E is **O**- α -orbital admissible and the following property holds:

- (E4) $\forall \theta, \mu \in Q$ with $\theta \perp \mu$, $\alpha(\theta, \mu) \geq 1$ and $\alpha(\mu, E\mu) \geq 1 \implies \alpha(\theta, E\mu) \geq 1$.

Example 2.16. Let $Q = \mathbb{R}, E\mu = \mu^3 + \sqrt{\mu}$, and $\alpha(\mu, \theta) = \mu^5 - \theta^5 + 1$, for all $\mu, \theta \in Q$ with $\mu \perp \theta$. Then, E is an **O**-triangular α -admissible mapping.

Lemma 2.17. Let $E : Q \rightarrow Q$ be an **O**-triangular- α -orbital admissible map. Consider that $\exists \theta_0 \in Q$ such that $\theta_0 \perp E\theta_0$ and $\alpha(\theta_0, E\theta_0) \geq 1$. An **O**-sequence $\{\theta_\eta\}$ is defined by $\theta_{\eta+1} = E\theta_\eta, \forall \eta \in \mathbb{N}$ with $\theta_\eta \perp E\theta_\eta$ or $E\theta_\eta \perp \theta_\eta$. Then, we get $\alpha(\theta_\eta, \theta_\nu) \geq 1, \forall \eta, \nu \in \mathbb{N}$ with $\eta < \nu$.

In this section, inspired by the concept of generalized α - ψ -GC type maps defined by Afshari et al. [28], we introduce a new orthogonal generalized α - ψ -GC type mapping and prove some FPT's for these contraction mappings in an **O**-complete **O**_b-MS.

3. Main results

Let Λ be a set of all increasing and continuous functions, and $\psi \in \Lambda$ is defined as $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, with $\psi^{-1}(\{0\}) = \{0\}$.

Let Γ be the family of all non-decreasing functions $\lambda : [0, \infty) \rightarrow [0, \frac{1}{g})$ which satisfy the condition

$$\lim_{\eta \rightarrow \infty} \lambda(\zeta_\eta) = \frac{1}{g} \implies \lim_{\eta \rightarrow \infty} \zeta_\eta = 0 \quad \text{for some } g \geq 1.$$

First, we explain the definition of an **O**-generalized α - ψ -GC type(A) map in an **O**-complete **O**_b-MS.

Definition 3.1. Let (Q, \perp, τ) be an **O**-complete **O**_b-MS, and let there be a map $E : Q \rightarrow Q$. E is called an **O**-generalized α - ψ -GC type(A) map whenever $\exists \alpha : Q \times Q \rightarrow \mathbb{R}^+$, and, for $\mathfrak{L} \geq 0$ such that

$$\mathbf{M}(\theta, \mu) = \max \left\{ \tau(\theta, \mu), \tau(\theta, E\theta), \tau(\mu, E\mu), \frac{\tau(\theta, E\mu) + \tau(\mu, E\theta)}{2g} \right\},$$

$$\mathcal{N}(\theta, \mu) = \min\{\tau(\theta, E\theta), \tau(\mu, E\theta)\},$$

we have

$$\alpha(\theta, \mu)\psi(g^3\tau(E\theta, E\mu)) \leq \lambda(\psi(\mathbf{M}(\theta, \mu)))\psi(\mathbf{M}(\theta, \mu)) + \mathfrak{L}\vartheta(\mathcal{N}(\theta, \mu)), \quad (3.1)$$

for all $\theta, \mu \in \mathcal{Q}$ with $\theta \perp \mu$, where $\lambda \in \Gamma$ and $\psi, \vartheta \in \Lambda$.

$$\lambda(\psi(\mathbf{M}(\theta, \mu))) < \frac{1}{g} \text{ for all } \theta, \mu \in \mathcal{Q}, \text{ with } \theta \perp \mu \text{ and } \theta \neq \mu.$$

Now, we generalize and improve our FPT from Afshari et al. [28] by introducing the notion of an \mathbf{O} -generalized α - ψ -GC type(A) map in \mathbf{O} -complete \mathbf{O}_b -MS.

Theorem 3.2. *Let $(\mathcal{Q}, \perp, \tau)$ be an \mathbf{O} -complete \mathbf{O}_b -MS, and $E : \mathcal{Q} \rightarrow \mathcal{Q}$ satisfies the below properties:*

- (i) *E is orthogonal-preserving,*
- (ii) *E is an \mathbf{O} -generalized α - ψ -GC type(A) map,*
- (iii) *E is \mathbf{O} -triangular α -orbital admissible,*
- (iv) *$\exists \theta_0 \in \mathcal{Q}$ such that $\theta_0 \perp E\theta_0$ and $\alpha(\theta_0, E\theta_0) \geq 1$,*
- (v) *E is \mathbf{O} -continuous.*

Then, E has a UFP (unique fixed point).

Proof. By the condition (iv), there exists $\theta_0 \in \mathcal{Q}$ such that

$$\theta_0 \perp E(\theta_0) \text{ or } E(\theta_0) \perp \theta_0 \text{ and } \alpha(\theta_0 \perp E\theta_0) \geq 1.$$

Let

$$\theta_1 = E(\theta_0), \theta_2 = E(\theta_1) = E^2(\theta_0), \dots, \theta_\eta = E(\theta_{\eta-1}) = E^\eta(\theta_0), \theta_{\eta+1} = E(\theta_\eta) = E^{\eta+1}(\theta_0), \quad \forall \eta \in \mathbb{N} \cup \{0\}.$$

If $\theta_{\eta^*} = \theta_{\eta^*+1}$ for $\eta^* \in \mathbb{N} \cup \{0\}$, then θ_{η^*} is a fixed point of E . Therefore, the proof is complete. So, we consider $\theta_\eta \neq \theta_{\eta+1}$. Thus, we have $\tau(E\theta_\eta, E\theta_{\eta+1}) > 0$. Since E is \mathbf{O} -preserving, we get

$$\theta_\eta \perp \theta_{\eta+1} \text{ or } \theta_{\eta+1} \perp \theta_\eta, \quad \forall \eta \in \mathbb{N} \cup \{0\}.$$

We construct an \mathbf{O} -sequence $\{\theta_\eta\}$. Since E is an \mathbf{O} -generalized α - ψ -GC type(A) map, we get

$$\theta_{\eta+1} = E\theta_\eta, \quad \forall \eta \in \mathbb{N}_0.$$

Since the map E is an orthogonal triangular α -orbital admissible, by Lemma 2.17, we get

$$\alpha(\theta_\eta, \theta_{\eta+1}) \geq 1, \quad \forall \eta \in \mathbb{N}_0. \quad (3.2)$$

By letting $\theta = \theta_{\eta-1}$ and $\mu = \theta_\eta$ in the inequality (3.1), using (3.2) and that ψ is an ascending map, we get

$$\begin{aligned}
\psi(\tau(\theta_\eta, \theta_{\eta+1})) &= \psi(\tau(E\theta_{\eta-1}, E\theta_\eta)) \\
&\leq \alpha(\theta_{\eta-1}, \theta_\eta)\psi(g^3(\tau(E\theta_{\eta-1}, E\theta_\eta))) \\
&\leq \lambda(\psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta)))\psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta)) + \mathfrak{L}\mathfrak{D}(\mathcal{N}(\theta_{\eta-1}, \theta_\eta)),
\end{aligned} \tag{3.3}$$

for all $\eta \in \mathbb{N}$, where

$$\begin{aligned}
\mathbf{M}(\theta_{\eta-1}, \theta_\eta) &= \max\left\{\tau(\theta_{\eta-1}, \theta_\eta), \tau(\theta_{\eta-1}, E\theta_{\eta-1}), \tau(\theta_\eta, E\theta_\eta), \frac{\tau(\theta_{\eta-1}, E\theta_\eta) + \tau(\theta_\eta, E\theta_{\eta-1})}{2g}\right\} \\
&= \max\left\{\tau(\theta_{\eta-1}, \theta_\eta), \tau(\theta_{\eta-1}, \theta_\eta), \tau(\theta_\eta, \theta_{\eta+1}), \frac{\tau(\theta_{\eta-1}, \theta_{\eta+1}) + \tau(\theta_\eta, \theta_\eta)}{2g}\right\} \\
&= \max\left\{\tau(\theta_{\eta-1}, \theta_\eta), \tau(\theta_\eta, \theta_{\eta+1}), \frac{\tau(\theta_{\eta-1}, \theta_{\eta+1})}{2g}\right\}
\end{aligned}$$

and

$$\mathcal{N}(\theta_{\eta-1}, \theta_\eta) = \min\{\tau(\theta_{\eta-1}, E\theta_{\eta-1}), \tau(\theta_\eta, E\theta_{\eta-1})\} = \min\{\tau(\theta_{\eta-1}, \theta_\eta), \tau(\theta_\eta, \theta_\eta)\} = 0. \tag{3.4}$$

Since

$$\frac{\tau(\theta_{\eta-1}, \theta_{\eta+1})}{2g} \leq \frac{g[\tau(\theta_{\eta-1}, \theta_\eta) + \tau(\theta_\eta, \theta_{\eta+1})]}{2g} \leq \max\{\tau(\theta_{\eta-1}, \theta_\eta) + \tau(\theta_\eta, \theta_{\eta+1})\},$$

we get

$$\mathbf{M}(\theta_{\eta-1}, \theta_\eta) \leq \max\{\tau(\theta_{\eta-1}, \theta_\eta), \tau(\theta_\eta, \theta_{\eta+1})\}. \tag{3.5}$$

Taking (3.5) and (3.4) into account, (3.3) yields

$$\begin{aligned}
\psi(\tau(\theta_\eta, \theta_{\eta+1})) &\leq \psi(g^3\tau(\theta_\eta, \theta_{\eta+1})) \\
&\leq \alpha(\theta_{\eta-1}, \theta_\eta)\psi(g^3\tau(\theta_\eta, \theta_{\eta+1})) \\
&\leq \lambda(\psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta)))\psi(\max\{\tau(\theta_{\eta-1}, \theta_\eta), \tau(\theta_\eta, \theta_{\eta+1})\}).
\end{aligned} \tag{3.6}$$

If $\eta \in \mathbb{N}$, we get $\max\{\tau(\theta_{\eta-1}, \theta_\eta), \tau(\theta_\eta, \theta_{\eta+1})\} = \tau(\theta_\eta, \theta_{\eta+1})$, and then by (3.6), we get

$$\psi(\tau(\theta_\eta, \theta_{\eta+1})) \leq \lambda(\psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta)))\psi(\tau(\theta_\eta, \theta_{\eta+1})) < \frac{1}{g}\psi(\tau(\theta_\eta, \theta_{\eta+1})) < \psi(\tau(\theta_\eta, \theta_{\eta+1})),$$

which is a contradiction. Thus, from (3.6) we conclude that

$$\begin{aligned}
\psi(\tau(\theta_\eta, \theta_{\eta+1})) &\leq \lambda(\psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta)))\psi(\tau(\theta_{\eta-1}, \theta_\eta)) \\
&< \frac{1}{g}\psi(\tau(\theta_{\eta-1}, \theta_\eta)) < \psi(\tau(\theta_{\eta-1}, \theta_\eta)), \quad \forall \eta \in \mathbb{N}.
\end{aligned} \tag{3.7}$$

Hence, $\{\psi(\tau(\theta_\eta, \theta_{\eta+1}))\}$ is a positive non-increasing sequence. Since ψ is ascending, the sequence $\{\tau(\theta_\eta, \theta_{\eta+1})\}$ is decreasing. Consequently, $\exists \epsilon \geq 0$ such that $\lim_{\eta \rightarrow \infty} \tau(\theta_\eta, \theta_{\eta+1}) = \epsilon$. We claim that $\epsilon = 0$.

Suppose, on the contrary, that

$$\lim_{\eta \rightarrow \infty} \tau(\theta_\eta, \theta_{\eta+1}) = \epsilon > 0.$$

Since $g \geq 1$, (3.7) can be approximated as

$$\frac{1}{g}\psi(\tau(\theta_\eta, \theta_{\eta+1})) \leq \psi(\tau(\theta_\eta, \theta_{\eta+1})) \leq \lambda(\psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta)))\psi(\tau(\theta_{\eta-1}, \theta_\eta)). \quad (3.8)$$

With regard to (3.2), in (3.8) we get

$$\frac{1}{g} \frac{\psi(\tau(\theta_\eta, \theta_{\eta+1}))}{\psi(\tau(\theta_{\eta-1}, \theta_\eta))} \leq \lambda(\psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta))) < \frac{1}{g}.$$

This yields $\lim_{\eta \rightarrow \infty} \lambda(\psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta))) = \frac{1}{g}$. Since $\lambda \in \Gamma$, we get $\lim_{\eta \rightarrow \infty} \psi(\mathbf{M}(\theta_{\eta-1}, \theta_\eta)) = 0$. We simplify that:

$$\lim_{\eta \rightarrow \infty} \psi(\tau(\theta_\eta, \theta_{\eta+1})) = 0.$$

Thus, by the fact that $\tau(\theta_\eta, \theta_{\eta+1}) \rightarrow \epsilon$ and the continuity of ψ , we get $\psi(\epsilon) = 0$. Since $\psi^{-1}(\{0\}) = \{0\}$, we have $\epsilon = 0$, and this is a contradiction. Thus, we get

$$\lim_{\eta \rightarrow \infty} \tau(\theta_\eta, \theta_{\eta+1}) = 0. \quad (3.9)$$

Now, we claim that $\lim_{\pi, \eta \rightarrow \infty} \tau(\theta_\eta, \theta_\pi) = 0$.

Consider, on the contrary, that $\exists \delta > 0$, and orthogonal subsequences $\{\theta_{\pi_i}\}$, and $\{\theta_{\eta_i}\}$ of $\{\theta_\eta\}$, with $\eta_i > \pi_i \geq j$, such that

$$\tau(\theta_{\pi_i}, \theta_{\eta_i}) \geq \delta. \quad (3.10)$$

Additionally, for every π_i , we choose the smallest integer η_i to fulfill (3.10), and $\eta_i > \pi_i \geq j$. Then, we get

$$\tau(\theta_{\pi_i}, \theta_{\eta_{i-1}}) < \delta. \quad (3.11)$$

From (3.10) and the condition (iii) in Definition 2.1, we get

$$\begin{aligned} \delta \leq \tau(\theta_{\eta_i}, \theta_{\pi_i}) &\leq g\tau(\theta_{\eta_i}, \theta_{\eta_{i+1}}) + g\tau(\theta_{\eta_{i+1}}, \theta_{\pi_i}) \\ &\leq g\tau(\theta_{\eta_i}, \theta_{\eta_{i+1}}) + g^2\tau(\theta_{\eta_{i+1}}, \theta_{\pi_{i+1}}) + g^2\tau(\theta_{\pi_{i+1}}, \theta_{\pi_i}). \end{aligned} \quad (3.12)$$

Taking $j \rightarrow \infty$ and (3.9) into account, Eq (3.12) yields

$$\begin{aligned} \delta &\leq g\tau(\theta_{\eta_i}, \theta_{\eta_{i+1}}) + g^2\tau(\theta_{\eta_{i+1}}, \theta_{\pi_{i+1}}) + g^2\tau(\theta_{\pi_{i+1}}, \theta_{\pi_i}) \\ &\leq \lim_{j \rightarrow \infty} (g\tau(\theta_{\eta_i}, \theta_{\eta_{i+1}}) + g^2\tau(\theta_{\eta_{i+1}}, \theta_{\pi_{i+1}}) + g^2\tau(\theta_{\pi_{i+1}}, \theta_{\pi_i})) \\ &\leq g^2 \lim_{j \rightarrow \infty} (\tau(\theta_{\eta_{i+1}}, \theta_{\pi_{i+1}})), \\ \frac{\delta}{g^2} &\leq \limsup_{j \rightarrow \infty} \tau(\theta_{\eta_{i+1}}, \theta_{\pi_{i+1}}), \end{aligned} \quad (3.13)$$

where $\lim_{j \rightarrow \infty} \tau(\theta_{\eta_j}, \theta_{\eta_{j+1}}) = \lim_{j \rightarrow \infty} \tau(\theta_{\pi_{j+1}}, \theta_{\pi_j}) = 0$. By Lemma 2.17, $\alpha(\theta_{\pi_j}, \theta_{\eta_j}) \geq 1$. Followed by (3.1), we get

$$\begin{aligned} \psi(\tau(\theta_{\eta_{j+1}}, \theta_{\pi_{j+1}})) &= \psi(\tau(E\theta_{\eta_j}, E\theta_{\pi_j})) \\ &\leq \psi(g^3 \tau(E\theta_{\eta_j}, E\theta_{\pi_j})) \\ &\leq \alpha(\theta_{\pi_j}, \theta_{\eta_j}) \psi(g^3 \tau(E\theta_{\eta_j}, E\theta_{\pi_j})) \\ &\leq \lambda(\psi(\mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j}))) \psi(\mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j})) + \mathfrak{L}\vartheta(\tau(\theta_{\pi_j}, E\theta_{\eta_j})), \end{aligned} \quad (3.14)$$

where

$$\begin{aligned} \mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j}) &= \max \left\{ \tau(\theta_{\eta_j}, \theta_{\pi_j}), \tau(\theta_{\eta_j}, E\theta_{\eta_j}), \tau(\theta_{\pi_j}, E\theta_{\pi_j}), \frac{\tau(\theta_{\eta_j}, E\theta_{\pi_j}) + \tau(\theta_{\pi_j}, E\theta_{\eta_j})}{2g} \right\} \\ &= \max \left\{ \tau(\theta_{\eta_j}, \theta_{\pi_j}), \tau(\theta_{\eta_j}, \theta_{\eta_{j+1}}), \tau(\theta_{\pi_j}, \theta_{\pi_{j+1}}), \frac{\tau(\theta_{\eta_j}, \theta_{\pi_{j+1}}) + \tau(\theta_{\pi_j}, \theta_{\eta_{j+1}})}{2g} \right\} \end{aligned}$$

and

$$\begin{aligned} \mathcal{N}(\theta_{\eta_j}, \theta_{\pi_j}) &= \min\{\tau(\theta_{\eta_j}, E\theta_{\eta_j}), \tau(\theta_{\pi_j}, E\theta_{\pi_j})\} \\ &= \min\{\tau(\theta_{\eta_j}, \theta_{\eta_{j+1}}), \tau(\theta_{\pi_j}, \theta_{\pi_{j+1}})\}. \end{aligned}$$

Notice that

$$\frac{\tau(\theta_{\eta_j}, \theta_{\pi_{j+1}}) + \tau(\theta_{\pi_j}, \theta_{\eta_{j+1}})}{2g} \leq \frac{g[\tau(\theta_{\eta_j}, \theta_{\pi_j}) + \tau(\theta_{\pi_j}, \theta_{\pi_{j+1}})] + g[\tau(\theta_{\pi_j}, \theta_{\eta_j}) + \tau(\theta_{\eta_j}, \theta_{\eta_{j+1}})]}{2g} \quad (3.15)$$

and

$$\tau(\theta_{\eta_j}, \theta_{\pi_j}) \leq g[\tau(\theta_{\eta_j}, \theta_{\eta_{j-1}}) + \tau(\theta_{\eta_{j-1}}, \theta_{\pi_j})] < g\tau(\theta_{\eta_j}, \theta_{\eta_{j-1}}) + g\delta. \quad (3.16)$$

Taking (3.11), (3.15) and (3.16) into account, we find that

$$\limsup_{j \rightarrow \infty} \mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j}) \leq g\delta, \quad (3.17)$$

$$\limsup_{j \rightarrow \infty} \mathcal{N}(\theta_{\eta_j}, \theta_{\pi_j}) = 0. \quad (3.18)$$

By allowing the upper limit to be $j \rightarrow \infty$ and using constraint (E4), (3.13), (3.17) and (3.18), inequality (3.14) becomes

$$\begin{aligned} \frac{1}{g} \psi(g\delta) &\leq \psi(g\delta) \leq \limsup_{j \rightarrow \infty} \psi(g^3 \tau(\theta_{\eta_{j+1}}, \theta_{\pi_{j+1}})) \\ &\leq \limsup_{j \rightarrow \infty} \alpha(\theta_{\pi_j}, \theta_{\eta_j}) \psi(g^3 \tau(\theta_{\eta_{j+1}}, \theta_{\pi_{j+1}})) \\ &\leq \limsup_{j \rightarrow \infty} \alpha(\theta_{\pi_j}, \theta_{\eta_j}) \psi(g^3 \tau(E\theta_{\eta_j}, E\theta_{\pi_j})) \\ &\leq \limsup_{j \rightarrow \infty} [\lambda(\psi(\mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j}))) \psi(\mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j})) + \mathfrak{L}\vartheta(\mathcal{N}(\tau(\theta_{\eta_j}, \theta_{\pi_j})))] \end{aligned}$$

$$\begin{aligned} &\leq \psi(g\delta) \limsup_{j \rightarrow \infty} \lambda(\psi(\mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j}))) \\ &\leq \frac{1}{g} \psi(g\delta). \end{aligned}$$

Then,

$$\limsup_{j \rightarrow \infty} \lambda(\psi(\mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j}))) = \frac{1}{g}.$$

Due to the fact that $\lambda \in \Gamma$, we get

$$\limsup_{j \rightarrow \infty} \lambda(\psi(\mathbf{M}(\theta_{\eta_j}, \theta_{\pi_j}))) = 0.$$

Thus, we assume that

$$\lim_{j \rightarrow \infty} \lambda(\psi(\tau(\theta_{\eta_j}, \theta_{\pi_j}))) = 0.$$

Consequently, due to the continuity of ψ and $\psi^{-1}(\{0\}) = \{0\}$, we obtain

$$\lim_{j \rightarrow \infty} \tau(\theta_{\eta_j}, \theta_{\pi_j}) = 0,$$

which contradicts (3.10). Therefore, $\{\theta_\eta\}$ is a \mathbf{O} -Cauchy sequence in \mathcal{Q} . Since \mathcal{Q} is an \mathbf{O} -complete \mathbf{O}_b -MS, $\exists \theta^* \in \mathcal{Q}$ such that $\lim_{\eta \rightarrow \infty} \theta_\eta = \theta^*$. The map E is \mathbf{O} -continuous, and it is obvious that $E\theta^* = \theta^*$. Hence, θ^* is a fixed point of E .

Now, we prove θ^* is a UFP of E . Suppose μ^* is another fixed point of E . If $\theta_\eta \rightarrow \mu^*$ as $\eta \rightarrow \infty$, we get $\theta^* = \mu^*$.

If $\lim_{\eta \rightarrow \infty} \{\theta_\eta\} \rightarrow \mu^*$, there is an orthogonal sub-sequence $\{\theta_{\eta_\gamma}\}$ such that $E\theta_{\eta_\gamma} \neq \mu^*$, $\forall \gamma \in \mathbb{N}$. By the choice of θ_0 , we get

$$\theta_0 \perp \mu^* \text{ (or) } \mu^* \perp \theta_0.$$

Since E is \mathbf{O} -preserving and $E^\eta \mu^* = \mu^*$, for all $\eta \in \mathbb{N}$, we get

$$E^\eta \theta_0 \perp \mu^* \text{ (or) } \mu^* \perp E^\eta \theta_0, \quad \forall \eta \in \mathbb{N}.$$

Since E is an \mathbf{O} -generalized α - ψ -GC type(A) map, we get

$$\psi(\tau(E^{\eta_\gamma} \theta_0, \mu^*)) = \psi(\tau(E^{\eta_\gamma} \theta_0, E^{\eta_\gamma} \mu^*)) \leq \psi(\tau(\theta_0, \mu^*)), \quad \gamma \in \mathbb{N}.$$

This implies $\psi(\tau(E^{\eta_\gamma} \theta_0, \mu^*)) \rightarrow -\infty$ as $\gamma \rightarrow \infty$. This yields that $\theta_\eta \rightarrow \mu^*$ as $\eta \rightarrow \infty$, which is a contradiction. Hence, E has a UFP. \square

We replace the continuity of map E in the above theorem by a suitable condition on \mathcal{Q} .

Theorem 3.3. *Let $(\mathcal{Q}, \perp, \tau)$ be an \mathbf{O} -complete \mathbf{O} -b-metric space, and $E : \mathcal{Q} \rightarrow \mathcal{Q}$ fulfills the following properties:*

- (i) E is orthogonal preserving,
(ii) E is an \mathbf{O} -generalized α - ψ -GC type(A) map,
(iii) E is \mathbf{O} -triangular- α orbital admissible,
(iv) $\exists \theta_0 \in \mathcal{Q}$ such that $\theta_0 \perp E\theta_0$ and $\alpha(\theta_0, E\theta_0) \geq 1$,
(v) E is \mathbf{O} - α -regular.

Then, E has a UFP.

Proof. From the proof of Theorem 3.2, we conclude that $\lim_{\eta \rightarrow \infty} \theta_\eta = \theta^*$. If \mathcal{Q} is \mathbf{O} - α -regular, then $\alpha(\theta_\eta, \theta_{\eta+1}) \geq 1$, \exists a subsequence $\{\theta_{\eta_\gamma}\}$ of $\{\theta_\eta\}$ such that

$$\alpha(\theta_{\eta_\gamma}, \theta^*) \geq 1, \quad \forall \gamma \in \mathbb{N}. \quad (3.19)$$

By the (iii) in Definition 2.1, we get

$$\tau(\theta^*, E\theta^*) \leq g\tau(\theta^*, \theta_{\eta_{\gamma+1}}) + g\tau(\theta_{\eta_{\gamma+1}}, E\theta^*) = g\tau(\theta^*, \theta_{\eta_\gamma}) + g\tau(E\theta_{\eta_\gamma}, E\theta^*).$$

Letting $\gamma \rightarrow \infty$, yields

$$\tau(\theta^*, E\theta^*) \leq \liminf_{\gamma \rightarrow \infty} g\tau(E\theta_{\eta_\gamma}, E\theta^*). \quad (3.20)$$

Using that $\psi \in \Lambda$, (3.19) and (3.20), we have

$$\begin{aligned} \psi(g^2\tau(\theta^*, E\theta^*)) &\leq \lim_{\gamma \rightarrow \infty} \psi(g^3\tau(E\theta_{\eta_\gamma}, E\theta^*)) \\ &\leq \lim_{\gamma \rightarrow \infty} \alpha(\theta_{\eta_{\gamma+1}}, \theta^*)\psi(g^3\tau(E\theta_{\eta_\gamma}, E\theta^*)) \\ &\leq \lim_{\gamma \rightarrow \infty} [\lambda(\psi(\mathbf{M}(\theta_{\eta_\gamma}, \theta^*)))\psi(\mathbf{M}(\theta_{\eta_\gamma}, \theta^*)) + \varrho\vartheta(\mathcal{N}(\theta_{\eta_\gamma}, \theta^*))]. \end{aligned} \quad (3.21)$$

We have

$$\begin{aligned} \mathbf{M}(\theta_{\eta_\gamma}, \theta^*) &= \max \left\{ \tau(\theta_{\eta_\gamma}, \theta^*), \tau(\theta_{\eta_\gamma}, E\theta_{\eta_\gamma}), \tau(\theta^*, E\theta^*), \frac{\tau(\theta_{\eta_\gamma}, E\theta^*) + \tau(\theta^*, E\theta_{\eta_\gamma})}{2g} \right\} \\ &= \max \left\{ \tau(\theta_{\eta_\gamma}, \theta^*), \tau(\theta_{\eta_\gamma}, \theta_{\eta_{\gamma+1}}), \tau(\theta^*, E\theta^*), \frac{\tau(\theta_{\eta_\gamma}, E\theta^*) + \tau(\theta^*, \theta_{\eta_{\gamma+1}})}{2g} \right\} \end{aligned}$$

and

$$\mathcal{N}(\theta_{\eta_\gamma}, \theta^*) = \min\{\tau(\theta_{\eta_\gamma}, E\theta_{\eta_\gamma}), \tau(\theta^*, E\theta_{\eta_\gamma})\} = \min\{\tau(\theta_{\eta_\gamma}, \theta_{\eta_{\gamma+1}}), \tau(\theta^*, \theta_{\eta_{\gamma+1}})\}.$$

Recall that

$$\frac{\tau(\theta_{\eta_\gamma}, E\theta^*) + \tau(\theta^*, \theta_{\eta_{\gamma+1}})}{2g} \leq \frac{g\tau(\theta_{\eta_\gamma}, \theta^*) + g\tau(\theta^*, E\theta^*) + \tau(\theta^*, \theta_{\eta_{\gamma+1}})}{2g}.$$

Then, by (3.9), we get

$$\limsup_{\gamma \rightarrow \infty} \frac{\tau(\theta_{\eta_\gamma}, E\theta^*) + \tau(\theta^*, \theta_{\eta_{\gamma+1}})}{2g} \leq \frac{\tau(\theta^*, E\theta^*)}{2}.$$

When $\gamma \rightarrow \infty$, we deduce

$$\begin{aligned} \lim_{\gamma \rightarrow \infty} \mathbf{M}(\theta_{\eta_\gamma}, \theta^*) &= \tau(\theta^*, E\theta^*), \\ \lim_{\gamma \rightarrow \infty} \mathcal{N}(\theta_{\eta_\gamma}, \theta^*) &= 0. \end{aligned}$$

Since $\lambda(\psi(\mathbf{M}(\theta_{\eta_\gamma}, \theta^*))) \leq \frac{1}{g}$, for all $\gamma \in \mathbb{N}$, from (3.21), we obtain

$$\psi(g^2\tau(\theta^*, E\theta^*)) \leq \frac{1}{g}\psi(\tau(\theta^*, E\theta^*)) \leq \psi(\tau(\theta^*, E\theta^*)).$$

Hence, $\tau(\theta^*, E\theta^*) = 0$, that is $E\theta^* = \theta^*$. Therefore, E has a fixed point.

Now, we prove θ^* is a UFP of E . Suppose μ^* is another fixed point of E . If $\theta_\eta \rightarrow \mu^*$ as $\eta \rightarrow \infty$, we get $\theta^* = \mu^*$.

If $\lim_{\eta \rightarrow \infty} \{\theta_\eta\} \not\rightarrow \mu^*$, there is a subsequence $\{\theta_{\eta_\gamma}\}$ such that $E\theta_{\eta_\gamma} \neq \mu^*$, $\forall \gamma \in \mathbb{N}$. By the choice θ_0 , we get

$$\theta_0 \perp \mu^* \text{ or } \mu^* \perp \theta_0.$$

Since E is \mathbf{O} -preserving and $E^\eta \mu^* = \mu^*$, $\forall \eta \in \mathbb{N}$, we get

$$E^\eta \theta_0 \perp \mu^* \text{ or } \mu^* \perp E^\eta \theta_0, \quad \forall \eta \in \mathbb{N}.$$

Since E is an \mathbf{O} -generalized α - ψ -GC type(A) map, we get

$$\psi(\tau(E^{\eta_\gamma} \theta_0, \mu^*)) = \psi(\tau(E^{\eta_\gamma} \theta_0, E^{\eta_\gamma} \mu^*)) \leq \psi(\tau(\theta_0, \mu^*)), \quad \gamma \in \mathbb{N}.$$

This implies $\psi(\tau(E^{\eta_\gamma} \theta_0, \mu^*)) \rightarrow -\infty$ as $\gamma \rightarrow \infty$. This yields that $\theta_\eta \rightarrow \mu^*$ as $\eta \rightarrow \infty$, and this is contradiction. Hence, E has a UFP. \square

We initiate the definition of an \mathbf{O} -generalized α - ψ -GC type(B) map as follows:

Definition 3.4. Let $(\mathcal{Q}, \perp, \tau)$ be an \mathbf{O} -complete \mathbf{O}_b -MS and let $E : \mathcal{Q} \rightarrow \mathcal{Q}$ be a map. E is called \mathbf{O} -generalized α - ψ -GC map of type(B) whenever $\exists \alpha : \mathcal{Q} \times \mathcal{Q} \rightarrow \mathbb{R}^+$ such that for all $\theta, \mu \in \mathcal{Q}$ with $\theta \perp \mu$ or $\mu \perp \theta$,

$$\alpha(\theta, \mu)\psi(g^3\tau(E\theta, E\mu)) \leq \lambda(\psi(\mathbf{M}(\theta, \mu)))\psi(\mathbf{M}(\theta, \mu)), \quad (3.22)$$

where

$$\mathbf{M}(\theta, \mu) = \max \left\{ \tau(\theta, \mu), \tau(\theta, E\theta), \tau(\mu, E\mu), \frac{\tau(\theta, E\mu) + \tau(\mu, E\theta)}{2g} \right\},$$

$\lambda \in \Gamma$ and $\psi \in \Lambda$.

Now, we generalize and improve our fixed point theorem from Afshari et al. [28] by introducing the notion of an \mathbf{O} -generalized α - ψ -GC type(B) map in \mathbf{O} -complete \mathbf{O}_b -MS.

Theorem 3.5. Let (Q, \perp, τ) be an \mathbf{O} -complete \mathbf{O}_b -MS, and let $E : Q \rightarrow Q$ satisfy the below properties:

- (i) E is orthogonal preserving,
- (ii) E is an \mathbf{O} -generalized α - ψ -GC map of type(B),
- (iii) E is \mathbf{O} -triangular- α orbital admissible,
- (iv) $\exists \theta_0 \in Q$ such that $\theta_0 \perp E\theta_0$ and $\alpha(\theta_0, E\theta_0) \geq 1$,
- (v) E is \mathbf{O} -continuous or Q is \mathbf{O} - α -regular.

Then, E has a UFP.

Now, we provide the example for Theorem 3.3.

Example 3.6. Let Q be a set of Lebesgue measurable functions $[0,1]$ such that $\int_0^1 |\theta(\zeta)|\tau\zeta < 1$. Define a relation \perp on Q by

$$\theta \perp \mu \text{ if } \theta(\zeta)\mu(\zeta) \leq \theta(\zeta) \vee \mu(\zeta),$$

where $\theta(\zeta) \vee \mu(\zeta) = \theta(\zeta)$ or $\mu(\zeta)$. Define $\tau : Q \times Q \rightarrow \mathbb{R}^+$ by

$$\tau(\theta, \mu) = \left(\int_0^1 |\theta(\zeta) - \mu(\zeta)|\tau\zeta \right)^2.$$

Then, (Q, τ) is an \mathbf{O} -complete \mathbf{O}_b -MS with $g = 2$. The operator $E : Q \times Q \rightarrow \mathbb{R}^+$ is defined by

$$E\theta(\zeta) = \frac{1}{4} \ln(1 + |\theta(\zeta)|).$$

Consider the map $\alpha : Q \times Q \rightarrow \mathbb{R}^+$, with $\lambda : \mathbb{R}^+ \rightarrow [0, \frac{1}{2})$ and $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, defined by

$$\alpha(\theta, \mu) = \begin{cases} 1, & \text{if } \theta(\zeta) \geq \mu(\zeta), \forall \zeta \in [0, 1], \\ 0, & \text{otherwise.} \end{cases}$$

$$\lambda(\zeta) = \frac{(\ln(1 + \sqrt{\zeta}))^2}{2\zeta} \text{ and } \psi(\zeta) = \zeta.$$

Obviously, $\psi \in \Lambda$, and $\lambda \in \Gamma$. Moreover, E is an \mathbf{O} -triangular- α orbital admissible map, and $\alpha(1, E1) \geq 1$.

Now, we prove E is an \mathbf{O} -generalized α - ψ -GC type(A) map. Certainly, $\forall \zeta \in [0, 1]$, we get

$$\begin{aligned} \sqrt{\alpha(\theta(\zeta), \mu(\zeta))\psi(g^3\tau(E\theta(\zeta), E\mu(\zeta)))} &\leq \sqrt{2^3 \left(\int_0^1 |E\theta(\zeta) - E\mu(\zeta)|\tau\zeta \right)^2} \\ &\leq 2\sqrt{2} \int_0^1 \left| \frac{1}{4} \ln(1 + |\theta(\zeta)|) - \frac{1}{4} \ln(1 + |\mu(\zeta)|) \right| \tau\zeta \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\sqrt{2}} \int_0^1 \left| \operatorname{In} \left(\frac{1 + |\theta(\zeta)|}{1 + |\mu(\zeta)|} \right) \right| \tau \zeta \\
&= \frac{1}{\sqrt{2}} \int_0^1 \left| \operatorname{In} \left(1 + \frac{|\theta(\zeta)| - |\mu(\zeta)|}{1 + |\mu(\zeta)|} \right) \right| \tau \zeta \\
&\leq \frac{1}{\sqrt{2}} \int_0^1 |\operatorname{In}(1 + |\theta(\zeta)| - |\mu(\zeta)|)| \tau \zeta,
\end{aligned}$$

and we have

$$\begin{aligned}
\int_0^1 |\operatorname{In}(1 + |\theta(\zeta)| - |\mu(\zeta)|)| \tau \zeta &\leq \operatorname{In} \left(\int_0^1 |(1 + |\theta(\zeta)| - |\mu(\zeta)|)| \tau \zeta \right) \\
&= \operatorname{In} \left(1 + \int_0^1 |\theta(\zeta) - \mu(\zeta)| \tau \zeta \right).
\end{aligned}$$

Therefore,

$$\begin{aligned}
\sqrt{\alpha(\theta(\zeta), \mu(\zeta)) \psi(g^3 \tau(E\theta(\zeta), E\mu(\zeta)))} &\leq \frac{1}{\sqrt{2}} \operatorname{In} \left(1 + \int_0^1 |\theta(\zeta) - \mu(\zeta)| \tau \zeta \right) \\
&\leq \frac{1}{\sqrt{2}} \operatorname{In} \left(1 + \sqrt{\tau(\theta, \mu)} \right).
\end{aligned}$$

So, we get

$$\begin{aligned}
\alpha(\theta(\zeta), \mu(\zeta)) \psi(g^3 \tau(E\theta(\zeta), E\mu(\zeta))) &\leq \frac{1}{2} (\operatorname{In}(1 + \sqrt{\tau(\theta, \mu)}))^2 \\
&\leq \frac{1}{2} (\operatorname{In}(1 + \sqrt{\mathbf{M}(\theta, \mu)}))^2 \\
&\leq \frac{(\operatorname{In}(1 + \sqrt{\mathbf{M}(\theta, \mu)}))^2}{2\mathbf{M}(\theta, \mu)} \mathbf{M}(\theta, \mu) \\
&= \lambda(\psi(\mathbf{M}(\theta, \mu))) \psi(\mathbf{M}(\theta, \mu)).
\end{aligned}$$

Hence by Theorem 3.3, we get that E has a UFP.

4. Application

As an application of Theorem 3.2, we find an existence and uniqueness result of the following type of integral equation:

$$\omega(\theta) = \lambda(\theta) + \int_0^{\alpha} E(\theta, s) \mathcal{H}(\theta, s, \omega(s)) ds, \quad \theta \in [0, \alpha], \alpha > 0. \quad (4.1)$$

Consider $\mathcal{Q} = C([0, \alpha], \mathbb{R})$ to be the real continuous functions on $[0, \alpha]$, and a mapping $\mathcal{D} : \mathcal{Q} \rightarrow \mathcal{Q}$ is defined by

$$\tau(\omega, \mu) = \max_{0 \leq \theta \leq \alpha} |\omega(\theta) - \mu(\theta)|^2, \quad \omega, \mu \in \mathcal{Q}. \quad (4.2)$$

Obviously, (\mathcal{Q}, τ) is a complete b -metric space, and $\omega(\theta)$ is a solution of integral equation (4.1) iff $\omega(\theta)$ is a fixed point of \mathcal{D} .

Theorem 4.1. *Suppose the following.*

(1) *The mappings $E : [0, \alpha] \times \mathbb{R} \rightarrow \mathbb{R}^+$, $\mathcal{H} : [0, \alpha] \times \mathbb{R} \rightarrow \mathbb{R}$, and $\lambda : [0, \alpha] \rightarrow \mathbb{R}$ are O -continuous functions.*

(2) *There exists $\mathcal{K} > 0$, such that, for all $\theta, s \in [0, \alpha]$ and $\omega, \mu \in \mathcal{Q}$,*

$$|\mathcal{H}(\theta, s, \omega(s)) - \mathcal{H}(\theta, s, \mu(s))| \leq \sqrt{\frac{e^{-\mathcal{K}(\omega, \mu)} \mathcal{K}(\omega, \mu)}{2}}. \quad (4.3)$$

(3) *For all $\theta, s \in [0, \alpha]$, we have*

$$\max \int_0^\alpha E(\theta, s)^2 ds \leq \frac{1}{\alpha}.$$

Then, (4.1) has a unique solution in \mathcal{Q} .

Proof. Define the O -relation \perp on \mathcal{Q} by

$$\omega \perp \mu \iff \omega(\theta)\mu(\theta) \geq \omega(\theta) \quad \text{or} \quad \omega(\theta)\mu(\theta) \geq \mu(\theta), \quad \forall \theta \in [0, \alpha].$$

Define $\tau : \mathcal{Q} \times \mathcal{Q} \rightarrow \mathbb{R}^+$, given by

$$\tau(\omega, \mu) = \max_{0 \leq \theta \leq \alpha} |\omega(\theta) - \mu(\theta)|^2,$$

for all $\omega, \mu \in \mathcal{Q}$. It is easy to see that, $(\mathcal{Q}, \perp, \tau)$ is an \mathbf{O} -complete \mathbf{O}_b -MS. For all $\omega, \mu \in \mathcal{Q}$ with $\omega \perp \mu$ and $\theta \in [0, \alpha]$, we have

$$\mathcal{D}(\omega(\theta)) = \lambda(\theta) + \int_0^\alpha E(\theta, s) \mathcal{H}(\theta, s, \omega(s)) ds \geq 1. \quad (4.4)$$

Accordingly, $[(\mathcal{D}\omega)(\theta)][(\mathcal{D}\mu)(\theta)] \geq (\mathcal{D}\mu)(\theta)$, and so $(\mathcal{D}\omega)(\theta) \perp (\mathcal{D}\mu)(\theta)$. Then, \mathcal{D} is \perp -preserving.

Let $\omega, \mu \in \mathcal{Q}$ with $\omega \perp \mu$. Suppose that $\mathcal{D}(\omega) \neq \mathcal{D}(\mu)$. For each $\theta \in [0, \alpha]$, we have

$$\begin{aligned} \tau(\mathcal{D}\omega, \mathcal{D}\mu) &= \max_{\theta \in [0, \alpha]} |\mathcal{D}\omega(\theta) - \mathcal{D}\mu(\theta)|^2 \\ &= \max_{\theta \in [0, \alpha]} \left\{ \left| \lambda(\theta) + \int_0^\alpha E(\theta, s) \mathcal{H}(\theta, s, \omega(s)) ds - \lambda(\theta) - \int_0^\alpha E(\theta, s) \mathcal{H}(\theta, s, \mu(s)) ds \right|^2 \right\} \\ &= \max_{\theta \in [0, \alpha]} \left\{ \left| \int_0^\alpha E(\theta, s) (\mathcal{H}(\theta, s, \omega(s)) - \mathcal{H}(\theta, s, \mu(s))) ds \right|^2 \right\} \\ &\leq \max_{\theta \in [0, \alpha]} \left\{ \int_0^\alpha E(\theta, s)^2 ds \int_0^\alpha |\mathcal{H}(\theta, s, \omega(s)) - \mathcal{H}(\theta, s, \mu(s))|^2 ds \right\} \\ &\leq \frac{1}{\alpha} \int_0^\alpha \left| \sqrt{\frac{e^{-\mathcal{K}(\omega, \mu)} \mathcal{K}(\omega, \mu)}{2}} \right|^2 ds \\ &\leq \frac{e^{-\mathcal{K}(\omega, \mu)}}{2} \mathcal{K}(\omega, \mu). \end{aligned}$$

Thus,

$$\tau(\mathcal{D}\omega, \mathcal{D}\mu) \leq \gamma(\mathcal{K}(\omega, \mu)) \mathcal{K}(\omega, \mu),$$

for all $\omega, \mu \in \mathcal{Q}$. Therefore, all the conditions of Theorem (3.2) are satisfied. Hence, (4.1) has a unique solution. \square

Example 4.2. Consider the integral equation as follows:

$$g(\theta) = \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \int_0^x \theta^2 \sigma g(\sigma) \delta\sigma, \quad 0 \leq x \leq 1. \quad (4.5)$$

Clearly, the above Eq (4.5) satisfies the assumption of Theorem 4.1, that is, $\sin(\pi\theta^2) - \frac{\theta^2}{\pi}$ is an orthogonal continuous function on $[0, 1]$. Kernel $K(\theta, \sigma)$ is orthogonal continuous on $\mathbb{R} = \{(\theta, \sigma), 0 < \theta, \sigma < 1\}$.

The solution will be determined from Eq (4.5) by the fixed point iteration method:

$$g_{\theta+1}(\theta) = \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \int_0^x \theta^2 \sigma g_{\theta}(\sigma) \delta\sigma, \quad 0 \leq x \leq 1.$$

By choosing $\sin(\pi\theta^2) - \frac{\theta^2}{\pi}$ as the initial function, we can apply the fixed point iteration method to get a numerical solution:

$$\begin{aligned} g_1(\theta) &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \int_0^x \theta^2 \sigma g_0(\sigma) \delta\sigma \\ &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \int_0^x \theta^2 \sigma \sin(\pi\sigma^2) \delta\sigma \\ &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \theta^2 \frac{1}{2\pi} (1 - \cos(\pi\theta^2)), \\ g_2(\theta) &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \int_0^x \theta^2 \sigma g_1(\sigma) \delta\sigma \\ &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \int_0^x \theta^2 \sigma \left(\sin(\pi\sigma^2) - \frac{\sigma^2}{\pi} + \sigma^2 \frac{1}{2\pi} (1 - \cos(\pi\sigma^2)) \right) \delta\sigma \\ &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \frac{\theta^2}{8\pi^3} \left(-4\pi^2 - 2 + 4\pi^2 \cos(\pi\theta^2) + \pi^2 \theta^4 + 2\theta^2 \pi \sin(\pi\theta^2) + 2 \cos(\pi\theta^2) \right), \\ g_3(\theta) &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \int_0^x \theta^2 \sigma g_2(\sigma) \delta\sigma \\ &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \int_0^x \theta^2 \sigma \left(\sin(\pi\sigma^2) - \frac{\sigma^2}{\pi} + \frac{\sigma^2}{8\pi^2} (-4\pi^2 - 2 + 4\pi^2 \cos(\pi\sigma^2) + \sigma^4 \pi^2 \right. \\ &\quad \left. + 2\theta^2 \pi \sin(\pi\theta^2) + 2 \cos(\pi\theta^2)) \right) \delta\sigma \\ &= \sin(\pi\theta^2) - \frac{\theta^2}{\pi} + \theta^2 \left\{ -\frac{1}{64\pi^6} \left(-32\pi^5 - 16\pi^3 - 24\pi + 32\pi^5 \cos(\pi\theta^2) + 8\theta^4 \pi^5 - 4\theta^4 \pi^3 \right. \right. \\ &\quad \left. \left. + 16\theta^2 \pi^4 (\sin(\pi\theta^2) + 16\pi^3 \cos(\pi\theta^2) + \theta^8 \pi^5 - 8\theta^4 \pi^3 \cos(\pi\theta^2) + 24\theta^2 \pi^2 \sin(\pi\theta^2) \right. \right. \\ &\quad \left. \left. + 24\pi \cos(\pi\theta^2) \right) \right\}. \end{aligned}$$

Consider that for $|\theta| \leq 1$, an O -sequence $\{g_{\theta}(\theta)\}$ will converge to $g(\theta) = \sin(\pi\theta^2) - \frac{\theta^2}{\pi}$.

Error calculation for an approximate solution compared to an exact solution is given in Figure 1. Table 1 shows that the error of an approximate solution compared to an exact solution is relatively small.

Table 1. Comparison of an approximate solution and an exact solution.

θ_j	approximate solution	exact solution	error
0.000	0.000	0.000	0.000
0.100	0.023	0.028	0.005
0.200	0.102	0.113	0.011
0.300	0.234	0.250	0.016
0.400	0.412	0.431	0.019
0.500	0.609	0.628	0.018
0.600	0.779	0.790	0.011
0.700	0.848	0.844	0.004
0.800	0.730	0.701	0.029
0.900	0.358	0.304	0.054
1.000	-0.251	-0.318	0.067




Figure 1. Graph of an approximate solution compare to an exact solution with $h=0.1$.

5. Conclusions

In this paper, we proved fixed point theorems for \mathbf{O} -generalized α - ψ -GC type contraction mappings in an \mathbf{O} -complete \mathbf{O}_b -MS. Furthermore, we presented some examples to strengthen our main results. Also, we provided an application to the existence of the solution of an integral equation and we have compared the approximate solution with the exact solution.

Khalehoghli et al. [30, 31] presented a real generalization of the mentioned Banach's contraction principle by introducing \mathbb{R} -metric spaces, where \mathbb{R} is an arbitrary relation on L . We note that in a special case, \mathbb{R} can be considered as $\mathbb{R} = \leq$ [partially ordered relation], $\mathbb{R} = \perp$ [orthogonal relation], etc. If one can find a suitable replacement for a Banach theorem that may determine the values of fixed

points, then many problems can be solved in this \mathbb{R} -relation. This will provide a structural method for finding a value of a fixed point. It is an interesting open problem to study the fixed-point results on \mathbb{R} -complete \mathbb{R} -metric spaces.

Conflict of interest

The authors declare that they have no competing interests.

References

1. S. Banach, Sur les opérations dans ensembles abstraits et leur application aux équations intégrales, *Fund. Math.*, **3** (1922), 51–57.
2. S. Czerwik, Contraction mappings in b -metric spaces, *Acta. Math. Inform. Univ. Ostrav.*, **1** (1993), 5–11.
3. H. Aydi, M. Bota, E. Karapinar, S. Moradi, A common fixed points for weak ψ -contractions on b -metric spaces, *Fixed Point Theory*, **13** (2012), 337–346.
4. M. Pacurar, A fixed point result for ψ -contractions and fixed points on b -metric spaces without the boundness assumption, *Fasc. Math.*, **43** (2010), 127–136.
5. M. B. Zada, M. Sarwar, P. Kumam, Fixed point results for rational type contraction in b -metric spaces, *Int. J. Anal. Appl.*, **16** (2018), 904–920.
6. M. A. Geraghty, On contractive mappings, *Proc. Amer. Math. Soc.*, **40** (1973), 604–608. <https://doi.org/10.1090/S0002-9939-1973-0334176-5>
7. B. Samet, C. Vetro, P. Vetro, Fixed point theorems for α - ψ -contractive type mappings, *Nonlinear Anal.: Theory Methods Appl.*, **75** (2012), 2154–2165. <https://doi.org/10.1016/j.na.2011.10.014>
8. S. H. Cho, J. S. Bae, E. Karapinar, Fixed point theorems for α -Geraghty contraction type maps in metric spaces, *Fixed Point Theory Appl.*, **2013** (2013), 329. <https://doi.org/10.1186/1687-1812-2013-329>
9. O. Popescu, Some new fixed point theorems for α -Geraghty contraction type mappings in metric spaces, *Fixed Point Theory Appl.*, **2014** (2014), 190. <https://doi.org/10.1186/1687-1812-2014-190>
10. E. Karapinar, P. Kumam, P. Salimi, On α - ψ -Meir-Keeler contractive mappings, *Fixed Point Theory Appl.*, **2013** (2013), 94. <https://doi.org/10.1186/1687-1812-2013-94>
11. E. Karapinar, A discussion on $(\alpha$ - ψ)-Geraghty contraction type maps, *Filomat*, **28** (2014), 761–766.
12. E. Karapinar, $(\alpha$ - ψ)-Geraghty contraction type mappings and some related fixed point results, *Filomat*, **28** (2014), 37–48. <https://doi.org/10.2298/FIL1401037K>
13. M. Mehmood, H. Aydi, M. U. Ali, Fahimuddin, A. Shoaib, M. De La Sen, Solutions of integral equations via fixed-point results on orthogonal gauge structure, *Math. Probl. Eng.*, **2021** (2021), 8387262. <https://doi.org/10.1155/2021/8387262>

14. H. Guan, J. Li, Y. Hao, Common fixed point theorems for weakly contractions in rectangular b -metric spaces with supportive applications, *J. Funct. Spaces*, **2022** (2022), 8476040. <https://doi.org/10.1155/2022/8476040>
15. L. Chen H. Guan, Common fixed point and coincidence point results for generalized α - φ_E -Geraghty contraction mappings in b -metric spaces, *AIMS Math.*, **7** (2022), 14513–14531. <https://doi.org/10.3934/math.2022800>
16. H. Guan, J. Li, Common fixed-point theorems of generalized (ψ, φ) weakly contractive mappings in b -metric-like spaces and application, *J. Math.*, **2021** (2021), 6680381. <https://doi.org/10.1155/2021/6680381>
17. Y. Hao, H. Guan, On some common fixed point results for weakly contraction mappings with application, *J. Funct. Spaces*, **2021** (2021), 5573983. <https://doi.org/10.1155/2021/5573983>
18. J. Li, H. Guan, Common fixed point results for generalized $(g - \alpha_{sp}, \psi, \phi)$ contractive mappings with applications, *J. Funct. Spaces*, **2021** (2021), 5020027. <https://doi.org/10.1155/2021/5020027>
19. F. Zhang, X. Zhang, Y. Hao, Common fixed point theorems for contractive mappings of integral type in G -metric spaces and applications, *J. Funct. Spaces*, **2021** (2021), 6619964. <https://doi.org/10.1155/2021/6619964>
20. M. E. Gordji, M. Ramezani, M. De La Sen, Y. J. Cho, On orthogonal sets and Banach fixed point theorem, *Fixed Point Theory*, **18** (2017), 569–578. <https://doi.org/10.24193/fpt-ro.2017.2.45>
21. M. E. Gordji, H. Habibi, Fixed point theory in generalized orthogonal metric space, *J. Linear Topol. Algebra*, **6** (2017), 251–260.
22. A. J. Gnanaprakasam, G. Mani, J. R. Lee, C. Park, Solving a nonlinear integral equation via orthogonal metric space, *AIMS Math.*, **7** (2022), 1198–1210. <https://doi.org/10.3934/math.2022070>
23. G. Mani, A. J. Gnanaprakasam, N. Kausar, M. Munir, Salahuddin, Orthogonal F -contraction mapping on O -complete metric space with applications, *Int. J. Fuzzy Log. Intell. Syst.*, **21** (2021), 243–250. <https://doi.org/10.5391/IJFIS.2021.21.3.243>
24. G. Mani, A. J. Gnanaprakasam, C. Park, S. Yun, Orthogonal F -contractions on O -complete b -metric space, *AIMS Math.*, **6** (2021), 8315–8330. <https://doi.org/10.3934/math.2021481>
25. A. J. Gnanaprakasam, G. Mani, V. Parvaneh, H. Aydi, Solving a nonlinear Fredholm integral equation via an orthogonal metric, *Adv. Math. Phys.*, **2021** (2021), 1202527. <https://doi.org/10.1155/2021/1202527>
26. M. Ramezani, Orthogonal metric space and convex contractions, *Int. J. Nonlinear Anal. Appl.*, **6** (2015), 127–132. <http://dx.doi.org/10.22075/ijnaa.2015.261>
27. S. K. Prakasam, A. J. Gnanaprakasam, N. Kausar, G. Mani, M. Munir, Salahuddin, Solution of integral equation via orthogonally modified F -contraction mappings on O -complete metric-like space, *Int. J. Fuzzy Logic Intell. Syst.*, **22** (2022), 287–295. <http://doi.org/10.5391/IJFIS.2022.22.3.287>

-
28. H. Afshari, H. Aydi, E. Karapınar, On generalized α - ψ -Geraghty contractions on b -metric spaces, *Georgian Math. J.*, **27** (2018), 9–21. <http://doi.org/10.1515/gmj-2017-0063>
29. S. K. Prakasam, A. J. Gnanaprakasam, O. Ege, G. Mani, S. Haque, N. Mlaiki, Fixed point for an $\mathbb{O}g\mathfrak{F}$ -c in O -complete b -metric-like spaces, *AIMS Math.*, **8** (2022), 1022–1039, <http://doi.org/10.3934/math.2023050>
30. S. Khalehoghli, H. Rahimi, M. E. Gordji, Fixed point theorems in R -metric spaces with applications, *AIMS Math.*, **5** (2020), 3125–3137. <https://doi.org/10.3934/math.2020201>
31. S. Khalehoghli, H. Rahimi, M. E. Gordji, R -topological spaces and SR -topological spaces with their applications, *Math. Sci.*, **14** (2020), 249–255. <https://doi.org/10.1007/s40096-020-00338-5>



AIMS Press

©2023 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)