



Research article

Dynamics of drug on-drug off models with mutations in morbidostat — Dedicated to the seventieth birthday of Professor Gail Wolkowicz

Ziran Cheng¹ and Sze-Bi Hsu^{2,*}

¹ Shen Zhen senior high school, Shenzhen, Guan-Dong, China

² Department of Mathematics, National Tsing-Hua University, Taiwan

* **Correspondence:** Email: sbhsu@math.nthu.edu.tw.

Abstract: The morbidostat is a bacteria culture device that progressively increases antibiotic drug concentration. It is used to study the evolutionary pathway. In this article, we construct mathematical models for the morbidostat. First we consider the case of no mutations, we study limiting systems and obtain criteria for the large time behavior of the solutions. From the theoretical results and numerical simulations, we conclude that there are two competitive exclusion states of either wild type or mutant type as the threshold parameter U varies. There are three cases, wild type bacteria excludes all mutants; a mutant dominates in the competition; oscillation between the above two states.

Next we study the systems of forward mutations and forward-backward mutations. Then we apply a result of perturbation for globally stable state.

Keywords: morbidostat; chemostat; wild type; mutants; drug inhibition; exclusion principle; coexistence; differential inequalities; impulse systems

Mathematics Subject Classification: 34C, 92B, 94D

1. Introduction and models

1.1. Introduction

Antibiotic drug resistance is a global health problem [9]. Five years after the clinical application of penicillin, staphylococcus aureus with penicillin resistance [8] was found in humans, which caused penicillin to fail in some patients. These years, human beings may face the dilemma of infecting multiple drug-resistant bacteria (commonly known as superbugs) without any help of antibiotics because of the antibiotic drug resistance.

In [10, 11] the authors presented a devices for building and operating an automated fluid system for continuous culture called “morbidostat” (See Figure 1.1). The morbidostat is used to study evolution of microbial drug resistance in real time, It constantly measure the growth rates of evolving microbial

populations inside culture in order to maintain a constant drug-induced inhibition. The growth rate measurements are done by using an optical detection system. Similar to chemostat rate D lower than the maximal growth rate of population in morbidostat. In contrast to a chemostat in which the bacterial growth is inherently the cell density is kept low such that the population is not nutrient limited, its growth rate is also controlled by externally adjusting drug concentration.

In this work, we follow the same mathematical model of morbidostat in [5, 6]. When the total population of all bacteria are less than a threshold value U , there is no drug injection into the morbidostat and $\frac{dP}{dt} = -DP$. However if the total population of all bacteria reaches over the threshold value U , there will be continuous drug injection into morbidostat. The growth rate of species i will be $g_i(S)f_i(P)$ where S is the concentration of nutrient, $g_i(S)$ takes the form of Holling type II $g_i(S) = \frac{m_i S}{a_i + S}$ and P is the concentration of drug, the inhibition $f_i(P)$ takes the form of Hill function $f_i(P) = \frac{1}{1 + (\frac{P}{K_i})^L}$ and $\frac{dP}{dt} = D(P^{(0)} - P)$.

Remark: We note that in [5, 6], we assume

$$f_0(P)g_0(S) \leq f_1(P)g_1(S) \leq \dots \leq f_n(P)g_n(S) \quad (H^*)$$

i.e, the n -th mutant has largest growth rate.

In [6] we consider periodic resetting for nutrient and drug; while in [5] we consider drug on-drug off mechanism.

In this paper we relax the assumption (H^*) with drug on-drug off dynamics in the morbidostat.




Figure 1. A schematic of morbidostat.

1.2. Description of the models

In this work, we analyze the transition between the population of wild type u and those of mutant v_i ($i = 1, 2, \dots, n$). Consider the drug on-drug off model with forward mutations (see Figure 1.2) and forward-backward mutations (see Figure 1.3).



Figure 2. Forward mutations between species.




Figure 3. Forward-backward mutations between species.

The growth dynamics with the nutrient substrate S in a chemostat under the influence of the drug inhibitor P with forward mutations and forward-backward mutations are given by (1) and (2) respectively:

$$\begin{cases} \frac{dS}{dt} = (S^{(0)} - S)D - \frac{1}{\gamma}g_0(S)f_0(P)u - \frac{1}{\gamma}\sum_{i=1}^n g_i(S)f_i(P)v_i \\ \frac{du}{dt} = (g_0(S)f_0(P) - D)u - q_0u \\ \frac{dv_i}{dt} = (g_i(S)f_i(P) - D)v_i + q_{i-1}v_{i-1} - q_iv_i \\ \frac{dv_n}{dt} = (g_n(S)f_n(P) - D)v_n + q_{n-1}v_{n-1} \\ \frac{dP}{dt} = \begin{cases} -DP, & \text{if } u + \sum_{j=1}^n v_j < U \\ (P^{(0)} - P)D, & \text{if } u + \sum_{j=1}^n v_j \geq U \end{cases} \end{cases} \quad (1)$$

$$\begin{cases} \frac{dS}{dt} = (S^{(0)} - S)D - \frac{1}{\gamma}g_0(S)f_0(P)u - \frac{1}{\gamma}\sum_{i=1}^n g_i(S)f_i(P)v_i \\ \frac{du}{dt} = (g_0(S)f_0(P) - D)u - q_0u + \tilde{q}_0v_1 \\ \frac{dv_i}{dt} = (g_i(S)f_i(P) - D)v_i + q_{i-1}v_{i-1} - q_iv_i - \tilde{q}_{i-1}v_i + \tilde{q}_iv_{i+1} \\ \frac{dv_n}{dt} = (g_n(S)f_n(P) - D)v_n + q_{n-1}v_{n-1} - \tilde{q}_{n-1}v_n \\ \frac{dP}{dt} = \begin{cases} -DP, & \text{if } u + \sum_{j=1}^n v_j < U \\ (P^{(0)} - P)D, & \text{if } u + \sum_{j=1}^n v_j \geq U \end{cases} \end{cases} \quad (2)$$

where $i = 1, 2, 3, \dots, n-1$. Let $v_0 = u$ and v_i are the volume densities of the wild type and mutant populations, respectively. γ denotes the yield constant, reflecting the conversion of nutrient to bacteria. S is the concentration of nutrient, while $S^{(0)}$ is the input concentration of nutrient. D is the dilution rate. $g_0(S)$ and $g_i(S)$ are the growth rates of the wild type and mutants, which satisfy $g_i(0) = 0$, $g'_i(S) > 0$ for $i = 0, 1, 2, \dots, n$. It implies that bacteria grows only when there has nutrient substrate in the device, and a higher concentration of nutrient leads to the bacteria's higher growth rates. Here we consider

the important special case $g_i(S) = \frac{m_i S}{a_i + S}$. q_i denotes the forward mutation from v_i to v_{i+1} , and \tilde{q}_i is the backward mutation from v_{i+1} to v_i . q_i and \tilde{q}_i are small positive quantities. Besides, the drug inhibitions for v_i are described by $f_i(P)$ for $i = 0, 1, 2, \dots, n$. It is easy to know, the higher concentration of drug leads to a stronger inhibition to the growth of bacteria. Based on the fact that mutants always have stronger resistances to the inhibitor than wild-type, and the i -th mutant v_i has stronger resistances than the $(i - 1)$ -th mutant v_{i-1} . So we assume

$$f_i(0) = 1, f'_i(P) < 0, f_0(P) \leq f_1(P) \leq \dots \leq f_n(P) \quad (H1)$$

for $i = 0, 1, 2, \dots, n$.

Let $g_i(\lambda_i) = D$, i.e. $\lambda_i = \frac{a_i}{(\frac{m_i}{D} - 1)}$. We also assume that:

$$\lambda_0 < \lambda_1 < \dots < \lambda_n < S^{(0)} \quad (H2)$$

which implies the wild type u which has smallest break-even concentration, is the most superior species than the mutants v_i in the absence of drug inhibition [1].

The rest of this paper is organized as follows. In Section 2 we study the dynamics of drug on-drug off models with no mutations, In Section 3 we study the dynamics of drug on-drug off models with forward mutations. In Section 4 we study the dynamics of drug on-drug off models with forward-backward mutations. Section 5 is a section of numerical simulations for various threshold parameters U . Section 6 is the discussion section.

2. Dynamics of drug on-drug off model with no mutations

Without loss of generality by scaling, we assume the yield constant, $\gamma = 1$.

Lemma 2.1. (Conservation Property) $S(t) + \sum_{i=0}^n v_i(t) \rightarrow S^{(0)}$ as $t \rightarrow \infty$.

Proof. Let $M(t) = S(t) + \sum_{i=0}^n v_i(t)$.

Adding the first $n + 2$ equations of (1) and (2), yields

$$\frac{dM(t)}{dt} = (S^{(0)} - S)D - D \sum_{i=0}^n v_i = (S^{(0)} - M(t))D$$

So $M(t) = S(t) + \sum_{i=0}^n v_i(t) \rightarrow S^{(0)}$ as $t \rightarrow \infty$. □

For simplicity in (1) we assume that $q_i = 0$ for all $i = 0, 1, 2, \dots, n$. Let $u = v_0$, the system (1) becomes

$$\begin{cases} \frac{dS}{dt} = (S^{(0)} - S)D - \sum_{i=0}^n g_i(S)f_i(P)v_i \\ \frac{dv_i}{dt} = (g_i(S)f_i(P) - D)v_i, \quad i = 0, 1, 2, \dots, n \\ \frac{dP}{dt} = \begin{cases} -DP, & \text{if } \sum_{j=0}^n v_j < U \\ (P^{(0)} - P)D, & \text{if } \sum_{j=0}^n v_j \geq U \end{cases} \end{cases} \quad (3)$$

$$S(0) > 0, v_i(0) > 0, i = 0, 1, \dots, n, P(0) > 0.$$

Next we state two theorems which will be used in Section 2.1 below.

Theorem A (Competitive exclusion in simple chemostat) ([1] p. 30 or p. 35)

Consider the simple chemostat equation

$$\begin{aligned} \frac{dS}{dt} &= (S^{(0)} - S)D - \sum_{i=0}^n g_i(S)v_i \\ \frac{dv_i}{dt} &= (g_i(S) - D)v_i, i = 0, 1, 2, \dots, n, \end{aligned}$$

with $S(0) \geq 0$ and $v_i(0) > 0$. Let (H2) hold. Then

$$\begin{aligned} \lim_{t \rightarrow \infty} S(t) &= \lambda_0 \\ \lim_{t \rightarrow \infty} v_0(t) &= v_0^* = S^{(0)} - \lambda_0 \\ \lim_{t \rightarrow \infty} v_i(t) &= 0, i = 1, 2, \dots, n. \end{aligned} \quad \square$$

Consider two systems of ordinary differential equations of the form

$$z' = Az, \quad y' = f(y, z) \tag{F.1}$$

and

$$x' = f(x, 0) \tag{F.2}$$

where

$$\begin{aligned} z &\in \mathbb{R}^m, (y, z) \in D \subset \mathbb{R}^n \times \mathbb{R}^m \\ x &\in \Omega = \{x \mid (x, 0) \in D\} \subseteq \mathbb{R}^n. \end{aligned}$$

We assume A is an $m \times m$ matrix, f is continuous differentiable, D is positively invariant for (F.1) and (F.2) is dissipative. Let (B1)-(B5) be additional hypothesis:

- (B1) All of eigenvalues of A have negative real parts.
- (B2) Equation (F.2) has finite number of rest points in Ω , each of which is hyperbolic ([2], p. 88) for (F.2). Denote these rest points by x_1, \dots, x_p .
- (B3) The dimension of the stable manifold ([2], p. 88) is n for $1 \leq i \leq r$, and the dimension of the stable manifold of x_j is less than n for $j = r + 1, \dots, p$. In symbols, $\dim(M^+(x_j)) = n, i = 1, \dots, r$; $\dim(M^+(x_j)) < n$ for $j = r + 1, \dots, p$.
- (B4) $\Omega = \cup_{i=1}^p M^+(x_j)$.
- (B5) Equation (F.2) does not possess a cycle of rest points.

Theorem B (A convergence theorem)([1], p. 294)

Let (B1)-(B5) hold and $(y(t), z(t))$ be a solution of (F.1). Then, for some i ,

$$\lim_{t \rightarrow \infty} (y(t), z(t)) = (x_i, 0).$$

2.1. The behavior of solutions of the system (3)

For the solution of the system (3), we discuss the following three cases:

Case 1 : $\sum_{i=0}^n v_i(t) < U$ for all $t > t_0 > 0$ for some $t_0 > 0$. Then we have

$$\frac{dP}{dt} = -DP, \text{ for } t \geq t_0$$

it is obvious that

$$P(t) \longrightarrow 0 \text{ as } t \longrightarrow \infty$$

Let

$$\Sigma_1(t) = S^{(0)} - S(t) - \sum_{i=0}^n v_i(t)$$

$$\Sigma_2(t) = P(t).$$

Then by Lemma 2.1, we have

$$\Sigma_1'(t) = -D\Sigma_1(t)$$

$$\Sigma_2'(t) = -D\Sigma_2(t)$$

and $\Sigma_1(t) \rightarrow 0$ and $\Sigma_2(t) \rightarrow 0$ as $t \rightarrow \infty$. Rewrite the system (3) with $\frac{dP}{dt} = -DP$ as

$$\begin{aligned} \Sigma_1'(t) &= -D\Sigma_1(t) \\ \Sigma_2'(t) &= -D\Sigma_2(t) \\ v_i'(t) &= \left(g_i \left(S^{(0)} - \Sigma_1(t) - \sum_{i=0}^n v_i(t) \right) f_i(\Sigma_2(t)) - D \right) v_i, i = 0, 1, 2, \dots, n. \end{aligned} \tag{2.1}$$

$$D = \left\{ (\Sigma_1, \Sigma_2, v_0, \dots, v_n) : v_i > 0, i = 0, 1, \dots, n, \sum_{i=0}^n v_i + \Sigma_1 \leq S^{(0)}, \Sigma_2 > 0 \right\}.$$

The system corresponding to (F.2) in Theorem B is

$$\begin{aligned} v_i'(t) &= \left(g_i \left(S^{(0)} - \sum_{i=0}^n v_i(t) \right) f_i(0) - D \right) v_i \\ &= \left(g_i \left(S^{(0)} - \sum_{i=0}^n v_i(t) \right) - D \right) v_i, i = 0, 1, 2, \dots, n. \end{aligned} \tag{2.2}$$

Apply Theorem B with $A = \text{diag}(-D, -D)$, $m = 2$ and

$$\Omega = \left\{ (v_0, v_1, \dots, v_n) : v_i > 0, \sum_{i=1}^n v_i \leq S^{(0)} \right\}.$$

Under the hypothesis (H2), we apply Theorem A and obtain a result of global stability,

$$\begin{aligned} \lim_{t \rightarrow \infty} v_0(t) &= v_0^* = S^{(0)} - \lambda_0 \\ \lim_{t \rightarrow \infty} v_i(t) &= 0, i = 1, 2, \dots, n. \end{aligned}$$

Now we verify the hypothesis (B1)-(B5) holds for the systems (2.1) and (2.2). Obviously the 2×2 matrix A satisfies hypothesis (B1). For hypotheses (B2), (B3), (B4), equation (2.2) has finite number of rest points in Ω , namely, $O = (0, 0, \dots, 0)$, $E_0 = (v_0^*, 0, \dots, 0)$, $E_i = (0, \dots, v_i^*, \dots, 0)$, $v_i^* = S^{(0)} - \lambda_i$, $i = 1, 2, \dots, n$. $\dim(M^+(E_0)) = n + 1$, $\dim(M^+(E_i)) = n - (i - 1)$, $i = 1, 2, \dots, n$. To verify the cyclic condition (B5), we note that from (H2) E_j is chained to E_i , $E_j \rightarrow E_i$, $j > i$, $i, j = 0, 1, 2, \dots, n$ if and only if $\lambda_j < \lambda_i$ and $E_i \rightarrow O$ for all $i = 1, 2, \dots, n$. Suppose there is a cycle in Ω , $E_{k(1)} \rightarrow E_{k(2)} \rightarrow \dots \rightarrow E_{k(1)}$ then it follows that $\lambda_{k(1)} < \lambda_{k(2)} < \dots < \lambda_{k(1)}$ which is obviously a contradiction to (H2)

Case 2 : $\sum_{i=0}^n v_i(t) \geq U$ for all $t > t_0 > 0$ for some $t_0 \geq 0$. Then we have

$$\frac{dP}{dt} = D(P^{(0)} - P)$$

we have

$$P(t) \rightarrow P^{(0)} \text{ as } t \rightarrow \infty.$$

Let

$$\Sigma_1(t) = S^{(0)} - S(t) - \sum_{i=0}^n v_i(t)$$

$$\Sigma_2(t) = P^{(0)} - P(t)$$

Then we have

$$\Sigma_1'(t) = -D\Sigma_1(t)$$

$$\Sigma_2'(t) = -D\Sigma_2(t)$$

and $\Sigma_1(t) \rightarrow 0$ and $\Sigma_2(t) \rightarrow 0$ as $t \rightarrow \infty$. Rewrite the system (3) with $\frac{dP}{dt} = (P^{(0)} - P)$,

$$\begin{aligned} \Sigma_1'(t) &= -D\Sigma_1(t) \\ \Sigma_2'(t) &= -D\Sigma_2(t) \\ v_i'(t) &= \left(g_i \left(S^{(0)} - \Sigma_1(t) - \sum_{i=0}^n v_i(t) \right) f_i(P^{(0)} - \Sigma_2(t)) \right) v_i, i = 0, 1, 2, \dots, n. \end{aligned} \quad (2.3)$$

$$D = \left\{ (\Sigma_1, \Sigma_2, v_0, \dots, v_n) : v_i > 0, i = 0, \dots, n, \sum_{i=0}^n v_i + \Sigma_1 \leq S^{(0)}, \Sigma_2 > 0 \right\}.$$

The system corresponding to (F.2) in Theorem B is

$$v_i'(t) = \left(g_i \left(S^{(0)} - \sum_{i=0}^n v_i(t) \right) f_i(P^{(0)}) - D \right) v_i, i = 0, 1, 2, \dots, n. \quad (2.4)$$

We may verify the hypotheses (B1)-(B5) as in the Case 1.

If $m_i f_i(P^{(0)}) \leq D$, then it is easy to show $\lim_{t \rightarrow \infty} v_i(t) = 0$. Thus we assume that

$$m_i f_i(P^{(0)}) > D, \quad i = 0, 1, 2, \dots, n \quad (H3)$$

Replace m_i in $g_i(S)$ by $m_i f_i(P^{(0)})$, we have a new simple chemostat equation.

Let

$$\hat{\lambda}_k = \min_{0 \leq i \leq n} \hat{\lambda}_i, \text{ where } \hat{\lambda}_i = \frac{a_i}{\frac{m_i f_i(P^{(0)})}{D} - 1},$$

then

$$\begin{aligned} S(t) &\rightarrow \hat{\lambda}_k \\ v_k(t) &\rightarrow \hat{v}_k^* = S^{(0)} - \hat{\lambda}_k \\ v_i(t) &\rightarrow 0, i \neq k \end{aligned} \quad (2.5)$$

as $t \rightarrow \infty$ or

$$\lim_{t \rightarrow \infty} (v_0(t), v_1(t), \dots, v_n(t)) = (0, \dots, S^{(0)} - \hat{\lambda}_k, \dots, 0)$$

Case 3 : $\sum_{i=0}^n v_i < U$ at some sequence $\{t_k\}$, $t_k \rightarrow \infty$ and $\sum_{i=0}^n v_i \geq U$ at some other sequence $\{t_l\}$, $t_l \rightarrow \infty$, $k = 1, 2, \dots, l = 1, 2, \dots, k \neq l$, $\sum_{i=0}^n v_i$ oscillates around U .

2.2. The results of the asymptotic behavior of the solutions of model with no mutations

Lemma 2.2. Let $v_i, i = 0, \dots, n$ be the solution of system (2.4). Let (H1),(H2),(H3) hold, then $\hat{v}_k^* < v_0^*$.

Proof. Since

$$\hat{\lambda}_k = \frac{a_k}{\frac{m_k f_k(P^{(0)})}{D} - 1} \text{ and } \lambda_k = \frac{a_k}{\frac{m_k}{D} - 1}$$

and $0 < f_i(P^{(0)}) < 1$, then $\hat{\lambda}_k > \lambda_k$.

Obviously, we obtain

$$\hat{v}_k^* = S^{(0)} - \hat{\lambda}_k < S^{(0)} - \lambda_k < S^{(0)} - \lambda_0 = v_0^*$$

Hence,

$$\hat{v}_k^* < v_0^*$$

□

In the following Theorem 2.3 we shall give sufficient conditions for the Case 1, Case 2 and Case 3 for the solutions of the system (3).

Theorem 2.3. Let (H1), (H2), (H3) hold and $v_i, i = 0, 1, \dots, n$ be the solutions of the system (3), then

- (I) if $U > v_0^*$, then there exists $t_0 > 0$, such that for all $t \geq t_0$, $\sum_{i=0}^n v_i(t) < U$. From Case 1 in section 2.1, $\lim_{t \rightarrow \infty} S(t) = \lambda_0$, $\lim_{t \rightarrow \infty} u(t) = S^{(0)} - \lambda_0$, $\lim_{t \rightarrow \infty} v_i(t) = 0, i = 1, 2, \dots, n$.
- (II) Let $\hat{\lambda}_l = \max_{0 \leq i \leq n} \hat{\lambda}_i$ and $\hat{v}_l^* = S^{(0)} - \hat{\lambda}_l < \hat{v}_k^*$. If $U < \hat{v}_l^*$, then there exists $t_0 > 0$, such that for all $t \geq t_0$, $\sum_{i=0}^n v_i(t) \geq U$. From Case 2 in section 2.1, $\lim_{t \rightarrow \infty} S(t) = \hat{\lambda}_k$, $\lim_{t \rightarrow \infty} v_k(t) = \hat{v}_k^* = S^{(0)} - \hat{\lambda}_k$, $\lim_{t \rightarrow \infty} v_i(t) = 0, i \neq k, i = 0, 1, 2, \dots, n$.
- (III) if $\hat{v}_k^* < U < v_0^*$, then $\sum_{i=0}^n v_i(t_k) \geq U$ for some $\{t_k\}$, $t_k \rightarrow \infty$ and $\sum_{i=0}^n v_i(t_l) < U$ for some other sequence $\{t_l\}$, $t_l \rightarrow \infty$. This is Case 3 in section 2.1, $\sum_{i=0}^n v_i(t)$ oscillates around U .

Proof.

1. We shall exclude the following two cases:

(i) $\sum_{i=0}^n v_i(t) \geq U$ for all $t > 0$.

From Case 2 in section 2.1, we have $v_k \rightarrow \hat{v}_k^*$, $v_i \rightarrow 0$, where $i \neq k$.

So $\sum_{i=0}^n v_i \rightarrow \hat{v}_k^* < U$ as $t \rightarrow \infty$. By Lemma 2.2 $\sum_{i=1}^n v_i(t) \rightarrow \hat{v}_0^* < v_0^* < U$ a contradiction to $\sum_{i=0}^n v_i(t) \geq U$ for all $t \geq 0$.

(ii) There exists $\hat{t} > 0$, such that

$$\sum_{i=0}^n v_i(\hat{t}) = U, \text{ and } \left. \frac{d}{dt} \left(\sum_{i=0}^n v_i \right) \right|_{t=\hat{t}} > 0 \quad (2.6)$$

In this case, consider the system (2.1),

we obtain that

$$\begin{aligned} \left. \frac{d}{dt} \left(\sum_{i=0}^n v_i \right) \right|_{t=\hat{t}} &= \sum_{i=0}^n (g_i(S) f_i(P) - D) v_i \\ &\leq \sum_{i=0}^n (g_i(S) - D) v_i \\ &= \sum_{i=0}^n (g_i(S^{(0)}) - \sum v_i - D) v_i \\ &= \sum_{i=0}^n (g_i(S^{(0)}) - \sum v_i - g_i(\lambda_i)) v_i \\ \sum_{i=0}^n v_i(\hat{t}) &= U > v_0^* = S^{(0)} - \lambda_0 \end{aligned}$$

It follows that $S^{(0)} - \sum_{i=1}^n v_i(\hat{t}) = S^{(0)} - U < \lambda_0 < \lambda_1 < \dots < \lambda_n$,

$g_i(S^{(0)} - \sum_{i=0}^n v_i(\hat{t})) < g_i(\lambda_i)$ for all $i = 0, 1, \dots, n$.

Hence $\left. \frac{d}{dt} \left(\sum_{i=0}^n v_i \right) \right|_{t=\hat{t}} < 0$.

This is a contradiction to (2.6).

Hence, we exclude (i) and (ii). It follows that $\sum_{i=0}^n v_i < U$ for $t \geq t_0$ for some $t_0 > 0$.

2. We shall exclude the following two cases:

(i) $\sum_{i=0}^n v_i(t) \leq U$ for all $t > 0$.

In this case, we have $v_0 \rightarrow v_0^*$, $v_i \rightarrow 0$, where $i \neq k$.

So $\sum_{i=0}^n v_i \rightarrow v_0^* > U$ as $t \rightarrow \infty$. Since $U < \hat{v}_l^* < \hat{v}_k^* < v_0^*$, we obtain a contradiction to $\sum_{i=0}^n v_i(t) \leq U$ for all $t > 0$

(ii) There exists $\bar{t} > 0$, such that

$$\sum_{i=0}^n v_i(\bar{t}) = U, \text{ and } \left. \frac{d}{dt} \left(\sum_{i=0}^n v_i \right) \right|_{t=\bar{t}} < 0 \quad (2.7)$$

In this case, we consider the system (3),
we obtain that

$$\begin{aligned}
 \frac{d}{dt} \left(\sum_{i=0}^n v_i \right) \Big|_{t=\bar{t}} &= \sum_{i=0}^n (g_i(S) f_i(P) - D) v_i \\
 &\geq \sum_{i=0}^n (g_i(S) f_i(P^{(0)}) - D) v_i \\
 &= \sum_{i=0}^n (g_i(S^{(0)} - \sum v_i) f_i(P^{(0)}) - D) v_i \\
 &= \sum_{i=0}^n (g_i(S^{(0)} - \sum v_i) f_i(P^{(0)}) - g_i(\hat{\lambda}_i) f_i(P^{(0)})) v_i \\
 &= \sum_{i=0}^n (g_i(S^{(0)} - \sum v_i) - g_i(\hat{\lambda}_i)) f_i(P^{(0)}) v_i
 \end{aligned} \tag{2.8}$$

Let

$$\hat{\lambda}_l = \max_{0 \leq i \leq n} \hat{\lambda}_i, \text{ and } \hat{v}_l^* = S^{(0)} - \hat{\lambda}_l < \hat{v}_k^*$$

When $U \leq \hat{v}_l^*$, $\sum_{i=0}^n v_i(\bar{t}) = U \leq \hat{v}_l^* = S^{(0)} - \hat{\lambda}_l$

It follows that $S^{(0)} - \sum_{i=0}^n v_i(\bar{t}) \geq \hat{\lambda}_l > \hat{\lambda}_i$, $i = 0, 1, \dots, n$ and $i \neq l$

The monotonicity of $g_i(S)$ implies that $g_i(S^{(0)} - \sum v_i) \geq g_i(\hat{\lambda}_i)$, from (2.8)

$$\frac{d}{dt} \left(\sum_{i=0}^n v_i \right) \Big|_{t=\bar{t}} \geq 0$$

This is a contradiction to (2.7)

Hence $\sum_{i=0}^n v_i(t) \geq U$ for all t large.

3. Assume $\hat{v}_k^* < U < \hat{v}_0^*$, $\sum_{i=0}^n v_i \geq U$ for all $t > 0$.

Then from (H3) and (2.6), $v_k \rightarrow \hat{v}_k^*$, $v_i \rightarrow 0$, where $i \neq k$.

So $\sum_{i=0}^n v_i \rightarrow \hat{v}_k^* < U$ as $t \rightarrow \infty$, a contradiction to the assumption $\sum_{i=0}^n v_i \geq U$ for all t large.

Suppose $\sum_{i=0}^n v_i < U$ for t large.

Then from (H2) $v_0 \rightarrow v_0^*$, $v_i \rightarrow 0$, $i = 1, 2, \dots, n$, as $t \rightarrow \infty$.

So $\sum_{i=0}^n v_i \rightarrow v_0^* > U$ as $t \rightarrow \infty$, a contradiction.

Hence $\sum_{i=0}^n v_i$ oscillates around U .

□

Remark 2.4. When $\hat{v}_l^* \leq U < \hat{v}_k^*$, from the numerical studies in section 5, we conjecture that the conclusion in (II) in Theorem 2.3 holds, i.e. $\lim_{t \rightarrow \infty} S(t) = \hat{\lambda}_k$, $\lim_{t \rightarrow \infty} v_k(t) = \hat{v}_k^* = S^{(0)} - \hat{\lambda}_k$, $\lim_{t \rightarrow \infty} v_i(t) = 0$ for $i \neq k$, $i = 0, 1, \dots, n$.

Remark 2.5. When the threshold value U satisfies $\hat{v}_k^* < U < \hat{v}_0^*$, we conjecture that the competitive exclusion principle holds. In this case, we conjecture that there exist a bifurcation point U^* such that for $\hat{v}_k^* < U < U^*$, species v_k wins the competition; for $U^* < U < \hat{v}_0^*$ the wild type species $u := v_0$ wins the competition. We note that from (III) of Theorem 2.3 for $\hat{v}_k^* < U < \hat{v}_0^*$, $\sum_{i=0}^n v_i$ oscillates around U . Thus the solution of the system (3) will not tend to equilibrium $E_0 = (\lambda_0, \hat{v}_0^*, 0, \dots, 0)$ or $E_1 = (\hat{\lambda}_0, 0, \dots, 0, \hat{v}_k^*, 0, \dots, 0)$. See Figure 5.3 (a), 5.3 (b), 5.3 (c) and Figure 5.4(a), 5.4(b), 5.4(c) in Section 5.

3. Dynamics of drug on-drug off model with forward mutations

Consider the following drug on-drug off model with forward mutation:

$$\begin{cases} \frac{dS}{dt} = (S^{(0)} - S)D - \sum_{i=0}^n g_i(S)f_i(P)v_i \\ \frac{du}{dt} = (g_0(S)f_0(P) - D)u - q_0u \\ \frac{dv_i}{dt} = (g_i(S)f_i(P) - D)v_i + q_{i-1}v_{i-1} - q_iv_i, i = 1, \dots, n-1 \\ \frac{dv_n}{dt} = (g_n(S)f_n(P) - D)v_n + q_{n-1}v_{n-1} \\ \frac{dP}{dt} = \begin{cases} -DP, & \text{if } \sum_{j=0}^n v_j < U \\ (P^{(0)} - P)D, & \text{if } \sum_{j=0}^n v_j \geq U \end{cases} \end{cases} \quad (1)$$

3.1. The behavior of the solutions of the system (1)

As we did in Section 2, we discuss three cases.

Case 1 : $\sum_{i=0}^n v_i(t) < U$ for all $t > t_0$ for some $t_0 > 0$. Then we have

$$\frac{dP}{dt} = -DP.$$

It is obvious that

$$P(t) \rightarrow 0 \text{ as } t \rightarrow \infty$$

As we did in case 1 in Section 2.1, it suffices to consider the limiting system of (1)

$$\begin{cases} S = S^{(0)} - \sum_{i=0}^n v_i \\ \frac{dv_0}{dt} = (g_0(S) - D - q_0)v_0 \\ \frac{dv_i}{dt} = (g_i(S) - D - q_i)v_i + q_{i-1}v_{i-1}, i = 1, 2, \dots, n-1 \\ \frac{dv_n}{dt} = (g_n(S) - D)v_n + q_{n-1}v_{n-1} \end{cases} \quad (3.1)$$

Theorem 3.1. Let $q = (q_0, q_1, \dots, q_n)$, then (3.1) has a unique positive equilibrium $E(q) = (v_0^*(q), \dots, v_n^*(q))$ satisfying

$$\begin{cases} S^* = g_0^{-1}(D + q_0) \\ (g_i(S^*) - D - q_i)v_i^* + q_{i-1}v_{i-1}^* = 0, \quad i = 1, 2, \dots, n-1 \\ (g_n(S^*) - D)v_n^* + q_{n-1}v_{n-1}^* = 0 \\ \sum_{i=0}^n g_i(S^*)v_i^* = (S^{(0)} - S^*)D \end{cases} \quad (3.2)$$

Proof. Consider the first equation of (3.1) :

$$\frac{dv_0}{dt} = (g_0(S) - D - q_0)v_0.$$

Then the positive equilibrium, $v_0 = v_0^*(q) > 0$, satisfies

$$g_0(S) - D - q_0 = 0.$$

The monotocity of $g_0(S)$ implies that $S^* = g_0^{-1}(D + q_0)$ is unique.

Since q_0 is close to 0, we have S^* is close to $g_0^{-1}(D) = \lambda_0$.

Let $S^* = \lambda_0 + \epsilon$, $\epsilon > 0$ is small enough.

From (H2), it follows that $g_i(S^*) = g_i(\lambda_0 + \epsilon) < g_i(\lambda_i) = D$, $i = 1, 2, \dots, n-1$.

Then we have for $i = 1, 2, \dots, n-1$,

$$g_i(S^*) - D - q_i < 0 \text{ and } g_n(S^*) - D < 0 \quad (3.3)$$

Consider the $n + 1$ equations in (3.2), we denote them in the form of $Ax = y$ as following,

$$\begin{bmatrix} q_0 & g_1(S^*) - D - q_1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & q_1 & g_2(S^*) - D - q_2 & \cdots & 0 & 0 & 0 \\ 0 & 0 & q_2 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & q_{n-1} & g_n(S^*) - D & 0 \\ g_0(S^*) & g_1(S^*) & g_2(S^*) & \cdots & g_{n-1}(S^*) & g_n(S^*) & 0 \end{bmatrix} \begin{bmatrix} v_0 \\ v_1 \\ v_2 \\ \vdots \\ v_{n-1} \\ v_n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ D(S^{(0)} - S^*) \end{bmatrix}$$

Let $A^{(0)} =$

$$\left[\begin{array}{ccccccc|c} q_0 & g_1(S^*) - D - q_1 & 0 & \cdots & 0 & 0 & 0 & 0 \\ 0 & q_1 & g_2(S^*) - D - q_2 & \cdots & 0 & 0 & 0 & 0 \\ 0 & 0 & q_2 & \cdots & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & q_{n-1} & g_n(S^*) - D & 0 & 0 \\ g_0(S^*) & g_1(S^*) & g_2(S^*) & \cdots & g_{n-1}(S^*) & g_n(S^*) & 0 & D(S^{(0)} - S^*) \end{array} \right]$$

We apply the following elementary transformations on $A^{(0)}$.

(Step 1) Consider the $(n + 1)$ -th row in $A^{(0)}$, multiplying $\frac{g_0(S^*)}{q_0}$ to 1st row and subtracting it from $(n + 1)$ -th row, and we have

$A^{(1)} =$

$$\left[\begin{array}{cccccc|c} q_0 & g_1(S^*) - D - q_1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & q_1 & g_2(S^*) - D - q_2 & \cdots & 0 & 0 & 0 \\ 0 & 0 & q_2 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & q_{n-1} & g_n(S^*) - D & 0 \\ 0 & m_1 & g_2(S^*) & \cdots & g_{n-1}(S^*) & g_n(S^*) & D(S^{(0)} - S^*) \end{array} \right]$$

where $m_1 = g_1(S^*) - \frac{g_0(S^*)}{q_0} \cdot (g_1(S^*) - D - q_1)$

Since $g_1(S^*) > 0$, $g_0(S^*) > 0$, $q_0 > 0$, we have $m_1 > 0$.

After n elementary transformations, we get an upper triangular matrix

$A^{(n)} =$

$$\left[\begin{array}{cccccc|c} q_0 & g_1(S^*) - D - q_1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & q_1 & g_2(S^*) - D - q_2 & \cdots & 0 & 0 & 0 \\ 0 & 0 & q_2 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & q_{n-1} & g_n(S^*) - D & 0 \\ 0 & 0 & 0 & \cdots & 0 & m_n & D(S^{(0)} - S^*) \end{array} \right] \quad (3.4)$$

So the solution $(v_0^*, v_1^*, v_2^*, \dots, v_n^*)$ exists and is unique.

By mathematical induction, we can show that, m_n in $A^{(n)}$ is positive.

From (3.4), we have

$$m_n v_n^* = D(S^{(0)} - S^*)$$

and it follows that $v_n^* > 0$

From (3.2), we obtain

$$v_{n-1}^* = \frac{-(g_n(S^*) - D)v_n^*}{q_{n-1}} > 0.$$

For each $1 \leq i \leq n-1$,

$$\frac{v_i^*}{v_{i-1}^*} = \frac{q_{i-1}}{-g_i(S^*) + D + q_i} > 0.$$

Hence $v_i^* > 0$ for all $0 \leq i \leq n$.

and the equilibrium $E(q) = (v_0^*(q), v_1^*(q), v_2^*(q), \dots, v_n^*(q))$ is positive.

□

Since $q \approx 0$, $E(q)$ is closed to E_0 , so we use the following perturbation theory to discuss the local and global stability of $E(q)$.

Theorem 3.2. [4] Let $A, B \in \mathbb{C}^{n \times n}$, with A simple. If A has eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ and μ is an eigenvalue of $A + B$, and if for a matrix norm induced by an absolute vector norm $\|\cdot\|_v$, we have $r = \|B\| \|A\|_v$, then μ lies in at least one of the disks $z : |z - \lambda_j| \leq r$, $j = 1, 2, \dots, n$, of the complex z -plane.

Theorem 3.3. [7] Assume that $(x_0, \lambda_0) \in U \times \Lambda$, $x_0 \in \text{Int}U$, $f(x_0, \lambda_0) = 0$, all eigenvalues of $D_x f(x_0, \lambda_0)$ have negative real part, and x_0 is globally attracting for solutions of $x' = f(x, \lambda)$ with $\lambda = \lambda_0$. If there exists a compact set $D \subset U$ such that for each $\lambda \in \Lambda$ and each $z \in U$, $x(t, z, \lambda) \in D$ for all large t , then there exists $\epsilon > 0$, and a unique point $\hat{x}(\lambda) \in U$ for $\lambda \in B_\Lambda(\lambda_0, \epsilon)$ such that $f(\hat{x}(\lambda), \lambda) = 0$ and $x(t, z, \lambda) \rightarrow \hat{x}(\lambda)$ as $t \rightarrow \infty$ for all $z \in U$.

Theorem 3.4. $E(q)$ is locally stable.

Proof. From (2.2) in Section 2, E_0 is a stable equilibrium where $E_0 = (v_0^*, 0, \dots, 0)$, $v_0^* = S^{(0)} - \lambda_0$. The Jacobi matrix of system (2.2) evaluated at E_0 , is

$$J(E_0) = \begin{bmatrix} -v_0^* g_0'(\lambda_0) & -v_0^* g_0'(\lambda_0) & \cdots & \cdots & -v_0^* g_0'(\lambda_0) \\ 0 & g_1(\lambda_0) - D & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & g_n(\lambda_0) - D \end{bmatrix} \quad (3.5)$$

The eigenvalues $\mu_0, \mu_2, \dots, \mu_{n+1}$ of $J(E_0)$ are $\mu_0 = -v_0^* g_0'(\lambda_0) < 0, \mu_1 = g_1(\lambda_0) - D < 0, \dots, \mu_n = g_n(\lambda_0) - D < 0$.

From (3.2), we have the Jacobi matrix of system (3.1) evaluated at $E(q)$,

$$J(E(q)) = \begin{bmatrix} -g_0'(S^*)v_0^* & \cdots & \cdots & \cdots & -g_0'(S^*)v_0^* \\ -g_1'(S^*)v_1^* + q_0 & (g_1(S^*) - D - q_1) & \cdots & \cdots & -g_1'(S^*)v_1^* \\ -g_2'(S^*)v_2^* & -g_2'(S^*)v_2^* + q_1 & (g_2(S^*) - D - q_2) & \cdots & -g_2'(S^*)v_2^* \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ -g_n'(S^*)v_n^* & -g_n'(S^*)v_n^* & \cdots & -g_n'(S^*) + q_{n-1} & (g_n(S^*) - D) \\ & & & & + (-g_n'(S^*))v_n^* \end{bmatrix} \quad (3.6)$$

Since q_i is sufficiently small,

we have $g_i(S^*) \approx g_i(\lambda_0)$ and $g_i'(S^*) \approx g_i'(\lambda_0)$.

Let $J(E(q)) = J(E_0) + B$, then $|B| \approx 0$.

From Theorem 3.2, the eigenvalues of $J(E(q))$ lies in at least one of the disks $z : |z - \mu_i| \leq \epsilon, i = 0, 1, \dots, n$ (ϵ is small enough).

Since $\mu_i < 0$ for all $i = 0, 1, \dots, n$, all eigenvalues of $J(E(q))$ have negative real part.

Hence $E(q)$ is locally stable. \square

Theorem 3.5. $E(q)$ is globally stable.

Proof. Write (3.2) as $\frac{dx}{dt} = f(x, q)$, when $x = (S, u, v_1, \dots, v_n)$ and $q = (q_0, q_1, \dots, q_{n-1})$.

Take $U = \mathbb{R}_+ \cup B_\epsilon(E_0)$, and $\Lambda(q) = [0, \delta]^{n-1}$.

when $q = 0$, i.e. all the $q_i = 0$ for each $0 \leq i \leq n$, $f(E_0, 0) = 0$, and $D_x f(E_0, 0)$ has all negative eigenvalues.

Take $D = [0, S^{(0)}]^{n+2}$, then D is compact.

when $q_i \approx 0$, and $x = E(q)$, we have $f(E(q), q) = 0$.

By Theorem 3.3, $x(t, z, q) \rightarrow E(q)$ as $t \rightarrow \infty$ for all $z \in U$.

This concludes that $E(q)$ is globally stable. \square

Next we discuss Case 2.

Case 2 : $\sum_{i=0}^n v_i(t) \geq U$ for all $t > t_0$ for some $t_0 > 0$.

Then

$$\frac{dP}{dt} = D(P^{(0)} - P),$$

and hence

$$P(t) \longrightarrow P^{(0)} \text{ as } t \rightarrow \infty.$$

As we did in Case 2 in Section 2.1, it suffice to consider the limiting system of (1)

$$\begin{cases} S = S^{(0)} - \sum_{i=0}^n v_i \\ \frac{dv_0}{dt} = (g_0(S)f_0(P^{(0)}) - D - q_0)v_0 \\ \frac{dv_i}{dt} = (g_i(S)f_i(P^{(0)}) - D - q_i)v_i + q_{i-1}v_{i-1} \\ \frac{dv_n}{dt} = (g_n(S)f_n(P^{(0)}) - D)v_n + q_{n-1}v_{n-1} \end{cases} \quad (3.7)$$

We shall study (3.7) by using the same method for the system (2.4) in section 2.1, Suppose (H3) holds, let

$$\hat{\lambda}_i = \frac{a_i}{\frac{m_i f_i(P^{(0)})}{D} - 1} > 0, \quad i = 0, 1, 2, \dots, n.$$

Theorem 3.6.

1. System (3.7) has an equilibrium $E_* = (v_{0*}, \dots, v_{n*})$.

(i) If $\hat{\lambda}_0 = \min_{0 \leq i \leq n} \hat{\lambda}_i$, then $v_{i*} > 0$ for all $i = 0, 1, \dots, n$. Write $E_* = (\bar{v}_0, \bar{v}_1, \dots, \bar{v}_n)$ satisfying

$$\begin{cases} \bar{S} = g_0^{-1} \left(\frac{D + q_0}{f_0(P^{(0)})} \right) \\ (g_i(\bar{S})f_i(P^{(0)}) - D - q_i)\bar{v}_i + q_{i-1}\bar{v}_{i-1} = 0, \quad i = 1, \dots, n-1 \\ (g_n(\bar{S})f_n(P^{(0)}) - D)\bar{v}_n + q_{n-1}\bar{v}_{n-1} = 0 \\ \sum_{i=0}^n g_i(\bar{S})f_i(P^{(0)})\bar{v}_i = (S^{(0)} - \bar{S})D \end{cases} \quad (3.8)$$

(ii) If $\hat{\lambda}_k = \min_{0 \leq i \leq n} \hat{\lambda}_i$ for some $k, 0 < k < n$, then $v_{i*} = 0$ for $i = 0, 1, \dots, k-1$ and $v_{j*} > 0$ for $k \leq j \leq n$. Write $E_* = (0, \dots, \bar{v}_k, \dots, \bar{v}_n)$ satisfying

$$\begin{cases} \bar{S} = g_k^{-1} \left(\frac{D + q_k}{f_k(P^{(0)})} \right) \\ \bar{v}_i = 0 \quad (0 \leq i < k) \\ (g_i(\bar{S})f_i(P^{(0)}) - D - q_i)\bar{v}_i + q_{i-1}\bar{v}_{i-1} = 0 \quad (k \leq i \leq n-1) \\ (g_n(\bar{S})f_n(P^{(0)}) - D)\bar{v}_n + q_{n-1}\bar{v}_{n-1} = 0 \\ \sum_{i=0}^n g_i(\bar{S})f_i(P^{(0)})\bar{v}_i = (S^{(0)} - \bar{S})D \end{cases} \quad (3.9)$$

(iii) If $\hat{\lambda}_n = \min_{0 \leq i \leq n} \hat{\lambda}_i$, then $v_{i^*} = 0$ for all $i = 0, 1, \dots, n - 1$ and $v_{n^*} > 0$. Write $E_* = (0, 0, \dots, 0, v_{n^*})$, $v_{n^*} = S^{(0)} - \hat{\lambda}_n$.

2. For the system (3.7), E_* is global stable.

Proof.

1.

(i) Since $\hat{\lambda}_0 < \hat{\lambda}_i, i = 1, 2, \dots, n$, replacing $g_i(S)$ by $f_i(P^{(0)})g_i(S)$ in Theorem 3.1, we complete the proof of (i).

(ii) Consider the steady state of the first equation in (3.7),

$$(g_0(S^*)f_0(P^{(0)}) - D - q_0)v_0^* = 0.$$

Claim: $v_0^* = 0$

if not, $v_0^* > 0$ then $g_0(S^*)f_0(P^{(0)}) - D - q_0 = 0$. Since the mutation rate $q_i, i = 0, 1, \dots, n$ are sufficiently small, $g_0(S^*) \approx g_0(\hat{\lambda}_k)$.

$$\begin{aligned} g_0(S^*)f_0(P^{(0)}) - D - q_0 &\approx g_0(\hat{\lambda}_k)f_0(P^{(0)}) - D - q_0 \\ &< g_0(\hat{\lambda}_0)f_0(P^{(0)}) - D - q_0 = -q_0 < 0. \end{aligned}$$

This leads to a contradiction $g_0(S^*)f_0(P^{(0)}) - D - q_0 = 0$. Hence we have $v_0^* = 0$.

Similarly from the steady state equation of (3.7)

$$(g_i(S^*)f_i(P^{(0)}) - D - q_i)v_{i^*} + q_{i-1}v_{(i-1)^*} = 0, i = 1, 2, \dots, k - 1,$$

we obtain $v_{i^*} = 0, i = 1, 2, \dots, k - 1$. Thus $E_* = (0, \dots, 0, \hat{v}_k, \dots, \hat{v}_n)$ satisfies (3.9).

Consider the $n - k + 1$ equations in (3.9), we denote them in the form of $Ax = y$ as following,

$$\begin{aligned} &\begin{bmatrix} q_k & g_{k+1}(\tilde{S})f_{k+1}(P^{(0)}) & \dots & 0 & 0 \\ 0 & q_{k+1} & g_{k+1}(\tilde{S})f_{k+2}(P^{(0)}) & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & q_{n-1} & g_n(\tilde{S})f_n(P^{(0)}) - D \\ g_k(\tilde{S})f_k(P^{(0)}) & g_{k+1}(\tilde{S})f_{k+1}(P^{(0)}) & \dots & g_{n-1}(\tilde{S})f_{n-1}(P^{(0)}) & g_n(\tilde{S})f_n(P^{(0)}) \end{bmatrix} \begin{bmatrix} \tilde{v}_k \\ \tilde{v}_{k+1} \\ \vdots \\ \tilde{v}_{n-1} \\ \tilde{v}_n \end{bmatrix} \\ &= \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ D(S^{(0)} - \tilde{S}) \end{bmatrix} \end{aligned} \tag{3.10}$$

Since q_k is close to 0, we have \tilde{S} is close to $g_k^{-1}(\frac{D}{f_k(P^{(0)})}) = \hat{\lambda}_k$.

Let $\tilde{S} = \hat{\lambda}_k + \epsilon$, ϵ is small enough,

From (H2), get $g_i(\tilde{S})f_i(P^{(0)}) = g_i(\hat{\lambda}_k + \epsilon)f_i(P^{(0)}) < g_i(\hat{\lambda}_i)f_i(P^{(0)}) = D$.

It follows that $g_i(\tilde{S})f_i(P^{(0)}) - D - q_i < 0$, and $g_n(\tilde{S}) - D < 0$.

Use the same method as Theorem 3.1, we have $\tilde{v}_i > 0$ when $k \leq i \leq n$.

(iii) Particularly, when $\hat{\lambda}_n = \min_{0 \leq i \leq n} \hat{\lambda}_i$,

Let $g_n(S)f_n(P^{(0)}) - D = 0$, we can get $S = \hat{\lambda}_n$.

Similarly, we can use mathematical induction as the second part of Theorem 3.6 to prove that $v_{i^*} = 0$ for all $0 \leq i \leq n - 1$, and then $v_{n^*} = S^{(0)} - \lambda_n$.

2. Since q_i is small enough and close to 0, E_* is close to E_1 .

We can also use the perturbation theory (Theorem 3.2 and Theorem 3.3) to prove the local and global stability of E_* . The proof is similar to Theorem 3.4 and Theorem 3.5.

□

Case 3 :

If $\sum_{i=0}^n v_i(t_k) < U$ at some sequence $\{t_k\}$, $t_k \rightarrow \infty$ and $\sum_{i=0}^n v_i(t_l) \geq U$ at some other sequence $\{t_l\}$, $t_l \rightarrow \infty$, $k = 1, 2, \dots, l = 1, 2, \dots, k \neq l$, $\sum_{i=0}^n v_i$ oscillates around U .

3.2. The results of the asymptotic behavior of the solutions of model with forward mutations

From Lemma 2.1, we have $S(t) + \sum_{i=0}^n v_i(t) \rightarrow S^{(0)}$ as $t \rightarrow \infty$.

In case 1, $S(t) \rightarrow S^* = g_0^{-1}(D + q_0)$, which is close to λ_0 .

So $\sum_{i=0}^n v_i(t)$ is close to $S^{(0)} - \lambda_0 (= v_0^*)$ as $t \rightarrow \infty$.

In case 2, $S \rightarrow S_* = g_k^{-1}(D + q_k)$, which is approximately equal to $\hat{\lambda}_k$, but a little greater than $\hat{\lambda}_k$.

So $\sum_{i=0}^n v_i(t)$ is close to $S^{(0)} - \hat{\lambda}_k (= \hat{v}_k^*)$ as $t \rightarrow \infty$.

By the theory of perturbation, we can get the same relation between $\sum_{i=0}^n v_i$ and U as Theorem 2.3 in Section 2.

Theorem 3.7. Let (H1), (H2), (H3) hold, then

- (I) if $U > v_0^*$, then there exists t_0 , such that for all $t \geq t_0$, $\sum_{i=0}^n v_i(t) < U$
- (II) Let $\hat{\lambda}_l = \max_{0 \leq i \leq n} \hat{\lambda}_i$ and $\hat{v}_l^* = S^{(0)} - \hat{\lambda}_l < \hat{v}_k^*$. If $U < \hat{v}_l^*$, then there exists $t_0 > 0$, such that for all $t \geq t_0$, $\sum_{i=0}^n v_i(t) \geq U$.
- (III) if $\hat{v}_k^* < U < v_0^*$, then $\sum_{i=0}^n v_i(t_k) \geq U$ for some sequence $\{t_k\}$, $t_k \rightarrow \infty$ and $\sum_{i=0}^n v_i(t_l) < U$ for some other sequence $\{t_l\}$, $t_l \rightarrow \infty$.

Proof.

(I) Assume $U > v_0^*$, we shall exclude the following two cases.

- (i) $\sum_{i=0}^n v_i(t) \geq U$ for $t > 0$. Then $P(t) \rightarrow P^{(0)}$ as $t \rightarrow \infty$. Consider the limiting system (3.7). By perturbation method in Theorem 3.3 and the result in Case 2 of section 2.1, we have

$$v_k(t) \rightarrow v_k^* + c_1(q), v_i(t) \rightarrow c_i(q)$$

where $i \neq k$, $c_1(q)$ and $c_i(q)$ are small, $q = (q_1, \dots, q_n)$. Then

$$\sum_{i=0}^n v_i \rightarrow v_k^* + c_1(q) + \sum_{i \neq 1} c_i(q)$$

and it follows that

$$\sum_{i=0}^n v_i(t) < v_k^* + c_1(q) + \sum_{i \neq 1} c_i(q) < v_0^* < U, \text{ for } t \text{ large.}$$

This is a contradiction to the assumption (i):

$$\sum_{i=0}^n v_i(t) \geq U \text{ for } t \geq 0.$$

(ii) There exists $\hat{t} > 0$ such that

$$\sum_{i=0}^n v_i(\hat{t}) = U \text{ and } \left. \frac{d}{dt} \left(\sum_{i=0}^n v_i(t) \right) \right|_{t=\hat{t}} > 0.$$

From the system (1),

$$\begin{aligned} \left. \frac{d}{dt} \left(\sum v_i \right) \right|_{t=\hat{t}} &= \sum_{i=0}^n (g_i(S) f_i(P) - D) v_i \\ &\leq \sum (g_i(S) - D) v_i \\ &= \sum (g_i(S^{(0)} - \sum v_i) - D) \\ &= \sum (g_i(S^{(0)} - \sum v_i) - g_i(\lambda_i)), \end{aligned}$$

$$\sum_{i=0}^n v_i(\hat{t}) = U > v_0^* = S^{(0)} - \lambda_0.$$

It follows that

$$\begin{aligned} S^{(0)} - \sum v_i(\hat{t}) &< \lambda_0 < \lambda_1 < \dots < \lambda_n \\ g_i \left(S^{(0)} - \sum_{i=0}^n v_i(\hat{t}) \right) &< g_i(\lambda_i), i = 0, 1, \dots, n. \end{aligned}$$

Hence

$$\left. \frac{d}{dt} \left(\sum_{i=0}^n v_i \right) \right|_{t=\hat{t}} < 0.$$

This is a contradiction. Excluding (i) and (ii), we obtain $\sum_{i=0}^n v_i < U$ for t large.

- (II) Assume $U < \hat{v}_l^*$. Then $\sum_{i=0}^n v_i(t) \geq U$ for $t \geq t_0$. Applying the perturbation method and the method in Theorem 2.3 (II), we finish the proof of (II).
- (III) Assume $v_k^* < U < v_0^*$. Suppose $\sum_{i=0}^n v_i \geq U$ for t large. Then by perturbation method in Theorem 3.3 and the results in Case 2 of section 2.1,

$$v_k(t) \rightarrow v_k^* + c_1(q), v_i(t) \rightarrow c_i(q), i \neq k,$$

$$\sum v_i \rightarrow v_k^* + c_1(q) + \sum_{i \neq k} c_i(q) < U,$$

a contradiction.

Suppose $\sum_{i=0}^n v_i < U$ for t large. Then

$$v_0(t) \rightarrow v_0^* + c_0(q), v_i(t) \rightarrow c_i(q), i = 1, 2, \dots, n \text{ as } t \rightarrow \infty.$$

Hence $\sum v_i \rightarrow v_0^* + c_0(q) + \sum_{i=1}^n c_i(q) > U$, a contradiction.

Hence $\sum_{i=0}^n v_i$ oscillates around U .

□

4. Dynamics of drug on-drug off model with forward-backward mutations

Consider the model

$$\begin{cases} \frac{dS}{dt} = (S^{(0)} - S)D - \sum_{i=0}^n g_i(S)f_i(P)v_i \\ \frac{du}{dt} = (g_0(S)f_0(P) - D)u - q_0u + \widetilde{q}_0v_1 \\ \frac{dv_i}{dt} = (g_i(S)f_i(P) - D)v_i + q_{i-1}v_{i-1} - q_iv_i - \widetilde{q}_{i-1}v_i + \widetilde{q}_iv_{i+1}, 1 \leq i \leq n-1 \\ \frac{dv_n}{dt} = (g_n(S)f_n(P) - D)v_n + q_{n-1}v_{n-1} - \widetilde{q}_{n-1}v_n \\ \frac{dP}{dt} = \begin{cases} -DP, & \text{if } \sum_{j=0}^n v_j < U \\ (P^{(0)} - P)D, & \text{if } \sum_{j=0}^n v_j \geq U \end{cases} \end{cases} \quad (2)$$

Lemma 4.1. ([3], p. 141) If $y(t)$ has a finite limit as $t \rightarrow \infty$ and $y^{(n)}$ is bounded for $t \geq t_0$, then $y^{(k)}(t) \rightarrow 0$ as $t \rightarrow \infty$ for $0 < k < n$.

In this section we only consider Case 1 and Case 2, i.e. we assume either $U > v_0^*$ or $U < \widehat{v}_1^*$. Then the solutions of (2) are smooth.

Lemma 4.2. In (2), if $v_m(t) \rightarrow 0$ as $t \rightarrow \infty$ for some m , $0 \leq m \leq n$, then we can get $v_k(t) \rightarrow 0$ as $t \rightarrow \infty$, for all $k \neq m$ and $0 \leq k \leq n$. That means all the species will go extinction as long as one of them goes extinction when time is long enough. Otherwise, all species coexists.

Proof. We just prove the cases when $0 < m < n$, since $m = 0$ and $m = n$ are similar.

Consider the model (2), we have

$$\begin{aligned} \frac{dv_m}{dt} &= (g_m(S)f_m(P) - D - q_m - \widetilde{q}_{m-1})v_m + q_{m-1}v_{m-1} + \widetilde{q}_mv_{m+1}. \\ \frac{d^2v_m}{dt^2} &= (g_m(S)f_m(P) - D - q_m - \widetilde{q}_{m-1})v_m' + (g_m'(S)S'f_m(P) + g_m(S)f_m'(P)P')v_m \\ &\quad + q_{m-1}v_{m-1}' + \widetilde{q}_mv_{m+1}' \end{aligned} \quad (4.1)$$

From Lemma 2.1, we have

$$S(t) + \sum_{i=0}^n v_i(t) \rightarrow S^{(0)} \text{ as } t \rightarrow \infty, \quad (4.2)$$

and $0 < S \leq S^{(0)}$ and $0 < P \leq P^{(0)}$.

From the first equation in (2),

$$\begin{aligned} |S'| &\leq S^{(0)}D + \max_{0 \leq j \leq n} \{g_j(S^{(0)})\} \cdot \sum_{i=1}^n v_i \\ &\leq S^{(0)}D + M_s S^{(0)}. \end{aligned}$$

For $1 \leq i \leq n-1$,

$$\begin{aligned} |v'_i| &\leq |(g_i(S)f_i(D) - D - q_i - \widetilde{q}_{i-1})v_i| + q_{i-1}v_{i-1} + \widetilde{q}_i v_{i+1} \\ &\leq \max \left\{ g_i(S^{(0)}) + D + g_i + \widetilde{g}_{i-1}, q_{i-1}, \widetilde{g}_i \right\} \sum_{i=0}^n v_i, \end{aligned}$$

from (4.2),

$$\sum_{i=0}^n v_i \leq S^{(0)}, 0 < S < S^{(0)}, 0 < P < P^{(0)},$$

it is easy to see that $|v'_i|$, $|P'| \leq DP^{(0)}$, $|S'|$, $|f'_m(P)|$, $|g'_m(S)|$ are bounded.

It follows from (4.1), we conclude that $|\frac{d^2 v_m}{dt^2}|$ is bounded.

Since $v_m \rightarrow 0$ as $t \rightarrow \infty$, and $|\frac{d^2 v_m}{dt^2}|$ is bounded.

From Lemma 4.1, we have $\frac{dv_m}{dt} \rightarrow 0$ as $t \rightarrow \infty$.

It implies that $q_{m-1}v_{m-1} + \widetilde{q}_m v_{m+1} \rightarrow 0$ as $t \rightarrow \infty$, $v_{m-1} \rightarrow 0$, $v_{m+1} \rightarrow 0$ as $t \rightarrow \infty$.

Hence $v_k \rightarrow 0$ as $t \rightarrow \infty$ for $k \neq m$. □

From Lemma 4.2, there are two cases: Either all the bacteria v_i ($0 \leq i \leq n$) go extinction or all of them coexist. In the drug on-drug off model with forward-backward mutations, we get the next results.

Theorem 4.3. Let the hypothesis (H1) and (H2) hold and v_i ($0 \leq i \leq n$) be the solutions of system (2), then the species v_i ($0 \leq i \leq n$) coexists.

Proof. If all the species v_i ($i = 0, 1, \dots, n$) go extinction, then $\sum_{i=0}^n v_i \rightarrow 0 < U$ as $t \rightarrow \infty$.

We have

$$S \rightarrow S^{(0)} \text{ and } P \rightarrow 0.$$

Consider the second equation of the system (2):

$$\frac{du}{dt} = (g_0(S^{(0)}) - D - q_0)u + \widetilde{q}_0 v_1$$

Since $g_0(S^{(0)}) - D = g_0(S^{(0)}) - g_0(\lambda_0) > 0$ and q_i, \widetilde{q}_i are sufficiently small.

$\frac{du}{dt} > 0$ as $t \rightarrow \infty$, which contradicts to $u \rightarrow 0$ as $t \rightarrow \infty$.

Hence all the species v_i ($i = 0, 1, \dots, n$) coexist for t large in the system (2) with forward-backward mutation.

□

Remark 4.4. We note that since the mutant rates \tilde{q}_i are quite small, the most resistant v_k whose $\hat{\lambda}_k = \min_{0 \leq i \leq n} \hat{\lambda}_i$, where $\hat{\lambda}_i = \frac{a_i}{\frac{m_i f_i(P)}{D} - 1}$ dominates the rest species.

5. Numerical simulations

In this numerical simulation, we consider the case $n = 2$ and verify the conjecture in Remark 2.4 and Remark 2.5.

For simplicity, we assume $S^{(0)} = 10$, $D = 0.9$ and

$$g_i(S) = \frac{m_i S}{a_i + S} \text{ for } i = 0, 1, \dots, n$$

where $a_0 = 2$, $a_1 = 2$, $a_2 = 3$, $m_0 = 3$, $m_1 = 2$, $m_2 = 1.5$.

Since $g_i(\lambda_i) = D$, we have $\lambda_0 \approx 0.86$, $\lambda_1 \approx 2.44$ and $\lambda_2 = 4.5$.

Hence $v_0^* = S^{(0)} - \lambda_0 \approx 9.14$.

Let $P^{(0)} = 10$ and

$$f_i(P) = \frac{1}{1 + (\frac{P}{K_i})^L} \text{ for } i = 0, 1, 2$$

where $L = 1$, $K_0 = 6$, $K_1 = 15$, $K_2 = 40$.

After calculations, we have $\hat{\lambda}_0 = 8$, $\hat{\lambda}_1 = 6$, $\hat{\lambda}_2 = 9$, Then $\hat{\lambda}_1 = \min(\hat{\lambda}_0, \hat{\lambda}_1, \hat{\lambda}_2)$, $\hat{\lambda}_2 = \max(\hat{\lambda}_0, \hat{\lambda}_1, \hat{\lambda}_2)$, and

$\hat{v}_k^* = \hat{v}_1^* = 4$, and $\hat{v}_l^* = \hat{v}_2^* = 1$.

Take $U = 3$, then $\hat{v}_2^* \leq U < \hat{v}_1^*$.

1. When there is no mutation, under the assumption (H3), (2.5), the mutant v_2 survives, but the wide type v_0 and the other mutant v_1 go extinct. $V := \sum_{i=0}^2 v_i(t) \geq U$ for time t large, satisfying the conjecture in Remark 2.4.

In Figure 5.1 (a), (b), (c) we verify the conjecture that if $\hat{v}_l^* < U < \hat{v}_k^*$ then $v_k(t) \rightarrow \hat{v}_k^*$, $v_i \rightarrow 0$ for $i \neq k$.




Figure 5.1 (a). Shows the prediction $V := \sum_{i=0}^n v_i > U$ for t large. The numerical data is $U = 3, \hat{v}_l^* = 1 < U = 3 < \hat{v}_k^* = 4$.




Figure 5.1 (b). Shows the prediction $P(t) \rightarrow P^{(0)}$ for t large, $P^{(0)} = 10$.




Figure 5.1 (c). Shows $v_k(t) \rightarrow \hat{v}_k^*$ as $t \rightarrow \infty$ and $v_i(t) \rightarrow 0, i \neq k$ as $t \rightarrow \infty$, where $k = 1, v_k^* = v_1^* = 4$.

In the following Figure 5.2 and Figure 5.3, U satisfies $\hat{v}_k^* < U < v_0^*$. Thus the total population $V = \sum_{i=0}^n v_i$ oscillates around U . We conjecture that there exists U^* between \hat{v}_k^* and v_0^* such that

species v_k win the competition when $\hat{v}_k^* < U < U^*$ and species v_0 (wild Type) win the competition when $U^* < U < v_0^*$. For the numerical data is $n = 2$, $\hat{v}_k^* = 4$, $v_0^* = 9.14$, we compute the value U^* , $U^* \approx 5.846$.

2. Let $U = 5.5$, $\hat{v}_k^* < U < U^*$, $\hat{v}_k^* = 4$, $U^* \approx 5.846$ and the initial data $(9.2, 0.3, 0.3, 0.2)$ near equilibrium $E_0 = (v_0^*, 0, 0, 0) = (9.14, 0, 0, 0)$, then the trajectories of the system (3) are shown in Figure 5.2 (a), 5.2 (b), 5.2 (c).




Figure 5.2 (a). $V = \sum_{i=0}^2 v_i$ is the total population of the bacteria. Since $\hat{v}_k^* = 4 < U = 5.5 < v_0^* = 9.14$, from (III) of Theorem 2.3, V oscillates around U .




Figure 5.2 (b). The drug concentration $P(t)$ oscillates between 0 and $P^{(0)} = 10$.




Figure 5.2 (c). The population v_1 wins the competition however v_1 oscillates, not tends to v_1^* .

- Let $U = 6.5$, $U^* < U < v_0^*$, $U^* \approx 5.846$, $v_0^* = 9.14$, and the initial data $(6.2, 0.2, 4.3, 0.3, 9.5)$ near equilibrium $E_1 = (0, v_1^*, 0, P^{(0)}) = (0, 4, 0, 10)$, then the trajectories of the system (3) are shown in Figure 5.3 (a), 5.3 (b), and 5.3 (c).




Figure 5.3 (a). The total population V oscillates around $U = 6.5$.




Figure 5.3 (b). The drug concentration $P(t)$ oscillates between 0 and $P^{(0)} = 10$.




Figure 5.3 (c). The population v_0 wins the competition however v_0 oscillates, not tends to v_0^* , $v_0^* = 9.14$.

Acknowledgment

The authors would like to give their sincere thanks to the anonymous referee for his/her valuable suggestions to improvement of the paper. The second author's research is supported by MOST 111-2115-M-007-011.

Conflict of interest

There is no conflict of interest.

References

1. H. L. Smith, P. E. Waltman, *The theory of the chemostat*, Cambridge University Press, Cambridge, 1995. <https://doi.org/10.1017/CBO9780511530043>
2. S. B. Hsu, K. C. Chen, *Ordinary differential equations with applications*, Series on Applied Mathematics, Vol 23, World Scientific Press, 2022, 3rd Edition. <https://doi.org/10.1142/8744>
3. W. A. Coppel, *Stability and asymptotic behavior of differential equations*, Health. Math. Monograph, 1965.
4. P. Lancaster, M. Tismenetsky, *The Theory of Matrices-second edition: With Applications*, Academic Press, 1985.
5. Z. Chen, S. B. Hsu, Y. T. Yang, The continuous Morbidostat : a chemostat controlled drug application to select for drug resistance mutants, *Commun. Pur. Appl. Anal.*, **19** 2020, 203–220. <https://doi.org/10.3934/cpaa.2020011>
6. Z. Chen, S. B. Hsu, Y. T. Yang, The Morbidostat: A Bio-reactor That Promotes Selection for Drug Resistance in Bacteria, *SIAM J. Appl. Math.*, **77** (2017), 470–499. <https://doi.org/10.1137/16M105695X>

7. H. L. Smith, P. Waltman, Perturbation of a Globally Stable Steady State, *Proceedings of the American Mathematical Society*, **127** (1999), 447–453. <https://doi.org/10.1090/S0002-9939-99-04768-1>
8. M. Barber, Infection by penicillin resistant Staphylococci, *Lancet*, **2** (1948), 641–644. [https://doi.org/10.1016/S0140-6736\(48\)92166-7](https://doi.org/10.1016/S0140-6736(48)92166-7)
9. S. B. Levy, B. Marshall, Antibacterial resistance worldwide: causes, challenges and responses, *Nat. Med.*, **10** (2004), s122–s129. <https://doi.org/10.1038/nm1145>
10. E. Toprak, A. Veres, S. Yildiz, J. M. Pedraza, R. Chait, J. Paulsson, et al., Building a morbidostat: an automated continuous-culture device for studying bacterial drug resistance under dynamically sustained drug inhibition, *Nat. protoc.*, **8** (2013), 555–567. <https://doi.org/10.1038/nprot.2013.021>
11. E. Toprak, A. Veres, J. B. Michel, R. Chait, D. L. Hartl, R. Kishony, Evolutionary paths to antibiotic resistance under dynamically sustained drug selection, *Nat. Genet.*, **44** (2012), 101–105.



AIMS Press

© 2023 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)