

SMALL POPULATIONS CORRECTIONS FOR SELECTION-MUTATION MODELS

PIERRE-EMMANUEL JABIN

CSCAMM and Department of Mathematics
University of Maryland
College Park, MD 20742-4015, USA

ABSTRACT. We consider integro-differential models describing the evolution of a population structured by a quantitative trait. Individuals interact competitively, creating a strong selection pressure on the population. On the other hand, mutations are assumed to be small. Following the formalism of [20], this creates concentration phenomena, typically consisting in a sum of Dirac masses slowly evolving in time. We propose a modification to those classical models that takes the effect of small populations into accounts and corrects some abnormal behaviours.

1. Introduction.

1.1. The first model. We study the dynamics of a population subject to mutations and selection due to competition between individuals. Each individual in the population is characterized by a quantitative phenotypic trait x (for example the size of the individual, their age at maturity, or their rate of intake of nutrients). For simplicity, x is taken here in \mathbb{R} even though all the arguments could easily be extended to higher dimensional cases.

Probabilistic models are usually considered as the most realistic in that setting. They consist in birth and death processes for each individual X_i , Poisson processes more precisely, with for instance birth rate $b(X_i)$ and a death rate which increases with the competition between individuals, for example

$$d_i = d(X_i) + \sum_{j \neq i} I(X_i - X_j),$$

where I denotes the interaction kernel, *i.e.* the effect that an individual with trait X_j has on the death rate of an individual with trait X_i .

When a birth occurs, it simply adds another individual with the same trait, except when a mutation takes place with small probability. In that case the new individual has a different random trait, obtained through some distribution K . We refer to [28] and the references therein for a nice introduction to the probabilistic approach.

Of course this is only one possible model and there are many variants. One can modify the interaction between individuals for instance by introducing explicit

2000 *Mathematics Subject Classification.* Primary: 35B25, 35K55, 92D15.

Key words and phrases. Adaptive dynamics, Hamilton-Jacobi equation with constraints, Dirac concentration, small populations.

resources (with chemostat like interactions maybe). The competition could influence both the birth rate and the mortality rate...

When the total number of individuals is too large (it can easily reach $10^{10} - 10^{12}$ for some micro-organisms), it becomes prohibitive to compute numerically the solution to this process. In that case one expects to be able to derive a deterministic model as a limit of large populations. Such a derivation was proved in [10] and one obtains integro-differential equations like

$$\partial_t u(t, x) = \left(b(x) - d(x) - \int I(x - y) u(t, y) dy \right) u(t, x) + M(u)(t, x), \quad (1)$$

where the mutation kernel is for instance

$$M(u)(x) = \int_{\mathbb{R}} K(z) (b(x + z) u(x + z) - b(x) u(x)) dz.$$

1.2. The scaling of fast reaction and small mutations. Eq. (1) is easy and fast to solve. However in most situations, small parameters appear and complicate that. Three scales are naturally important for the model

- The reproduction rate, which is also the inverse of the typical time scale to observe one birth or one death.
- The rate of mutations or the inverse of the time scale to see one mutation.
- The size of one mutation when it occurs, *i.e.* the typical size of the change in trait due to one mutation.

The two small scales that are typical for this problem are:

- The size of each mutation is small.
- The change in traits due to mutations, which is the product of the rate of mutations by the size of each mutation, is very small in comparison to the reproduction rate. As we wish to see the evolution of traits generated by the mutations, this means that we need to rescale the equation in time and will hence get a large reproduction rate.

To simplify again our equations, let us now assume that $b = 1$ and denote $r(x) = 1 - d(x)$ the reproduction rate of an individual without competition.

Taking the two scalings into account, one has in fact to deal with an equation like

$$\partial_t u_\varepsilon(t, x) = \frac{1}{\varepsilon} (r(x) - I \star u_\varepsilon) u_\varepsilon(t, x) + M_\varepsilon(u_\varepsilon)(t, x), \quad (2)$$

where the mutation kernel now reads

$$M_\varepsilon(u)(x) = \frac{1}{\varepsilon} \int_{\mathbb{R}} K(z) (b(x + \varepsilon z) u(x + \varepsilon z) - b(x) u(x)) dz.$$

Note that $M_\varepsilon(f)$ is indeed of order 1 if $bu \in C^1$ for instance.

Eq. (2) is now much more delicate. The properties of the solution might depend on ε (its smoothness for instance) and solving it numerically can become again very costly if ε is too small (a typical value for many applications would be $\varepsilon \sim 10^{-4}$). Therefore one would wish to derive a new model as $\varepsilon \rightarrow 0$.

Eq. (2) is strongly similar to a reaction-diffusion equation with a strong reaction term. However a crucial difference here is that the reaction term is non local. As we will see this completely changes the behaviour of the solution.

This scaling was introduced in [20] and formally studied there. We briefly reproduce the main argument here. The starting point is to introduce a large deviation

scaling (which makes sense in view of the original probabilistic interpretation of the model)

$$\varphi_\varepsilon(t, x) = \varepsilon \log u_\varepsilon. \quad (3)$$

Then it is easy to see that Eq. (2) becomes

$$\partial_t \varphi_\varepsilon = r - I \star u_\varepsilon + H_\varepsilon(\varphi_\varepsilon), \quad (4)$$

where

$$H_\varepsilon(f) = \int_{\mathbb{R}} K(z) \left(e^{(bf(x+\varepsilon z) - bf(x))/\varepsilon} - 1 \right) dz. \quad (5)$$

Using the theory of viscosity solutions to Hamilton-Jacobi equations, one can pass to the limit in (4) and obtain for the limit φ of φ_ε

$$\partial_t \varphi = r - I \star u + H(\partial_x \varphi), \quad (6)$$

with u the weak limit (in the space of measure) of u_ε and

$$H_\varepsilon(f) = \int_{\mathbb{R}} K(z) \left(e^{(bf(x+\varepsilon z) - bf(x))/\varepsilon} - 1 \right) dz. \quad (7)$$

Of course at the limit, φ and u are no more connected by a relation like (3). Therefore Eq. (6) is no more closed and the question of how to recover u from φ is one of the main difficulties here, which is now fortunately better understood though.

1.3. The problem with Eq. (2). Let us focus here on one other delicate issue with this approach which is the main motivation for the current work. In the scaling under consideration, one has growth or decay of order $\exp(C/\varepsilon)$. In particular one can see that the ratio between the maximal value of u_ε and the value at most other points is of this order $\exp(C/\varepsilon)$. However if we come back to the starting point, which means a total population of $10^{10} - 10^{12}$ and $\varepsilon \sim 10^{-4}$, then there is an obvious problem. If (and it can be proved) u_ε is of order $\exp(C/\varepsilon)$ over a fixed interval of traits I then the total population over this interval includes in fact much less than one individual!

This has several consequences. First of all it means that given the scales under consideration here, the limit of the probabilistic models to the deterministic one is not fully justified (the total number of individuals would have to be much higher with respect to ε). This, in itself, could be ignored as anyway with 10^{12} individuals the obvious solution, use the probabilistic model, is not practical. Hence one could reasonably accept to still work with the deterministic model if its predictions were qualitatively in agreement with the behaviour of the stochastic one. That is not the case.

The first indication of a serious flaw comes from the case where we do not put any mutations in the model. Making $M_\varepsilon = 0$ in (2) still allows to perform the same analysis. In such a case, one would expect that nothing happens anymore at the limit as no evolution should be possible without mutations. However this is not the case and numerical simulations in particular show a remarkably similar behaviour between the cases with and without mutations (see [33] for example). This phenomenon is entirely due to the persistence of very small subpopulations which should actually be extinct but are kept by our deterministic model and can therefore re-emerge later if the conditions are right.

A second important problem concerns the issue of branching. Biologically speaking branching is the process by which one population divides itself into two (and

then possibly more) subpopulations. Mathematically one expects that at the limit, u will be a sum of moving Dirac masses

$$u = \sum_i \alpha_i(t) \delta_{x_i(t)}.$$

Branching then corresponds to the case when one Dirac mass becomes two. This phenomenon is one of the main motivation to study models like (2) instead of the adaptive dynamics approach for instance (see [18] for an introduction). At the limit (6), branching occurs at infinite speed, *i.e.* if Dirac mass $\alpha\delta_{x(t)}$ divides itself after some time t_0 into $\alpha_1\delta_{x_1(t)} + \alpha_2\delta_{x_2(t)}$ then

$$\frac{d}{dt}|x_2(t) - x_1(t)|_{t=t_0} = +\infty.$$

This is observed numerically but it can also be obtained through a formal analysis of the equation (we refer to [20]). Instead probabilistic models predict a finite speed at branching...

Those qualitative disagreements are important and should be corrected. One would then hope to derive models that are both able of dealing with very large populations and still treat correctly the small subpopulations. It is for the moment a completely open question of how to keep the stochastic effects for the small populations. But one could at least try to truncate the populations with less than 1 individual, which is expected to be enough to correct the qualitative flaws of the deterministic models.

There are already some attempts in this direction, see [33] and very recently [32]. The proposed correction in [33] consists in studying equations like

$$\partial_t u_\varepsilon(t, x) = \frac{1}{\varepsilon} (r(x) - I \star u_\varepsilon) u_\varepsilon(t, x) - \frac{1}{\varepsilon} \sqrt{\frac{u_\varepsilon}{\bar{u}_\varepsilon}} + M_\varepsilon(u_\varepsilon)(t, x). \quad (8)$$

The added mortality term has the effect of killing (in times of order ε) any population with a density less than \bar{u}_ε . From the modeling point of view, this is quite satisfactory as it corrects most of the problems with (2).

Unfortunately the mathematical analysis of an equation like (8) is for the moment untractable. The only situation that is understood is when \bar{u}_ε is chosen like $\exp(-\bar{\varphi}/\varepsilon)$. However this is exactly the scaling that was giving less than 1 individual and it is not satisfactory. Instead given our scalings, one would like to work with \bar{u}_ε that are polynomial in ε .

However that would mean truncating every value less than $C \varepsilon \log \varepsilon$ in φ_ε , which implies at the limit, as $\varphi \leq 0$, everything...

1.4. The proposed correction. We propose here a change in the model that that it completely corrects all the abnormal behaviours of (2) at the limit. In addition it is for the moment the only correction for which one can derive rigorously the limit. That means in particular that one can obtain numerical simulations for realistically low ε .

Unfortunately this correction is for the moment purely phenomenological. It cannot be derived while staying in the framework of asexual reproduction with pure competitive interactions.

It is inspired from the case with sexual reproduction where similar terms are rather common. There indeed exists a well known phenomenon in the case of sexual reproduction which drives to extinction small populations. If the population

is too small then the probability of meeting a partner is very low and hence the birth rate declines. This is the idea that we follow here.

Let us explain it first in this context of sexual reproduction. Consider a subpopulation with trait x , if the density of population for x , $u_\varepsilon(x)$ is below a critical value \bar{u}_ε then we assume that the probability for an individual with trait x to meet a partner is too low. Instead this individual will reproduce with an individual with a different trait y such that $u_\varepsilon(y)$ is large enough. Typically one can expect then that y should be the closest trait to x with a population large enough. However as the two traits are different, the individuals are not as compatible and the corresponding birth rate should decrease with the distance $|x - y|$.

Since we are mostly interested in organisms with asexual reproduction here, this phenomenon unfortunately does not seem to be very relevant. However it is possible to find the same type of effects by taking into account some cooperative interactions. Selection-mutation models rightly focus on competition between individuals as the main interaction mechanism in order to observe selection (and hence evolution). Nevertheless cooperation usually exists as well. In order to justify formally our correction term, we introduce some cooperative interactions in the model. We just assume that it takes place at smaller scales of population than the competitive interactions so that when looking at the dominant traits, competitive interactions still dominate.

We are for example led to add to the reproduction rate a cooperative effect of the kind

$$\min \left(C_0, \int_{\mathbb{R}} \Phi(|x - y|/\eta_\varepsilon) \frac{u_\varepsilon(y)}{\eta_\varepsilon} dy \right), \quad (9)$$

for a small parameter η_ε and a decreasing Φ . Note that the maximum effect of cooperation is capped (obviously birth rates cannot blow up).

The model would now become

$$\partial_t u_\varepsilon = \frac{1}{\varepsilon} \left(\tilde{r} + \min \left(C_0, \int_{\mathbb{R}} \Phi(|x - y|/\eta_\varepsilon) \frac{u_\varepsilon(y)}{\eta_\varepsilon} dy \right) - I \star u_\varepsilon \right) u_\varepsilon + M_\varepsilon(u_\varepsilon).$$

Of course to see an effect on the small population then it is necessary that the reproduction rate without cooperation be negative, which means here $\tilde{r} < 0$. In fact the actual reproduction rate observed for the large populations would be $r = \tilde{r} + C_0$ which then should be positive.

It is possible to simplify (9) (for numerical purposes for instance) while keeping the same structure and the same effects. When one combines it with the lower value of the reproduction rate, and assume for instance a linear Φ , then it is essentially possible to replace (9) by a regularization of

$$C_0 - K d(x, \{u_\varepsilon \geq \bar{u}_\varepsilon\}). \quad (10)$$

Those are the type of corrections that we consider here, with K large enough.

Adding a term of this kind to the model essentially keeps the competitive interaction between the individuals. Actually by taking $r = \tilde{r} + C_0$, one finds exactly the term given by Eq. (2) for populations $u_\varepsilon \geq \bar{u}_\varepsilon$. The new term only add a penalization for populations that are too small to benefit from some cooperation.

1.5. A brief overview of the various approaches for selection, mutation dynamics. The stochastic approach is based on individual-based models. As we mentioned before, they are related to evolutionary PDE models as those here or

in [16, 26] through a scaling of large population (see again [10]). Using a simultaneous scaling of large population and rare mutations, a stochastic limit process was obtained in [8] in the case of a monotype population (i.e. when the limit process can only be composed of a single Dirac mass), and in [13] when the limit population can be composed of finitely many Dirac masses. Other features can be added to those models, age-dependence for instance as in [29].

At the deterministic level, one approach consists in studying a simultaneous scaling of mutation and selection, in order to obtain a limit dynamics where transitions from a single Dirac mass to two Dirac masses could occur; the famous branching phenomenon that we have already mentioned. This is where the deterministic models presented here fit. More is said about the contributions in that area in the next subsection.

Another approach consists in completely separating the two scalings. One then tries to directly characterize evolutionary dynamics as sums of Dirac masses under biologically relevant parameter scalings, instead of obtaining it as some limit. This is a key point in *adaptive dynamics*—[25, 30, 31, 17, 9].

A classical way of justifying this form of the solution consists in studying the stationary behaviour of an evolutionary model involving a scaling parameter for mutations, and then letting this parameter converge to 0. The stationary state has been proved to be composed of one or several Dirac masses for various models (for deterministic PDE models, see [6, 7, 16, 26, 21], for Fokker-Planck PDEs corresponding to stochastic population genetics models, see [5], for stochastic models, see [36], for game-theoretic models, see [15]). Closely related to these works are the notions of ESS (evolutionarily stable strategies) and CSS (convergence stable strategies) [31, 18], which allow one in some cases to characterize stable stationary states [6, 16, 26, 15].

In this context the phenomenon of *evolutionary branching*, which is especially important for us, simply corresponds to the direct transition from a population composed of a single Dirac mass to a population composed of two Dirac masses, [31, 22, 23].

1.6. An overview of the works around Eq. (2). The scalings and the first formal results have been obtained in [20]. This was followed by several works on other models and on the corresponding Hamilton-Jacobi PDE [7, 34].

The main difficulty as mentioned before is the identification of the weak limit of u in terms of φ . There is usually one additional information which comes from uniform bounds on the total mass, it is that

$$\max_{\mathbb{R}} \varphi(t, \cdot) = 0, \quad \forall t.$$

One solution is then to try to see the undetermined $I \star u$ as a Lagrange multiplier for this additional constraint.

For example if φ solves (6) then our constraint should imply that $r - I \star u$ is non positive on the set $\{\varphi = 0\}$ and vanishes on at least one point.

Unfortunately, it is in general even formally not possible to identify $I \star u$ with that. For instance if φ attains its maximum at one point then one has only one constraint which is not enough in general.

There are however special cases of competitive interactions where this is enough. Suppose for instance that $I = 1$. Then $I \star u$ is just the total mass and formally one

simply expects that

$$\int_{\mathbb{R}} u(t, y) dy = \max_{x \text{ s.t. } \varphi(t,x)=0} r(x).$$

This is typical of so-called one resource interaction, meaning that the individuals interact only through one average quantity. In general that means considering interactions of the type

$$R \left(x, \int \eta(x) u(t, x) dx \right), \tag{11}$$

with R increasing in the x variable.

For models of the type we consider here, rigorous results (especially for the well posedness of the Hamilton-Jacobi eq. at the limit) mainly only exist in this case with just one resource, see [4] and [3], [27] (one resource but multidimensional traits).

However it is possible to extend the theory, see [11]. Formally one expects the limit u to satisfy the following conditions

$$\begin{aligned} (i) \quad & \text{supp } u(t, \cdot) \subset \{\varphi(t, \cdot) = 0\}, \\ (ii) \quad & r - I \star u \leq 0, \quad \text{on } \{\varphi(t, \cdot) = 0\}, \\ (iii) \quad & r - I \star u = 0, \quad \text{on } \text{supp } u(t, \cdot). \end{aligned} \tag{12}$$

This corresponds to the definition of an evolutionarily stable strategy. If one assumes a strong competition, *i.e.* that the operator $I \star$ is positive, then there exists a unique measure satisfying (12) (see [26]).

This is the approach followed here as well. With respect to the case with one resource, there are nevertheless additional difficulties. It is much harder to control the time oscillations of the reaction term, which is usually increasing and hence BV in time for one resource.

2. The model studied and the results. According to the previous considerations, we study the following equation

$$\partial_t u_\varepsilon(t, x) = \frac{1}{\varepsilon} (r - I \star_x u_\varepsilon(t, \cdot) - D_\varepsilon(u_\varepsilon)) u_\varepsilon(t, x) + M_\varepsilon(u_\varepsilon)(t, x), \tag{13}$$

where we recall that the mutation kernel M_ε reads

$$M_\varepsilon(f)(x) = \frac{1}{\varepsilon} \int_{\mathbb{R}} K(z) (f(x + \varepsilon z) - f(x)) dz, \tag{14}$$

for a smooth K such that $\int_{\mathbb{R}} zK(z) dz = 0$. Following [20], one defines φ_ε as

$$u_\varepsilon = e^{\varphi_\varepsilon/\varepsilon}, \quad \text{or } \varphi_\varepsilon = \varepsilon \log u_\varepsilon, \tag{15}$$

and obtains the equation

$$\partial_t \varphi_\varepsilon = r - I \star u_\varepsilon + H_\varepsilon(\varphi_\varepsilon), \tag{16}$$

with

$$H_\varepsilon(f) = \int_{\mathbb{R}} K(z) \left(e^{(f(x+\varepsilon z)-f(x))/\varepsilon} - 1 \right) dz. \tag{17}$$

We now detail the various assumptions that we require on the many terms present in the model.

Assumptions on the reproduction term and the distribution of initial traits. First of all we need to make sure that only the traits in a compact interval are important (to avoid traveling waves effects for instance). This can be simply ensured by asking all traits out of an interval to have a negative reproduction rate. Our assumption on r is hence

$$r \in L^\infty \cap C^2(\mathbb{R}), \quad \exists R > 0, \quad \exists r_0 > 0, \quad \forall |x| > R, \quad r(x) \leq -r_0. \quad (18)$$

Of course the whole population should not vanish immediately, and it is necessary that a non negligible part be concentrated on $[-R, R]$. The total population should also be bounded which leads to

$$\begin{aligned} \sup_\varepsilon \int_{\mathbb{R}} u_\varepsilon(t=0, \cdot) dx &< \infty, \\ \exists R_0 \quad \min_{|x| < R_0} r(x) &> 0, \quad \inf_\varepsilon \int_{-R_0}^{R_0} u_\varepsilon(t=0, x) dx > 0. \end{aligned} \quad (19)$$

Assumption on the interaction kernel I : Strong competition and concentration. We need an assumption of what we call “strong competition” on I , namely

$$I \in C_c^2, \quad \forall f \in M^1(\mathbb{R}) \setminus \{0\}, \quad \int_{\mathbb{R}^2} I(x-y) df(x) df(y) > 0, \quad (20)$$

which means that the Fourier transform \hat{I} of I is positive.

We also need to ensure some sort of symmetry for the interactions between individuals, for instance to guarantee that if an individual with trait x influences one with trait y then the converse is true (*i.e.* $I(y-x) \neq 0$ if $I(x-y) \neq 0$). In its most general form, see [26], this assumption reads

$$\exists M(x), N(x) \geq 0, \quad I(y-x) = M(x) I(x-y) N(y). \quad (21)$$

It is essential to obtain entropies for the competition term in the system. To keep things as straightforward as possible here, we assume the simpler

$$I(-x) = I(x). \quad (22)$$

This for instance allows us to define uniquely the ESS as per

Proposition 1. *Assume (20), (18), (22) and that $r, I \in C(\mathbb{R})$. For any closed $\Omega \subset \mathbb{R}$, there exists a unique finite nonnegative measure $\mu(\Omega)$ satisfying*

- i) $\text{supp } \mu \subset \Omega$*
- ii) $r - I \star \mu \leq 0$ in Ω , $r - I \star \mu = 0$ on $\text{supp } \mu$.*

Note however that sometimes, one may have uniqueness of the environmental variables whereas the population measure is not unique (and (20) is violated), most of our method would remain valid in such a case. This is the situation of a single resource, where almost nothing is required, see [4]. Nevertheless in more general situations, the conditions for which this kind of property holds without (20) are not currently identified.

We need an additional assumption to make sure that the ESS can only be concentrated on a set of measure 0

$$\begin{aligned} \exists S \in C(\mathbb{R}_+) \text{ with } S(0) = 0 \text{ s.t. } \forall \mu \in M_+^1(\mathbb{R}_+) \\ |\{x, |r(x) - I \star_x \mu(x)| \leq \nu\}| \leq S(\nu). \end{aligned} \quad (23)$$

This condition is in part technical and is required to avoid some time oscillations. However it also corresponds to the natural biological idea that only a few traits can be present at a given time.

It is probably hard to check (23) in specific models, but it is at least satisfied in large classes of parameters. One easy example is if the derivatives $r^{(k)}$ and $I^{(k)}$ are positive (or negative) on the interval $[-R, R]$ for some k . It has in fact been proved, see [24], that generically in r and I the ESS is discrete (a finite sum of Dirac masses) and hence (23) should be satisfied.

Assumptions on the correction D_ε . As for D_ε , we assume that there exists a critical scale $\bar{\varphi}_\varepsilon$ s.t.

$$\begin{aligned} \sup_\varepsilon \sup_{t,x} (|D_\varepsilon| + |\partial_x D_\varepsilon(t, x)|) < \infty, \quad \sup_\varepsilon \sup_{t,x} \bar{\varphi}_\varepsilon |\partial_{xx} D_\varepsilon(t, x)| < \infty, \\ D_\varepsilon(t, x) = 0 \quad \text{if} \quad \varphi_\varepsilon(t, x) \geq -\bar{\varphi}_\varepsilon \\ |\partial_x D_\varepsilon| \geq 2 |\partial_x r| \quad \text{if} \quad d(x, \{y, \varphi_\varepsilon(t, y) \geq -\bar{\varphi}_\varepsilon\}) \geq 2 \bar{\varphi}_\varepsilon. \end{aligned} \tag{24}$$

Those assumptions are compatible with corrections like (9) and (10). They introduce a new scale in the problem $-\bar{\varphi}_\varepsilon$ which corresponds to a critical population density of $\exp(-\bar{\varphi}_\varepsilon/\varepsilon)$. In line with our previous consideration, we would like to take this polynomial in ε which means $-\bar{\varphi}_\varepsilon \sim \varepsilon \log \varepsilon$ or more precisely for a constant C uniform in ε

$$\varepsilon \log \frac{1}{\varepsilon} \leq \bar{\varphi}_\varepsilon \leq C \varepsilon \log \frac{1}{\varepsilon}. \tag{25}$$

There is no need to be more precise on D_ε to study the properties of Eq. (16). However if one wants to identify the limit, it is necessary to specify what D_ε should look like at the limit. Therefore we make the additional assumption that there exists positive constants L, D_0 , for any fixed $f \in C(\mathbb{R})$, non positive

$$D_\varepsilon = D_\varepsilon(f) \longrightarrow \min(L d(x, \{f = 0\}), D_0) \quad \text{as} \quad \varepsilon \rightarrow 0. \tag{26}$$

This corresponds to (10) but other shapes would be possible and would essentially work the same.

Assumptions on the mutation kernel. We simply assume that

$$K \in C_c^2(\mathbb{R}), \quad K \geq 0, \quad \int_{\mathbb{R}} z K(z) dz = 0. \tag{27}$$

The main result. Formally, we can hence expect that as $\varepsilon \rightarrow 0$, Eq. (16) will lead to

$$\partial_t \varphi = r - I \star_x \mu(\{\varphi(t, \cdot) = 0\}) - \min(L d(x, \{f = 0\}), D_0) + H(\partial_x \varphi). \tag{28}$$

where the Hamiltonian H_ε became

$$H(p) = \int_{\mathbb{R}} K(z) (e^{p z} - 1) dz. \tag{29}$$

This is indeed what one can prove

Theorem 2.1. *Assume (18) on r , (19) on the bounds for the initial population, (20)-(22) on the interaction I together with (23), (24) on D_ε with (25)-(26) on the limit of D_ε , and (27) on the mutation kernel K . Finally assume that the initial data $u_\varepsilon(t = 0) > 0$ or $\varphi_\varepsilon(t = 0)$ are $W^{1,\infty} \cap C^2$, with*

$$\inf_\varepsilon \inf_{x \in \mathbb{R}} \partial_{xx} \varphi_\varepsilon(t = 0, x) > -\infty, \tag{30}$$

and that $\varphi_\varepsilon(t = 0, \cdot)$ converges to a function φ^0 for the norm $\|\cdot\|_{L^\infty(\mathbb{R})}$.

Then for any $T > 0$, up to the extraction of a subsequence in ε , φ_ε converges to some continuous φ uniformly on any compact subset of $[0, T] \times \mathbb{R}$ and φ is a solution to (28) almost everywhere in t, x and in the viscosity sense with initial condition $\varphi(t = 0, \cdot) = \varphi^0$. In particular the function $I \star u_\varepsilon$ converges to $I \star u$ in $L^p([0, T], C(\mathbb{R}))$ for any $p < \infty$, where $u(t, \cdot) = \mu(\{\varphi(t, \cdot) = 0\})$ is defined from φ by Prop. 1 and is continuous in time.

From a practical point of view, computing the solution u_ε of Eq. (13) is often too costly for small ε . This result allows to approximate the population density u_ε for small ε by the simpler $\mu(\{\varphi(t, \cdot) = 0\})$, where φ may be obtained by a discretization of (28), in the manner of those done in [20]. Rigorous numerical analysis of this kind of Hamilton-Jacobi equations is however still very preliminary.

Section 3 gives a short sketch of the proof of Prop. 1 and can be safely ignored if one is familiar with [26], [35].

The proof of Theorem 2.1 is given in the next section and being quite technical is divided into several lemmas. Lemma 4.0.1 just corresponds to the classical a priori estimates on equations like (16). Lemma 4.0.11 essentially follows the step of [11] and can also be skipped if the reader is familiar with that work.

In the proofs below, C denotes a numerical constant which may change from line to line but which only depends on T , norms of the initial data or the coefficients, but which is always uniform on ε .

We define as usual the distance between a point and a set

$$d(x, \Omega) = \inf_{y \in \Omega} |x - y|.$$

We also define the semi distance $\delta(O_1, O_2)$ between two sets O_1 and O_2 as usual by

$$\delta(O_1, O_2) = \sup_{x \in O_1} \inf_{y \in O_2} |x - y|.$$

A small $\delta(O_1, O_2)$ indicates that O_1 is almost included in \bar{O}_2 and in particular $\delta(O_1, O_2) = 0$ iff $O_1 \subset \bar{O}_2$.

We denote by $M^1(\omega)$ the set of signed Radon measures on the subset ω of \mathbb{R} equipped with the total variation norm.

3. Sketch of the proof of Prop. 1. A complete proof of this proposition can be found in [26], and [35]. We only show uniqueness and give a rough sketch of the existence part. For a given compact set Ω , assume that we can find two measures μ_1 and μ_2 with support in Ω and such that

$$r - I \star \mu_i = 0 \quad \text{on } \text{supp } \mu_i, \quad r - I \star \mu_i \leq 0 \quad \text{on } \Omega.$$

Compute

$$0 \geq \int (r - I \star \mu_1) d\mu_2 = \int I \star (\mu_2 - \mu_1) d\mu_2.$$

Summing with the symmetric term, one obtains

$$\int I \star (\mu_2 - \mu_1) (d\mu_2 - d\mu_1) \leq 0.$$

However since $\hat{I} > 0$ then the corresponding quadratic form is positive and we can conclude that $\mu_1 = \mu_2$.

For the existence, one considers the equation

$$\partial_t \mu_\varepsilon = \frac{1}{\varepsilon}(r - I \star \mu_\varepsilon) \mu_\varepsilon + \varepsilon M(u_\varepsilon), \quad \mu_\varepsilon(t = 0) = \mu(t = 0),$$

for a well chosen initial data $\mu(t = 0)$. If Ω is fully discrete, $\Omega = \{x_1, \dots, x_n\}$ then one may choose $\mu(t = 0) = \sum_i \delta_{x_i}$. If Ω is an interval then one simply takes $\mu(t = 0) = 1$. The general case is trickier and consists in taking an initial measure $\mu(t = 0)$ s.t. for any $\delta > 0$ and $x_0 \in \Omega$

$$\int_{\Omega \cap [x_0 - \delta, x_0 + \delta]} d\mu(t = 0) > 0.$$

It is then easy to obtain lower and upper bounds for the total mass $\int \mu_\varepsilon(t, dx)$.

Note that though there is also a mutation term here, its scaling is completely different from (13). The scaling does not involve small mutations and moreover the mutation rate is of order ε (instead of 1) thus vanishing at the limit.

Passing to the weak limit, $\mu_\varepsilon \rightarrow \mu$, gives a measure that has the desired properties (but is by no means easy to show).

4. Proof of Theorem 2.1.

4.1. A priori estimates. We start by stating and proving the obvious a priori estimates that one can obtain for the problem. Those essentially follow the lines of previous works.

We show the following estimates on the solution to (16)

Lemma 4.0.1. *Let φ_ε be a solution to (16) with the assumptions of Theorem 2.1. Then for any $T > 0$*

$$\|\partial_t \varphi_\varepsilon\|_{L^\infty([0, T] \times \mathbb{R})} + \|\partial_x \varphi_\varepsilon\|_{L^\infty([0, T], L^\infty(\mathbb{R}))} \leq C_T, \tag{31}$$

$$\forall t \leq T, x \in \mathbb{R}, \quad \partial_{xx} \varphi_\varepsilon(t, x) \geq -\frac{C}{\varphi_\varepsilon}, \quad H_\varepsilon(\varphi_\varepsilon) \geq -C \frac{\varepsilon}{\varphi_\varepsilon}, \tag{32}$$

$$\forall t \leq T, \frac{1}{C} \leq \int_{\mathbb{R}} u_\varepsilon(t, x) dx \leq C, \quad \varphi_\varepsilon(t, x) \leq \varepsilon \log 1/\varepsilon + C \varepsilon, \tag{33}$$

where C only depends on the time T , $\int_{\mathbb{R}} e^{\varphi_\varepsilon^0(x)/\varepsilon} dx$, $\|\partial_x \varphi_\varepsilon^0\|_{L^\infty(\mathbb{R})}$ and the infimum of $\partial_{xx} \varphi_\varepsilon^0(x)$. Finally φ_ε has level sets, uniformly bounded in ε .

Proof. Almost all proofs here are taken directly from [11] and reproduced for the sake of completeness. Some resulting bounds are different and much worse than in this former article, more precisely the lower bounds on $\partial_{xx} \varphi_\varepsilon$ and H_ε . Nevertheless even for those, the proofs, i.e. the way to obtain the bounds, are very close. As such we may skip some technical details.

Step 0: Upper Bound on the total mass. First notice that because of (18), there exists $R > 0$ s.t.

$$\forall |x| > R, \quad r(x) - I \star u_\varepsilon(t, x) \leq -r_0.$$

Let ψ be a smooth test function with support in $|x| > R$, taking values in $[0, 1]$ and equal to 1 on $|x| > R + 1$. Using the previous bound, we compute

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} \psi(x) u_\varepsilon(t, x) dx &\leq -\frac{r_0}{\varepsilon} \int_{\mathbb{R}} \psi(x) u_\varepsilon(t, x) dx \\ &\quad + \frac{1}{\varepsilon} \int_{\mathbb{R}^2} K(z) (\psi(x - \varepsilon z) - \psi(x)) u_\varepsilon(t, x) dz dx \\ &\leq -\frac{r_0}{\varepsilon} \int_{\mathbb{R}} \psi(x) u_\varepsilon(t, x) dx + C \int_{\mathbb{R}} u_\varepsilon(t, x) dx. \end{aligned}$$

On the other hand, on the bounded domain $[-R-1, R+1]$ as $I(x) > 0$ for all x , one has for some constant C

$$\forall |x| < R+1, \quad I \star u_\varepsilon(t, x) \geq C \int_{\mathbb{R}} (1 - \psi) u_\varepsilon dx.$$

Therefore with the same kind of estimate

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} (1 - \psi(x)) u_\varepsilon(t, x) dx &\leq C \int_{\mathbb{R}} u_\varepsilon(t, x) dx \\ &+ \frac{1}{\varepsilon} \left(\sup r - C \int_{\mathbb{R}} (1 - \psi) u_\varepsilon dx \right) \int_{\mathbb{R}} (1 - \psi(x)) u_\varepsilon(t, x) dx. \end{aligned}$$

Summing the two

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} u_\varepsilon(t, x) dx &\leq \frac{1}{\varepsilon} \left(\sup r - C \int_{\mathbb{R}} (1 - \psi) u_\varepsilon dx \right) \int_{\mathbb{R}} (1 - \psi) u_\varepsilon dx \\ &- \frac{r_0}{\varepsilon} \int_{\mathbb{R}} \psi(x) u_\varepsilon(t, x) dx + C \int_{\mathbb{R}} u_\varepsilon(t, x) dx. \end{aligned}$$

Since the sum of the first two terms of the r.h.s. is negative if $\int u_\varepsilon$ is larger than a constant independent of ε , this shows that $\int u_\varepsilon(t, x) dx$ remains uniformly bounded on any finite time interval.

Step 1: Bound on $\partial_x \varphi_\varepsilon$. This is a usual bound for solutions to Hamilton-Jacobi equations and we have to check that it remains true uniformly at the ε level. We follow exactly [11] for instance. Compute

$$\begin{aligned} \partial_t \partial_x \varphi_\varepsilon &= \partial_x r - \partial_x I \star u_\varepsilon - \partial_x D_\varepsilon \\ &+ \int_{\mathbb{R}} K(z) e^{\frac{\varphi_\varepsilon(t, x + \varepsilon z) - \varphi_\varepsilon(t, x)}{\varepsilon}} \frac{\partial_x \varphi_\varepsilon(t, x + \varepsilon z) - \partial_x \varphi_\varepsilon(t, x)}{\varepsilon} dz. \end{aligned} \quad (34)$$

By our assumptions and the upper bound on the total mass

$$|\partial_x r - \partial_x I \star u_\varepsilon - \partial_x D_\varepsilon| \leq C.$$

First note that this shows that $\|\partial_x \varphi_\varepsilon(t, \cdot)\|_{L^\infty}$ remains finite over a (possibly very short) time interval $[0, t_\varepsilon]$.

Now we use the classical maximum principle. Fix $t \in [0, T]$ such that $C_{\varepsilon, t} := \|\partial_x \varphi_\varepsilon(t, \cdot)\|_{L^\infty(\mathbb{R})} < \infty$. For any $x \in \mathbb{R}$ such that $\partial_x \varphi_\varepsilon(t, x) > \sup_y \partial_x \varphi_\varepsilon(t, y) - \alpha$, where the constant $\alpha > 0$ will be specified later, we have

$$\partial_t \partial_x \varphi_\varepsilon(t, x) \leq C + \int_{\mathbb{R}} K(z) e^{|z| C_{t, \varepsilon}} \frac{\alpha}{\varepsilon} dz \leq C \left(1 + \frac{\alpha}{\varepsilon} e^{C C_{t, \varepsilon}} \right).$$

Therefore, choosing $\alpha = \varepsilon e^{-C C_{t, \varepsilon}}$, we obtain

$$\frac{d}{dt} \sup_x \partial_x \varphi_\varepsilon(t, x) \leq C,$$

for a constant C independent of $t < t_\varepsilon$ and of ε . Using a similar argument for the minimum, we deduce that $t_\varepsilon > T$ and that $\partial_x \varphi_\varepsilon$ is bounded on $[0, T] \times \mathbb{R}$ by a constant depending only on T and $\|\partial_x \varphi_\varepsilon^0\|_{L^\infty(\mathbb{R})}$.

Step 2: First bound on $H_\varepsilon(\varphi_\varepsilon)$ and bounds on $\partial_t \varphi_\varepsilon$ and φ_ε . First remark that

$$\begin{aligned} - \int_{\mathbb{R}} K(z) dz &\leq H_\varepsilon(\varphi_\varepsilon(t))(x) = \int_{\mathbb{R}} K(z) e^{\frac{\varphi_\varepsilon(t, x + \varepsilon z) - \varphi_\varepsilon(t, x)}{\varepsilon}} dz - \int_{\mathbb{R}} K(z) dz \\ &\leq \int_{\mathbb{R}} K(z) e^{|z| \|\partial_x \varphi_\varepsilon\|_{L^\infty([0, T], \mathbb{R})}} dz \leq C. \end{aligned}$$

This is not optimal, more precisely the lower bound is atrocious, but will suffice for the moment.

Directly from Eq. (16),

$$|\partial_t \varphi_\varepsilon| \leq \sup |r| + \sup |I| \int_{\mathbb{R}} u_\varepsilon dx + |D_\varepsilon| + C \leq C,$$

hence ending the proof of the full Lipschitz bound on φ_ε .

To get the upper bound on φ_ε , we use this Lipschitz bound to get

$$\varphi_\varepsilon(t, y) \geq \varphi_\varepsilon(t, x) - C |y - x|,$$

so

$$\int_{\mathbb{R}} u_\varepsilon(t, y) dy \geq \int_{\mathbb{R}} e^{\varphi_\varepsilon(t, x)/\varepsilon} e^{-C |y-x|/\varepsilon} dy \geq 2C^{-1} \varepsilon e^{\varphi_\varepsilon(t, x)/\varepsilon}.$$

Hence the bound on the total mass yields that $\varphi_\varepsilon \leq \varepsilon \log 1/\varepsilon + C\varepsilon$.

Step 3: Lower bounds on $\partial_{xx} \varphi_\varepsilon$ and H_ε . Here we start paying for the introduction of an additional mortality term as the second derivative of D_ε is not bounded uniformly in ε . As before we use a maximum principle, from (16) and (24)

$$\begin{aligned} \partial_t \partial_{xx} \varphi_\varepsilon &\geq -\frac{C}{\bar{\varphi}_\varepsilon} + \int_{\mathbb{R}} K(z) e^{\frac{\varphi_\varepsilon(t, x+\varepsilon z) - \varphi_\varepsilon(t, x)}{\varepsilon}} \frac{\partial_{xx} \varphi_\varepsilon(t, x + \varepsilon z) - \partial_{xx} \varphi_\varepsilon(t, x)}{\varepsilon} dz \\ &\quad + \int_{\mathbb{R}} K(z) e^{\frac{\varphi_\varepsilon(t, x+\varepsilon z) - \varphi_\varepsilon(t, x)}{\varepsilon}} \frac{(\partial_x \varphi_\varepsilon(t, x + \varepsilon z) - \partial_x \varphi_\varepsilon(t, x))^2}{\varepsilon} dz. \end{aligned}$$

The last term is of course non negative and so with the same argument as before, we get

$$\frac{d}{dt} \inf_x \partial_{xx} \varphi_\varepsilon(t, x) \geq -\frac{C}{\bar{\varphi}_\varepsilon}.$$

This proves the uniform lower bound on $\partial_{xx} \varphi_\varepsilon$. Let us turn to the sharp lower bound (32) on $H_\varepsilon(\varphi_\varepsilon)$. Let us write

$$H_\varepsilon(\varphi_\varepsilon) = \int_{\mathbb{R}} K(z) \exp\left(\int_0^1 z \partial_x \varphi_\varepsilon(t, x + \theta z \varepsilon) d\theta\right) dz - \int_{\mathbb{R}} K(z) dz.$$

By differentiating once more

$$\begin{aligned} \int_0^1 z \partial_x \varphi_\varepsilon(t, x + \theta z \varepsilon) d\theta &\geq z \partial_x \varphi_\varepsilon(t, x) \\ &\quad + \int_0^1 z \int_0^1 \theta z \varepsilon \partial_{xx} \varphi_\varepsilon(t, x + \theta' \theta z \varepsilon) d\theta' d\theta \\ &\geq z \partial_x \varphi_\varepsilon(t, x) - C \frac{\varepsilon}{\bar{\varphi}_\varepsilon} z^2. \end{aligned}$$

Eventually

$$\begin{aligned} H_\varepsilon(\varphi_\varepsilon) &\geq \int_{\mathbb{R}} K(z) \exp(z \partial_x \varphi_\varepsilon(t, x) - C \varepsilon z^2 / \bar{\varphi}_\varepsilon) dz - \int_{\mathbb{R}} K(z) dz \\ &\geq H(\partial_x \varphi_\varepsilon(t, x)) - C \frac{\varepsilon}{\bar{\varphi}_\varepsilon}, \end{aligned}$$

where H is defined as in (29) and since K is compactly supported. Because we assumed that $\int_{\mathbb{R}} zK(z) dz = 0$, we have $H(p) \geq 0$ for any p , which gives the final bound.

Step 4: φ_ε has uniform compact level sets.

Observe that $\varphi_\varepsilon(t = 0, x) \rightarrow -\infty$ when $x \rightarrow \pm\infty$ since $\int_{\mathbb{R}} u_\varepsilon(t = 0, x) dx < \infty$ and

$\partial_x \varphi_\varepsilon(t = 0)$ is bounded. Because of the uniform convergence of $\varphi_\varepsilon(t = 0)$ to φ^0 on \mathbb{R} , one moreover deduces that the convergence $\varphi_\varepsilon(t = 0, x) \rightarrow -\infty$ is uniform in ε .

Since $\partial_x \varphi_\varepsilon \in L^\infty([0, T], \mathbb{R})$, this remains true at any time $t \in [0, T]$, uniformly in ε .

Therefore, the set

$$\Omega_\varepsilon := \{(t, x) \in [0, T] \times \mathbb{R} : \varphi_\varepsilon(t, x) \geq -l\}$$

is bounded for any l , uniformly in ε .

Step 5: Lower bound on the total mass and φ_ε .

By the previous steps we know that

$$M_\varepsilon(u_\varepsilon) = \frac{u_\varepsilon}{\varepsilon} H_\varepsilon \geq -C \frac{u_\varepsilon}{\bar{\varphi}_\varepsilon}.$$

Note that $D_\varepsilon = 0$ whenever $u_\varepsilon \geq e^{-\bar{\varphi}_\varepsilon/\varepsilon}$. Therefore by (19), choose R_0 s.t. $\min_{|x| \leq K} r > 0$ and integrate (13)

$$\frac{d}{dt} \int_{|x| \leq K} u_\varepsilon dx \geq \frac{1}{\varepsilon} \left(\min_{|x| \leq K} r - C \frac{\varepsilon}{\bar{\varphi}_\varepsilon} - C \int_{\mathbb{R}} u_\varepsilon dx \right) \int_{|x| \leq K} u_\varepsilon dx.$$

This implies that by (19)

$$\log \int_{|x| \leq K} u_\varepsilon(t, x) dx \geq -C + \frac{1}{\varepsilon} \int_0^t \left(\min_{|x| \leq K} r - C \frac{\varepsilon}{\bar{\varphi}_\varepsilon} - C \int_{\mathbb{R}} u_\varepsilon(s, x) dx \right) ds,$$

or obviously

$$\log \int_{\mathbb{R}} u_\varepsilon(t, x) dx \geq -C + \frac{1}{\varepsilon} \int_0^t \left(\min_{|x| \leq K} r - C \frac{\varepsilon}{\bar{\varphi}_\varepsilon} - C \int_{\mathbb{R}} u_\varepsilon(s, x) dx \right) ds.$$

This allows us to conclude that the total mass remains bounded from below uniformly in ε .

In particular by step 4, this lower bound means that $\max \varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon$.

4.2. Passing to the limit in the equation: First steps. By the uniform bounds provided by Lemma 4.0.1, we can extract a subsequence in ε (still denoted with ε), and find a function φ on $[0, T] \times \mathbb{R}$ such that $\partial_t \varphi \in L^\infty([0, T] \times \mathbb{R})$, $\partial_x \varphi \in L^\infty([0, T] \times \mathbb{R})$, with $\max \varphi = 0$ satisfying by the Arzela-Ascoli theorem

$$\varphi_\varepsilon \longrightarrow \varphi \quad \text{uniformly in } C(J) \text{ for any compact } J \text{ of } [0, T] \times \mathbb{R}, \tag{35}$$

Note that by the upper and lower bounds on φ_ε , one has $\max \varphi = 0$.

Since u_ε is uniformly bounded in L^1 , it converges (still after an extraction) in the weak-* topology of measures to some $u \in L^\infty([0, T], M^1(\mathbb{R}))$. This in turn implies a weak convergence of $I \star u_\varepsilon$. Note that it is indeed only a weak convergence in spite of the convolution because it regularizes only in x and time oscillations are still possible.

D_ε is uniformly bounded and can hence be assumed to converge weakly to some $D \in L^\infty$.

We wish to use the notion of viscosity solution (see [4] for instance) and the uniform bound on $\partial_x \varphi_\varepsilon$ to obtain the convergence of $H_\varepsilon(\varphi_\varepsilon)$ to $H(\partial_x \varphi)$.

In order to do that, a natural idea is to try and follow the strategy of [20] and get that

$$\psi = \varphi + \int_0^t I \star u(s, \cdot) ds$$

should be a viscosity solution to

$$\partial_t \psi = r - D + H(\partial_x \psi - \int_0^t \partial_x I \star u(s, \cdot) ds).$$

Unfortunately this requires more than just weak convergence in L^∞ for D_ε ; it requires some uniform convergence.

In summary our candidate for the limit is for the moment

$$\partial_t \varphi = r - I \star u(t, \cdot) - D + H(\partial_x \varphi). \tag{36}$$

Unfortunately D and u are still unidentified. This is now our main aim but it will require a much more precise understanding and control of the time oscillations of the set where the population is concentrated.

In addition in order to derive the limit, one needs to obtain some uniform convergence on D_ε . And finally in order to prove that φ is a viscosity solution and not only ψ (see again [20]), we will need continuity in time for u .

4.3. Continuity in time of the set $\{x \mid \varphi = 0\}$. At the limit, it is possible to show that the points where the population is concentrated move at most at a finite speed given by

$$V = 2 \sup_\varepsilon \sup_{|\xi| \leq 2 \|\partial_x \varphi_\varepsilon\|_{L^\infty}} \frac{1}{|\xi|} \int e^{z\xi} K(z) dz - 1.$$

We comment on that again at the very end of the proof. However we do not have yet enough tools to prove such a strong statement. For the time being we will be satisfied in proving a continuity result. The first step is to do it at the limit

Lemma 4.0.2. *Assume that for some point x_0 , some $t > t_0$ and $\delta > 0$*

$$d(x_0, \{\varphi(s, \cdot) = 0\}) \geq \delta, \quad \forall s \in [t_0, t],$$

then $\varphi(t, x_0) < 0$.

Before turning to the next result, let us point out that Lemma 4.0.2 actually implies the following

$$\begin{aligned} &\exists \tau \in C(\mathbb{R}_+) \text{ with } \tau(0) = 0, \text{ s.t. } \forall s \geq t, \\ &\forall x \in \{\varphi(s, \cdot) = 0\}, \quad \exists y \in \{\varphi(t, \cdot) = 0\} \text{ with } |y - x| \leq \tau(s - t). \end{aligned} \tag{37}$$

One of the main idea in the following proofs is to follow the characteristics corresponding to the Hamiltonian H . However Eq. (16) involves the modified Hamiltonian H_ε which does not have characteristics per se (it is non local for example). Some modifications are hence needed and follow the usual ideas for parabolic problems, which is why we use the following lemma

Lemma 4.0.3. *Introduce the intermediary scale $\sqrt{\varepsilon |\bar{\varphi}_\varepsilon|} = \varepsilon/\bar{\varepsilon} = \bar{\varepsilon} \bar{\varphi}_\varepsilon$ and consider any interval $[a, b]$. Define $a(t) = a + (t - t_0) V/2$, $b(t) = b - (t - t_0) V/2$. We denote*

$$m_\varepsilon(t) = \max_{[a(t), b(t)]} \varphi_\varepsilon(t, \cdot).$$

Then

$$\begin{aligned} m_\varepsilon(t) - m_\varepsilon(t_0) &\geq \int_{t_0}^t \left(\min_{[a(t), b(t)]} (r(x) - I \star_x u_\varepsilon(s, x) - D_\varepsilon) - \frac{C}{\varepsilon} \bar{\varphi}_\varepsilon \right) ds \\ m_\varepsilon(t) &\leq \max_{[a - \bar{\varepsilon} \bar{\varphi}_\varepsilon, b + \bar{\varepsilon} \bar{\varphi}_\varepsilon]} \varphi_\varepsilon(t_0, \cdot) \\ &\quad + \int_{t_0}^t \max_{[a(s) - \bar{\varepsilon} \bar{\varphi}_\varepsilon, b(s) + \bar{\varepsilon} \bar{\varphi}_\varepsilon]} (r(x) - I \star_x u_\varepsilon(t, x) - D_\varepsilon + 2\bar{\varepsilon}) ds. \end{aligned}$$

Proof of Lemma 4.0.3. The part

$$\min_{[a(t), b(t)]} (r(x) - I \star_x u_\varepsilon(s, x) - D_\varepsilon) - \frac{C}{\varepsilon} \bar{\varphi}_\varepsilon \leq \frac{d}{dt} m_\varepsilon(s)$$

is direct once one recalls that for any $x \in [a(t), b(t)]$ then

$$\begin{aligned} \partial_t \varphi_\varepsilon(s, x) &= r(x) - I \star_x u_\varepsilon(s, x) - D_\varepsilon + H_\varepsilon(\varphi_\varepsilon)(s, x) \\ &\geq \min_{[a(t), b(t)]} (r(x) - I \star_x u_\varepsilon(s, x) - D_\varepsilon) - \frac{C}{\varepsilon} \bar{\varphi}_\varepsilon, \end{aligned}$$

by the lower bound (32).

Let us turn now to the upper bound which is trickier and involves the velocity V .

Consider χ_ε a regularization of $\frac{3}{2} \|\partial_x \varphi_\varepsilon\|_{L^\infty} (\cdot)_+$, with $(\cdot)_+$ the positive part, at the intermediary scale $\bar{\varepsilon} \bar{\varphi}_\varepsilon$, *i.e.*

$$\begin{aligned} \chi_\varepsilon(x) &= 0, \quad \text{if } x \leq 0, \\ \chi'_\varepsilon(x) &= \frac{3}{2} \|\partial_x \varphi_\varepsilon\|_{L^\infty}, \quad \text{if } x \geq \bar{\varepsilon} \bar{\varphi}_\varepsilon, \end{aligned} \tag{38}$$

together with the general bounds

$$\begin{aligned} 0 \leq \chi_\varepsilon &\leq \frac{3}{2} \|\partial_x \varphi_\varepsilon\|_{L^\infty} (x)_+ + \bar{\varepsilon} \bar{\varphi}_\varepsilon, \quad \chi_\varepsilon \geq \frac{3}{2} \|\partial_x \varphi_\varepsilon\|_{L^\infty} (x)_+ - \bar{\varepsilon} \bar{\varphi}_\varepsilon, \\ 0 \leq \chi'_\varepsilon(x) &\leq 2 \|\partial_x \varphi_\varepsilon\|_{L^\infty} \mathbb{I}_{x \geq 0} + \bar{\varepsilon} \bar{\varphi}_\varepsilon, \quad |\chi''_\varepsilon(x)| \leq \frac{C}{\varepsilon} \bar{\varepsilon}. \end{aligned} \tag{39}$$

With that one defines

$$\psi_\varepsilon(t, x) = -\chi_\varepsilon(a(t) - x) - \chi_\varepsilon(x - b(t)).$$

Note that ψ_ε satisfies the following inequation

$$\partial_t \psi_\varepsilon \leq -\frac{V}{2} |\partial_x \psi_\varepsilon|.$$

This has for consequence that $\forall z \in \text{supp } K \subset [-1, 1]$

$$\begin{aligned} &\int_{\mathbb{R}} \exp\left(\frac{|\psi_\varepsilon(t, x + \varepsilon z) - \psi_\varepsilon(t, x)|}{2\varepsilon}\right) K(z) dz - 1 \\ &\leq \int_{\mathbb{R}} e^{z \partial_x \psi_\varepsilon(t, x)} K(z) dz - 1 + \bar{\varepsilon} \leq \frac{V}{2} |\partial_x \psi_\varepsilon| + \bar{\varepsilon}. \end{aligned} \tag{40}$$

As φ_ε and ψ_ε are smooth functions, it is easy to study

$$\tilde{m}_\varepsilon(t) = \sup(\varphi_\varepsilon(t, \cdot) + \psi_\varepsilon(t, \cdot)) = \max(\varphi_\varepsilon(t, \cdot) + \psi_\varepsilon(t, \cdot)).$$

In general this maximum is attained at one (or several) point $x \in \omega_m(t)$. First note that at such a point, one has

$$\partial_x(\varphi_\varepsilon(t, x) + \psi_\varepsilon(t, x)) = 0.$$

By (38) this implies that

$$\omega_m(t) \subset [a(t) - \bar{\varepsilon}\bar{\varphi}_\varepsilon, b(t) + \bar{\varepsilon}\bar{\varphi}_\varepsilon]. \tag{41}$$

Now in general

$$\frac{d}{dt} \tilde{m}_\varepsilon(t) \leq \sup_{x \in \omega_m(t)} \partial_t(\varphi_\varepsilon(t, x) + \psi_\varepsilon(t, x)).$$

Note that by the definition of the maximum, $\forall z \in \text{supp } K$

$$\varphi_\varepsilon(t, x + \varepsilon z) \leq \varphi_\varepsilon(t, x) + \psi_\varepsilon(t, x) - \psi_\varepsilon(t, x + \varepsilon z).$$

Now for $x \in \omega_m(t)$

$$\begin{aligned} H_\varepsilon(\varphi_\varepsilon)(t, x) &= \int_{\mathbb{R}} \exp\left(\frac{\varphi_\varepsilon(t, x + \varepsilon z) - \varphi_\varepsilon(t, x)}{\varepsilon}\right) K(z) dz - 1 \\ &\leq \int_{\mathbb{R}} \exp\left(\frac{-\psi_\varepsilon(t, x + \varepsilon z) + \psi_\varepsilon(t, x)}{\varepsilon}\right) K(z) dz - 1 \\ &\leq \frac{V}{2} |\partial_x \psi_\varepsilon| + \bar{\varepsilon} \leq -\partial_t \psi_\varepsilon + \bar{\varepsilon}. \end{aligned}$$

Therefore

$$\begin{aligned} \frac{d}{dt} \tilde{m}_\varepsilon(t) &\leq \sup_{x \in \omega_m(t)} (r(x) - I \star_x u_\varepsilon(t, x) - D_\varepsilon) + \bar{\varepsilon} \\ &\leq \sup_{x \in [a(t) - C\bar{\varepsilon}\bar{\varphi}_\varepsilon, b(t) + C\bar{\varepsilon}\bar{\varphi}_\varepsilon]} (r(x) - I \star_x u_\varepsilon(t, x) - D_\varepsilon) + \bar{\varepsilon}, \\ &\leq \sup_{x \in [a(t), b(t)]} (r(x) - I \star_x u_\varepsilon(t, x) - D_\varepsilon) + 2\bar{\varepsilon} \end{aligned}$$

by (41).

Note that by (38) and (41)

$$m_\varepsilon(t) \leq \tilde{m}_\varepsilon(t) \leq \max_{[a(t) - \bar{\varepsilon}\bar{\varphi}_\varepsilon, b(t) + \bar{\varepsilon}\bar{\varphi}_\varepsilon]} m_\varepsilon(t),$$

which allows to conclude. □

Let us now turn to Lemma 4.0.2

Proof of Lemma 4.0.2.

Step 1: Semi-continuity of $\{\varphi = 0\}$

Let us start with the following crucial observation

$$\begin{aligned} \forall t, \exists \tau_t \in C(\mathbb{R}_+) \text{ with } \tau_t(0) = 0, \text{ s.t. } \forall s \geq t, \\ \forall x \in \{\varphi(s, \cdot) = 0\}, \exists y \in \{\varphi(t, \cdot) = 0\} \text{ with } |y - x| \leq \tau_t(s - t). \end{aligned} \tag{42}$$

This is a sort of semi-continuity for $\{\varphi = 0\}$. It is proved very simply by contradiction. If it were not true, then

$$\begin{aligned} \exists t, \exists \tau_0 > 0, \exists s_n \rightarrow t, s_n \geq t, \exists x_n \in \{\varphi(s_n, \cdot) = 0\}, \\ d(x_n, \{\varphi(t, \cdot) = 0\}) \geq \tau_0, \end{aligned}$$

where $d(x, \omega) = \inf_{y \in \omega} |x - y|$ is the usual distance.

Since all the x_n belong to a compact set, we can extract a converging subsequence $x_n \rightarrow x$. As φ is continuous, $\varphi(t, x) = 0$ or $x \in \{\varphi(t, \cdot) = 0\}$. On the other hand one would also have $d(x, \{\varphi(t, \cdot) = 0\}) \geq \tau_0$ which is contradictory.

Therefore the result of the lemma is obviously true if t is such that $\tau_{t_0}(t) < \delta$.

Step 2: The connection between τ_t and the Lemma.

One does not have in general a uniform control on τ_{t_0} . In that case, one would find a sequence t_n , a number $\delta > 0$ s.t. $\forall \eta > 0, \limsup \tau_{t_n}(\eta) > \delta$.

The result of Lemma 4.0.2 would precisely rule this out and we argue by contradiction. Denote by t the first time when such a jump occurs. That means that

- τ can be chosen uniform till t , *i.e.* $\exists \tau$ s.t. $\forall s < t$ and $\forall \eta \in [0, t - s]$, $\tau_s(\eta) \leq \tau(\eta)$.
- There is a jump at t of size $\delta > 0$, which means $\exists x_0, \exists t_0 < t$ s.t. $\varphi(t, x_0) = 0$ but

$$d(x_0, \{\varphi(s, \cdot) = 0\}) \geq \delta, \quad \forall s \in [t_0, t].$$

Note that one can take t_0 as close to t as one wishes and in particular we may freely assume that $t - t_0$ is small enough s.t. $\tau(t - t_0) < \delta/8$.

Step 3: The contradiction.

First of all, note that as $\varphi_\varepsilon \rightarrow \varphi$ in L^∞ norm, one has that for ε small enough and some interval I_ε with $|I_\varepsilon| \rightarrow 0$ as $\varepsilon \rightarrow 0$

$$d(x_0, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) \geq \frac{\delta}{2}, \quad \forall s \in [t_0, t] \setminus I_\varepsilon.$$

Indeed one observes that

$$\{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\} \subset \{\varphi(s, \cdot) \geq -\bar{\varphi}_\varepsilon - \|\varphi_\varepsilon - \varphi\|_{L^\infty}\}.$$

And by the continuity of φ

$$\delta(\{\varphi(s, \cdot) \geq -\bar{\varphi}_\varepsilon - \|\varphi_\varepsilon - \varphi\|_{L^\infty}\}, \{\varphi(s, \cdot) = 0\}) \rightarrow 0,$$

for almost every s .

In particular we point out that one may define I_ε with $|I_\varepsilon| \rightarrow 0$ and $\eta_\varepsilon \rightarrow 0$ s.t.

$$\delta(\{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}, \{\varphi(s, \cdot) = 0\}) \leq \eta_\varepsilon, \quad \forall s \in [t_0, t] \setminus I_\varepsilon. \quad (43)$$

Now define the following interval

$$I_1(s) = [x_0 - \delta/4 + (s - t_0)V/2, x_0 + \delta/4 + (s - t_0)V/2].$$

Moreover denote by y_0 the closest point on the left from x_0 in $\{\varphi(t_0, \cdot) = 0\}$. Assume for instance that $y_0 < x_0$.

Next for any ε small enough, and any $s \in [t_0, t]$, denote

$$y_\varepsilon(s) = \sup\{x < x_0, \varphi_\varepsilon(s, x) \geq -\bar{\varphi}_\varepsilon\}.$$

By (43), one has that $y_\varepsilon(s) \leq y_0 + \eta_\varepsilon + \tau(s - t_0)$ for any $s \in [t_0, t] \setminus I_\varepsilon$. The same result can be obtained for the closest point on the right.

By the assumption on x_0 in the Lemma, this implies that I_1 remains at a distance larger than $\delta/4 > C\bar{\varphi}_\varepsilon$ of $\{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}$.

By the properties of D_ε , we deduce that for any $s \in [s_0, t]$,

$$\begin{aligned} \max_{I_1(s)} r - I \star_x u_\varepsilon(s, \cdot) - D_\varepsilon &\leq r(y_\varepsilon) - I \star_x u_\varepsilon(s, y_\varepsilon) - D_\varepsilon(s, y_\varepsilon) - C\delta/4 \\ &\leq r(y_0) - I \star_x u_\varepsilon(s, y_0) - D_\varepsilon(s, y_0) - C(\delta/4 - \eta_\varepsilon - \tau(s - t_0)) \\ &\leq r(y_0) - I \star_x u_\varepsilon(s, y_0) - D_\varepsilon(s, y_0) - C\delta/16, \end{aligned}$$

by choosing ε small enough.

Now let us apply Lemma 4.0.3 to I_1 to deduce that

$$\begin{aligned} \max_{I_1(t)} \varphi_\varepsilon(t, \cdot) &\leq \int_{t_0}^t (r(y_0) - I \star_x u_\varepsilon(s, y_0) - D_\varepsilon(s, y_0)) ds \\ &\quad - C \frac{\delta}{16} (t - t_0) + C|I_\varepsilon| + \max_{[x_0 - 3\delta/4, x_0 + 3\delta/4]} \varphi_\varepsilon(t_0, \cdot). \end{aligned}$$

Note that by (32)

$$\partial_t \varphi_\varepsilon(s, y_0) \geq r(y_0) - I \star_x u_\varepsilon(s, y_0) - D_\varepsilon(s, y_0) - \frac{C\varepsilon}{\bar{\varphi}_\varepsilon}.$$

So

$$\begin{aligned} \varphi(t, x_0) &\leq \|\varphi - \varphi_\varepsilon\|_{L^\infty} + \max_{I_1(t)} \varphi_\varepsilon(t, \cdot) \leq \varphi_\varepsilon(t, y_0) - \varphi_\varepsilon(t_0, y_0) - C \frac{\delta}{16} (t - t_0) \\ &\quad + C |I_\varepsilon| + \|\varphi - \varphi_\varepsilon\|_{L^\infty} + \max_{[x_0-3\delta/4, x_0+3\delta/4]} \varphi_\varepsilon(t_0, \cdot). \end{aligned}$$

To conclude, just observe that $\varphi(t_0, y_0) = 0$ so that

$$\varphi_\varepsilon(t, y_0) - \varphi_\varepsilon(t_0, y_0) + C |I_\varepsilon| \leq \|\varphi - \varphi_\varepsilon\|_{L^\infty} + \varepsilon |\log \varepsilon| + C |I_\varepsilon| \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

Therefore taking ε small enough, one concludes that

$$\varphi(t, x_0) \leq \max_{[x_0-3\delta/4, x_0+3\delta/4]} \varphi(t_0, \cdot) + \|\varphi - \varphi_\varepsilon\|_{L^\infty} < 0,$$

which gives the desired contradiction. \square

4.4. Control of the oscillations. Before obtaining the continuity of the set $\{\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$ which is our main goal, we have to control the oscillations in time of the reaction term.

Lemma 4.0.4. *There exists a positive function $\alpha_\varepsilon(s)$ with $\int_0^T \alpha_\varepsilon^2(s) ds \leq 1$, s.t. for any $t_0 \in [0, T]$ and any $t > t_0$*

$$\|I \star u_\varepsilon(t, \cdot) - I \star u_\varepsilon(t_0, \cdot)\|_{L^\infty} \leq \frac{1}{\sqrt{\varepsilon}} \int_{t_0}^t \alpha_\varepsilon(s) ds.$$

With this control, we can first show that the reaction term $r - I \star u_\varepsilon$ essentially vanishes on the set $\{\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$

Lemma 4.0.5. *There exists $\tilde{\varepsilon} \rightarrow 0$, I_ε composed of at most $C\varepsilon^{-3/8}$ intervals of size $\varepsilon^{3/4}$ s.t. for any $t_0 \in [0, T] \setminus I_\varepsilon$, for any x_0 s.t.*

$$\exists t \in [t_0, t_0 + \varepsilon^{3/4}], \quad \varphi_\varepsilon(t, x_0) \geq -\bar{\varphi}_\varepsilon$$

then one has for any $s \in [t_0, t_0 + \varepsilon^{3/4}]$,

$$r(x_0) - I \star u_\varepsilon(s, x_0) \leq \tilde{\varepsilon}.$$

This last lemma enables us to be very precise concerning the growth rate at any point without mutations

Lemma 4.0.6. *For any t_0 and $t > t_0$ with $t - t_0 \geq C\tilde{\varepsilon}$, for any x_0 and any $\nu > C\sqrt{\tilde{\varepsilon}}$ then*

$$\begin{aligned} \text{either } &\int_{t_0}^t (r(x_0) - I \star (s, x_0) - D_\varepsilon(s, x_0)) ds \leq -C(t - t_0)(\nu^2 - \tilde{\varepsilon}), \\ \text{or } &\int_{t_0}^t D_\varepsilon(s, x_0) ds \leq C(t - t_0)\nu. \end{aligned}$$

With this control on the growth rate, it is possible to make sure that at most times the set $\{\varphi(t, \cdot) = 0\}$ is of measure 0 or more precisely

Lemma 4.0.7. For any $t_0 \in [0, T]$, define the set Ω_{t_0} by $x_0 \in \Omega_{t_0}$ iff

$$x_0 \in \{\varphi(t_0, \cdot) = 0\}, \quad \exists t_n \rightarrow t, \quad x_n \in \{\varphi(t_n, \cdot) = 0\} \text{ s.t. } x_n \rightarrow x_0.$$

Then $|\Omega_t| = 0$.

We now turn to the proofs of those results

Proof of Lemma 4.0.4. The proof uses one of the entropies of the system (and a different one from [11]). Start by the following

$$\begin{aligned} & \frac{d}{dt} \left[\int u_\varepsilon(t, x) I(x-y) u_\varepsilon(t, y) dx dy - 2 \int r u_\varepsilon(t, x) dx \right] \\ &= \frac{2}{\varepsilon} \int (r - I \star u_\varepsilon(t, \cdot)) u_\varepsilon(t, x) (I \star (u_\varepsilon(t, \cdot) - r)) dx \\ &+ \frac{2}{\varepsilon} \int D_\varepsilon(t, x) u_\varepsilon(t, x) (I \star u_\varepsilon(t, \cdot) - r) dx \\ &+ \int M_\varepsilon(u_\varepsilon)(t, x) (I \star u_\varepsilon(t, \cdot) - r) dx. \end{aligned}$$

Note that as $D_\varepsilon(t, x) = 0$ if $\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon$ then

$$D_\varepsilon u_\varepsilon \leq C e^{-\bar{\varphi}_\varepsilon/\varepsilon} \leq C \varepsilon.$$

Similarly for any smooth function ψ , by a simple change of variable

$$\int M_\varepsilon(u_\varepsilon) \psi(x) dx = \frac{1}{\varepsilon} \int K(z) u_\varepsilon(t, x) (\psi(x - \varepsilon z) - \psi(x)) dx \leq C \varepsilon \|\nabla \psi\|_{L^\infty}.$$

Hence one first obtains that

$$\frac{1}{\varepsilon} \int_0^T \int (r - I \star u_\varepsilon(t, \cdot))^2 u_\varepsilon(t, x) dx dt \leq C. \quad (44)$$

Now note that

$$\begin{aligned} I \star u_\varepsilon(t, \cdot) - I \star u_\varepsilon(t_0, \cdot) &= \frac{1}{\varepsilon} \int_{t_0}^t I \star ((r - I \star u_\varepsilon(s, \cdot)) u_\varepsilon(s, \cdot)) ds \\ &\quad - \frac{1}{\varepsilon} \int_{t_0}^t I \star (D_\varepsilon(s, \cdot) u_\varepsilon(s, \cdot)) ds \\ &\quad + \int_{t_0}^t I \star M_\varepsilon(u_\varepsilon(s, \cdot)) ds. \end{aligned}$$

With the same bounds as before, one obtains that

$$\begin{aligned} \|I \star (u_\varepsilon(t) - u_\varepsilon(t_0))\|_{L^\infty} &\leq C(t - t_0) + \frac{1}{\varepsilon} \int_{t_0}^t \int |r - I \star u_\varepsilon(s, \cdot)| u_\varepsilon(s, \cdot) dx ds \\ &\leq C(t - t_0) + \frac{1}{\sqrt{\varepsilon}} \int_{t_0}^t \left(\frac{1}{\varepsilon} \int (r - I \star u_\varepsilon(s, \cdot))^2 u_\varepsilon(s, \cdot) dx \right)^{1/2}, \end{aligned}$$

by Cauchy-Schwarz. Defining

$$\alpha_\varepsilon = C + \left(\frac{1}{\varepsilon} \int (r - I \star u_\varepsilon(s, \cdot))^2 u_\varepsilon(s, \cdot) dx \right)^{1/2},$$

allows to conclude. \square

Proof of Lemma 4.0.5. First of all, consider the intervals $I_i = [2i\varepsilon^{3/4}, 2(i+1)\varepsilon^{3/4}]$ for $i = 0 \dots T\varepsilon^{-3/4}/2$. As α_ε is uniformly bounded in L^2 then there are at most $C\varepsilon^{-3/8}$ indices i s.t.

$$\int_{I_i} \alpha_\varepsilon^2(s) ds \geq \varepsilon^{-3/4} \left(\int_{I_i} \alpha_\varepsilon(s) ds \right)^2 > \varepsilon^{3/8},$$

that is

$$\frac{1}{\sqrt{\varepsilon}} \int_{I_i} \alpha_\varepsilon(s) ds > \varepsilon^{1/16}.$$

Define I_ε as the union of the intervals $I_i = [2i\varepsilon^{3/4} - 2\varepsilon^{3/4}, 2(i+1)\varepsilon^{3/4} + 2\varepsilon^{3/4}]$ for such indices i .

Now for any $t_0 \notin I_\varepsilon$ and any $t \in [t_0, t_0 + \varepsilon^{3/4}]$, one has that

$$\frac{1}{\varepsilon} \int_{t-\varepsilon^{3/4}}^{t+\varepsilon^{3/4}} \alpha_\varepsilon(s) ds < \varepsilon^{1/16}.$$

Therefore by Lemma 4.0.4, $I \star u_\varepsilon(s)$ changes by at most $\varepsilon^{1/16}$ over the interval $[t - \varepsilon^{3/4}, t + \varepsilon^{3/4}]$.

Define

$$\tilde{\varepsilon} = \varepsilon^{1/16} + C \frac{\varepsilon}{\bar{\varphi}_\varepsilon} + \varepsilon^{1/8}.$$

Choosing any x_0 s.t. $\varphi_\varepsilon(t, x_0) \geq -\bar{\varphi}_\varepsilon$ and any $s \in [t - \varepsilon^{3/4}, t + \varepsilon^{3/4}]$, we argue by contradiction. If

$$r(x_0) - I \star u_\varepsilon(s, x_0) \geq \tilde{\varepsilon}.$$

Then at any other $s' \in [t - \varepsilon^{3/4}, t + \varepsilon^{3/4}]$, and in particular for $s' = t_0$,

$$r(x_0) - I \star u_\varepsilon(s', x_0) \geq C \frac{\varepsilon}{\bar{\varphi}_\varepsilon} + \varepsilon^{1/8}.$$

Note that $D_\varepsilon(t, x_0) = 0$ and hence

$$\partial_t \varphi_\varepsilon(t, x_0) \geq r(x_0) - I \star u_\varepsilon(t_0, x_0) - C \frac{\varepsilon}{\bar{\varphi}_\varepsilon} \geq \varepsilon^{1/8}.$$

By continuity that means that $\varphi_\varepsilon(s', x_0)$ is increasing in s' over a (possibly very small) interval $[t, t + \nu]$. Define ν such as to make this interval maximal. At $t + \nu$, $\varphi_\varepsilon(t + \nu, x_0) > \varphi_\varepsilon(t, x_0) \geq -\bar{\varphi}_\varepsilon$ which means that $D_\varepsilon(t + \nu, x_0) = 0$ and if $t + \nu \leq t + \varepsilon^{3/4}$

$$\partial_t \varphi_\varepsilon(t + \nu, x_0) \geq r(x_0) - I \star u_\varepsilon(t + \nu, x_0) - C \frac{\varepsilon}{\bar{\varphi}_\varepsilon} \geq \varepsilon^{1/8}.$$

This implies that $t + \nu > t + \varepsilon^{3/4}$, and hence that $\varphi_\varepsilon(s', x_0) \geq -\bar{\varphi}_\varepsilon$ on any $s' \in [t, t + \varepsilon^{3/4}]$. Therefore

$$\partial_t \varphi_\varepsilon(s', x_0) \geq \varepsilon^{1/8},$$

or finally

$$\varphi_\varepsilon(t + \varepsilon^{3/4}, x_0) \geq -\bar{\varphi}_\varepsilon + \varepsilon^{7/8},$$

which is impossible as the maximum of φ_ε is of order $\varepsilon |\log \varepsilon|$. □

We turn to the next proof

Proof of Lemma 4.0.6. For any $s \in [t_0, t] \setminus I_\varepsilon$, one has either that x_0 is at distance less than ν of $\{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}$ or it is at distance larger than ν . Let us decompose accordingly $[t_0, t]$ into $I_\varepsilon \cup I_\nu \cup J_\nu$ where I_ν consists of the set where x_0 is at distance less than ν of $\{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}$.

We observe that on J_ν by the properties of D_ε and since on $\{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}$, one has by Lemma 4.0.5 that $r - I \star u_\varepsilon(s, \cdot)$ is less than $\tilde{\varepsilon}$ then

$$r(x_0) - I \star u_\varepsilon(s, x_0) - D_\varepsilon(s, x_0) \leq \tilde{\varepsilon} - C\nu \leq -C\nu.$$

On the other hand, on I_ν then still by 4.0.5, one has that

$$r(x_0) - I \star u_\varepsilon(s, x_0) - D_\varepsilon(s, x_0) \leq \tilde{\varepsilon},$$

and moreover

$$D_\varepsilon(s, x_0) \leq C\nu.$$

Therefore

$$\int_{t_0}^t (r(x_0) - I \star u_\varepsilon(s, x_0) - D_\varepsilon(s, x_0)) ds \leq |I_\varepsilon| + C\tilde{\varepsilon}|I_\nu| - C\nu|J_\nu|,$$

and

$$\int_{t_0}^t D_\varepsilon(s, x_0) \leq C\nu|I_\nu| + C|J_\nu|.$$

We recall that $|I_\varepsilon| \leq \varepsilon^{3/8}$ which is asymptotically much smaller than $\tilde{\varepsilon}^2$. Therefore either $|J_\nu| \geq (t - t_0)\nu$ and we find the first possibility or $|J_\nu| \leq (t - t_0)\nu$ and we find the second one. \square

The next proof is

Proof of Lemma 4.0.7. Note first of all that by Lemma 4.0.2, one may in fact take any sequence $t_n \rightarrow t_0$ for any $x_0 \in \Omega_{t_0}$.

Choose any t_0 , any $\eta > 0$ and any x_0 in $\{\varphi(t_0, \cdot) = 0\}$. Observe that for any $t > t_0$

$$\varphi_\varepsilon(t, x_0) \geq \varphi_\varepsilon(t_0, x_0) + \int_{t_0}^t (r - I \star u_\varepsilon(s, x_0) - D_\varepsilon(s, x_0)) ds - C \frac{\varepsilon}{\bar{\varphi}_\varepsilon}.$$

Therefore we first deduce that

$$\int_{t_0}^t (r - I \star u_\varepsilon(s, x_0) - D_\varepsilon(s, x_0)) ds \leq C\varepsilon |\log \varepsilon| + C(t - t_0) \frac{\varepsilon}{\bar{\varphi}_\varepsilon} + \|\varphi_\varepsilon - \varphi\|_{L^\infty}. \quad (45)$$

Next find x_t the closest point from x_0 s.t. $\varphi(x_t) = 0$. By definition $x_t - x_0 \rightarrow 0$. We define the small interval around x_t for $t_0 \leq s \leq t$

$$I_s = [x_t - (t - s)V/2, x_t + (t - s)V/2].$$

By Lemma 4.0.3

$$\begin{aligned} -\|\varphi - \varphi_\varepsilon\|_{L^\infty} &\leq \varepsilon |\log \varepsilon| + \int_{t_0}^t (r(x_0) - I \star_x u_\varepsilon(s, x_0)) ds \\ &\quad + C(\bar{\varepsilon} + |x_0 - x_t| + (t - t_0))(t - t_0). \end{aligned}$$

This implies that

$$\begin{aligned} \int_{t_0}^t (r(x_0) - I \star_x u_\varepsilon(s, x_0)) ds &\geq -\|\varphi - \varphi_\varepsilon\|_{L^\infty} - \varepsilon |\log \varepsilon| \\ &\quad - C(\bar{\varepsilon} + |x_0 - x_t| + (t - t_0))(t - t_0). \end{aligned} \quad (46)$$

Now define ν s.t.

$$C(\nu^2 - \varepsilon) = \frac{\|\varphi - \varphi_\varepsilon\|_{L^\infty}}{t - t_0} + \frac{\varepsilon |\log \varepsilon|}{t - t_0} + C(\bar{\varepsilon} + |x_0 - x_t| + (t - t_0)).$$

Note that as long as ε is small enough one has that $\nu \geq \tilde{\varepsilon}$. Moreover ν can be chosen arbitrarily small by taking t close enough to t_0 . By applying Lemma 4.0.6 for this ν , (46) means that we are necessarily in the second case which means

$$\int_{t_0}^t D_\varepsilon(s, x_0) ds \leq C(t - t_0) \nu.$$

We combine this inequality to (45) to deduce that

$$\int_{t_0}^t (r - I \star u_\varepsilon(s, x_0)) ds \leq C(t - t_0) \left(\frac{\varepsilon}{\bar{\varphi}_\varepsilon} + \nu^2 \right). \tag{47}$$

Now, we define the average

$$\mu_\varepsilon = \frac{1}{t - t_0} \int_{t_0}^t u_\varepsilon(s, x_0) ds$$

and from (46) and (47), we get

$$|r(x_0) - I \star \mu_\varepsilon(x_0)| \leq C \left(\frac{\varepsilon}{\bar{\varphi}_\varepsilon} + \nu^2 \right), \quad \forall x_0 \in \Omega_{t_0}.$$

By the fundamental property (23), we deduce that

$$|\Omega_{t_0}| \leq S \left(C \left(\frac{\varepsilon}{\bar{\varphi}_\varepsilon} + \nu^2 \right) \right).$$

To conclude take $t \rightarrow t_0$ s.t. $\nu \rightarrow 0$ and then ε small enough to obtain that $|\Omega_{t_0}| = 0$. \square

4.5. Continuity in time of the set $\{x | \varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$. As a first consequence of our control over the oscillations of the reaction term, one may simply obtain the continuity in time of the set $\{\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$

Lemma 4.0.8. *There exists \tilde{I}_ε with $|\tilde{I}_\varepsilon| \rightarrow 0$ as $\varepsilon \rightarrow 0$ s.t. if for some $t_0 \in [0, T] \setminus \tilde{I}_\varepsilon$, any point x_0 , any $t > t_0$ and $\delta > 0$*

$$d(x_0, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) \geq \delta, \quad \forall s \in [t_0, t],$$

then for ε small enough with respect to δ ,

$$\varphi_\varepsilon(t, x_0) < -\bar{\varphi}_\varepsilon - C \delta (t - t_0).$$

for some constant C independent of ε .

Unfortunately Lemma 4.0.8 does not imply a result like (37) with a function τ uniform in ε . It guarantees that there cannot be jumps of significant size in the set $\{\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$ but that set could still be propagated very fast.

However we can combine it with Lemma 4.0.7 to finally deduce a uniform (in ε) continuity in time for the support of $\{\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$.

Lemma 4.0.9. *There exists \bar{I}_ε with $|\bar{I}_\varepsilon| \rightarrow 0$ as $\varepsilon \rightarrow 0$ s.t. for any $t_0 \in [0, T] \setminus \bar{I}_\varepsilon$, $\forall \delta > 0$, there exists $\tau > 0$ s.t. $\forall x_0$, if*

$$d(x_0, \{\varphi_\varepsilon(t_0, \cdot) \geq -\bar{\varphi}_\varepsilon\}) \geq \delta,$$

then for ε small enough with respect to δ and τ , for any $t < t_0 + \tau$

$$\varphi_\varepsilon(t, x_0) < -\bar{\varphi}_\varepsilon - C(t - t_0) < -\bar{\varphi}_\varepsilon.$$

Lemma 4.0.9 in particular directly implies

Corollary 1. For any $t_0 \in [0, T] \setminus \bar{I}_\varepsilon$, $\exists \tau_{t_0} \in C([0, T])$ with $\tau_{t_0}(0) = 0$ s.t. for any $t \geq t_0$

$$\delta(\{\varphi_\varepsilon(t, \cdot) \geq -\bar{\varphi}_\varepsilon\}, \{\varphi_\varepsilon(t_0, \cdot) \geq -\bar{\varphi}_\varepsilon\}) \leq \tau_{t_0}(t - t_0).$$

We turn to the proofs

Proof of Lemma 4.0.8. Let first characterize the “good” t_0 . We define

$$\tilde{I}_\varepsilon = \{t_0, \exists t > 0 \text{ with } |I_\varepsilon \cap [t_0, t]| \geq \bar{\varepsilon}|t - t_0|\}. \quad (48)$$

Note that of course $I_\varepsilon \subset \tilde{I}_\varepsilon$ (just take $t = t_0 + \varepsilon^{3/4}$). We can estimate $|\tilde{I}_\varepsilon|$ the following way: If $t_0 \in \tilde{I}_\varepsilon$ then there exists an index $k \geq 0$ s.t. $t_0 \in \tilde{I}_\varepsilon^k$ where \tilde{I}_ε^k is defined as the union of the intervals $[i 2^k \varepsilon^{3/4}, (i+1) 2^k \varepsilon^{3/4}]$ for those i s.t.

$$|I_\varepsilon \cap [i 2^k \varepsilon^{3/4}, i 2^k \varepsilon^{3/4} + 2^k \varepsilon^{3/4}]| \geq 2^{k-3} \bar{\varepsilon} \varepsilon^{3/4}.$$

Each \tilde{I}_ε^k is composed of at most n_k such intervals and since $|I_\varepsilon| \leq C \varepsilon^{3/8}$, one has that

$$n_k \leq 2^{3-k} C \varepsilon^{-3/8} \bar{\varepsilon}^{-1}.$$

Consequently

$$|\tilde{I}_\varepsilon^k| \leq C \varepsilon^{3/8} \bar{\varepsilon}^{-1}.$$

Since $k \leq C |\log \varepsilon|$, we conclude that

$$|\tilde{I}_\varepsilon| \leq C \varepsilon^{3/8} \bar{\varepsilon}^{-1} |\log \varepsilon|,$$

which indeed converges to 0 as $\varepsilon \rightarrow 0$.

Now define $y_\varepsilon^-(s)$ similarly as in the proof of Lemma 4.0.2 by

$$y_\varepsilon^-(s) = \sup\{y < x_0, \varphi_\varepsilon(s, y) \geq -\bar{\varphi}_\varepsilon\},$$

and y_ε^+ by

$$y_\varepsilon^+(s) = \sup\{y > x_0, \varphi_\varepsilon(s, y) \geq -\bar{\varphi}_\varepsilon\}.$$

Define again the interval

$$I_1(s) = [x_0 - \delta/2 + (s - t_0)V/2, x_0 + \delta/2 + (s - t_0)V/2].$$

It remains non empty for $s \in [t_0, t]$.

For $s \in [t_0, t] \setminus I_\varepsilon$, one has that

$$\begin{aligned} \max_{I_1(s)} (r - I \star_x u_\varepsilon(s, \cdot) - D_\varepsilon) &\leq \max_{\pm} r - I \star_x u_\varepsilon(s, \cdot) - D_\varepsilon(s, \cdot) \Big|_{y_\varepsilon^\pm(s)} \\ &\quad - C \delta. \end{aligned}$$

Apply now Lemma 4.0.5, if $s \notin I_\varepsilon$ then this implies that

$$\max_{I_1(s)} (r - I \star_x u_\varepsilon(s, \cdot) - D_\varepsilon) \leq \tilde{\varepsilon} - C \delta \leq -C \delta,$$

provided that ε is small enough with respect to δ .

We now apply Lemma 4.0.3 to I_1 to deduce that

$$\begin{aligned} \max_{I_1(t)} \varphi_\varepsilon(t, \cdot) &\leq 2 \bar{\varepsilon} (t - t_0) - C \delta (t - t_0 - |I_\varepsilon \cap [t_0, t]|) \\ &\quad + C |I_\varepsilon \cap [t_0, t]| + \max_{[x_0 - 3\delta/4, x_0 + 3\delta/4]} \varphi_\varepsilon(t_0, \cdot). \end{aligned}$$

Now since $t_0 \notin \tilde{I}_\varepsilon$ then

$$|I_\varepsilon \cap [t_0, t]| \leq \bar{\varepsilon} (t - t_0), \quad t - t_0 - |I_\varepsilon \cap [t_0, t]| \geq \frac{t - t_0}{2}.$$

Hence

$$\max_{I_1(t)} \varphi_\varepsilon(t, \cdot) \leq -\bar{\varphi}_\varepsilon - (t - t_0)(C\delta - C\bar{\varepsilon}) \leq -\bar{\varphi}_\varepsilon - C\delta(t - t_0),$$

provided that ε is small enough with respect to δ , which concludes the proof. \square

Proof of Lemma 4.0.9. By Lemma 4.0.2 and Lemma 4.0.7, we know that for a.e. t_0 then

$$|\{\varphi(t_0, \cdot)\}| = 0.$$

Let us choose such a t_0 with as well $t_0 \notin \tilde{I}_\varepsilon$ and denote the corresponding interval \bar{I}_ε . Now for any δ , divide the domain in intervals I_i of size $\delta/2$. And denote

$$m = \sup_i \inf_{I_i} \varphi(t_0, \cdot).$$

By the previous property $m < 0$. Define $\tau = -m/(3\|\varphi\|_{W^{1,\infty}})$.

Now for any x_0 at distance larger than δ from $\{\varphi_\varepsilon(t_0, \cdot) \geq -\bar{\varphi}_\varepsilon\}$, there exists $y_1 < x_0$ and $y_2 > x_0$ s.t.

$$|y_i - x_0| \geq \delta/2, \quad \varphi(t_0, y_i) \leq m.$$

Obviously for any $t < t_0 + \tau$,

$$\varphi(t, y_i) \leq 2m/3,$$

and hence for any y s.t. $|y - y_i| \leq \tau$ then

$$\varphi(t, y) \leq m/3, \quad \varphi_\varepsilon(t, y) \leq m/3 + \|\varphi - \varphi_\varepsilon\|_{L^\infty} < -\bar{\varphi}_\varepsilon, \quad (49)$$

provided ε is chosen small enough.

Now we apply a first time Lemma 4.0.8 with $\delta = \tau$ which means that for ε small enough with respect to τ , there cannot be a jump of size τ in $\{\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$. If for some $t \leq t_0 + \tau$

$$[y_1, y_2] \cap \{\varphi_\varepsilon(t, \cdot) \geq -\bar{\varphi}_\varepsilon\} \neq \emptyset,$$

then by Lemma 4.0.8 with $\delta = \tau$, it means that for some $s \in [t_0, t]$, one can find a point y with $|y - y_1| \leq \tau$ or $|y - y_2| \leq \tau$ s.t.

$$\varphi_\varepsilon(s, y) \geq -\bar{\varphi}_\varepsilon,$$

which contradicts (49).

Hence we know that x_0 stays at distance $\delta/2$ of the set $\{\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$. We apply once more Lemma 4.0.8 to conclude. \square

4.6. Identifying the limit terms: Final steps. The first step is to characterize the limit of D_ε . Of course there is a natural candidate for that which is

$$D = \min(Ld(x, \{\varphi = 0\}), D_0). \quad (50)$$

By (43), we essentially know that at most times t , the set $\{\varphi_\varepsilon(t, \cdot) \geq -\bar{\varphi}_\varepsilon\}$ is included in a neighborhood of $\{\varphi(t, \cdot) = 0\}$. However to prove (50), we need to show the contrary namely that $\{\varphi(t, \cdot) = 0\}$ is included in a neighborhood of $\{\varphi_\varepsilon(t, \cdot) \geq -\bar{\varphi}_\varepsilon\}$ at most times t . This is what the following lemma shows

Lemma 4.0.10. *For any fixed $\delta > 0$, denote*

$$J_\varepsilon = \{t \in [0, T], \quad \delta(\{\varphi(t, \cdot) = 0\}, \{\varphi_\varepsilon(t, \cdot) \geq -\bar{\varphi}_\varepsilon\}) > \delta\}.$$

Then $J_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ and consequently (50) holds.

The last step is to identify the weak limit u of u_ε . Fortunately we now have all the required tools and we can follow the same ideas as in [11]. Therefore let us define

$$\mu_t = \mu(t) = \mu(\{\varphi(t, \cdot) = 0\}),$$

where μ is given by Prop. 1.

We prove the intermediary result

Lemma 4.0.11. *For any fixed $t_0 \in [0, T] \setminus \bar{I}_\varepsilon$, there exists a function $\sigma \in C(\mathbb{R}_+)$ with $\sigma(0) = 0$ s.t.*

$$\int_{t_0}^t \int_{\mathbb{R}} |I \star u_\varepsilon(s, x) - I \star \mu(t_0, x)|^2 dx ds \leq C(t - t_0) \tau(t - t_0) + \sigma(\varepsilon),$$

where τ is given by (37).

Then simply by passing to the limit one deduces that

Corollary 2. *For any fixed t_0 ,*

$$\int_{t_0}^t \int_{\mathbb{R}} |I \star u(s, x) - I \star \mu(t_0, x)|^2 dx ds \leq (t - t_0) \tau(t - t_0).$$

This means that any t_0 is a Lebesgue point of $I \star u$ and then necessarily that

$$I \star u(t_0, \cdot) = I \star \mu(t_0, \cdot),$$

which as $\hat{I} > 0$ lets us conclude that $u = \mu$. We have identified all the terms in the limiting equation. It remains to actually obtain this equation and explain in which sense, which is done in the next subsection. We conclude this subsection by giving the proofs of Lemmas 4.0.10 and 4.0.11.

Proof of Lemma 4.0.10. Choose a time $t_0 \in [0, T] \setminus \bar{I}_\varepsilon$. Apply Corollary 1 and Lemma 4.0.9 and find t s.t. $\tau_{t_0}(t - t_0) \leq \delta/2$ and the τ from Lemma 4.0.9 satisfies $t < t_0 + \tau$.

Then for any $s \in [t_0, t]$ and any x_0 s.t.

$$d(x_0, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) > \delta.$$

First by Corollary 1,

$$d(x_0, \{\varphi_\varepsilon(t_0, \cdot) \geq -\bar{\varphi}_\varepsilon\}) > \delta/2.$$

One has then by Lemma 4.0.9 that on $[t_0, t]$,

$$\varphi_\varepsilon(s, x_0) < -\bar{\varphi}_\varepsilon - C\delta(s - t_0).$$

That implies that on $s \in [t_0 + C\delta^{-1}\|\varphi - \varphi_\varepsilon\|_{L^\infty}, t]$ then

$$x_0 \notin \{\varphi(s, \cdot) = 0\}.$$

Therefore for any $s \in [t_0 + C\delta^{-1}\|\varphi - \varphi_\varepsilon\|_{L^\infty}, t]$

$$\delta(\{\varphi(s, \cdot) = 0\}, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) \leq \delta.$$

We conclude that for any $t_0 \notin \bar{I}_\varepsilon$ there exists $t > 0$ chosen uniformly in ε s.t.

$$J_\varepsilon \cap [t_0, t] \subset [t_0, t_0 + C\delta^{-1}\|\varphi - \varphi_\varepsilon\|_{L^\infty}].$$

which implies that $|J_\varepsilon| \rightarrow 0$. □

Proof of Lemma 4.0.11. The steps are mostly the same as in [11]. We give them for the sake of completeness but we do not repeat all the details here. Part of the proof can also be simplified by using the more precise results that were proved before in the present article.

Step 1: The functional.

We look at the evolution of

$$F_\varepsilon(t) = \int_{\mathbb{R}} \log u_\varepsilon(t, x) d\mu_{t_0}(x) = \frac{1}{\varepsilon} \int_{\mathbb{R}} \varphi_\varepsilon(t, x) d\mu_{t_0}(x),$$

for $s \geq t_0$. Compute

$$\begin{aligned} \frac{d}{dt} F_\varepsilon(t) &= \frac{1}{\varepsilon} \int_{\mathbb{R}} (r - I \star u_\varepsilon - D_\varepsilon) d\mu_{t_0}(x) \\ &\quad + \frac{1}{\varepsilon} \int_{\mathbb{R}} H_\varepsilon(\varphi_\varepsilon(t)) u_\varepsilon d\mu_{t_0}. \end{aligned}$$

Write

$$\begin{aligned} \frac{1}{\varepsilon} \int_{\mathbb{R}} (r - I \star u_\varepsilon - D_\varepsilon) d\mu_{t_0}(x) &= \frac{d}{dt} \int_{\mathbb{R}} u_\varepsilon(t, x) dx \\ &\quad - \frac{1}{\varepsilon} \int_{\mathbb{R}} (r - I \star u_\varepsilon - D_\varepsilon) (u_\varepsilon(t, x) dx - d\mu_{t_0}(x)). \end{aligned}$$

As $r - I \star \mu_{t_0}$ vanishes on the support of μ_{t_0} ,

$$\begin{aligned} \frac{1}{\varepsilon} \int_{\mathbb{R}} (r - I \star u_\varepsilon - D_\varepsilon) d\mu_{t_0}(x) &= \frac{d}{dt} \int_{\mathbb{R}} u_\varepsilon(t, x) - \frac{A(t)}{\varepsilon} \\ &\quad + \frac{1}{\varepsilon} \int_{\mathbb{R}} I \star (u_\varepsilon - \mu_{t_0}) (u_\varepsilon(t, x) dx - d\mu_{t_0}(x)) + \frac{1}{\varepsilon} \int_{\mathbb{R}} D_\varepsilon u_\varepsilon dx, \end{aligned}$$

with

$$A(t) = \int_{\mathbb{R}} (r - I \star \mu_s) u_\varepsilon(t, x) dx + \int_{\mathbb{R}} D_\varepsilon d\mu_{t_0}.$$

Notice that

$$\int_{\mathbb{R}} I \star (u_\varepsilon(t, \cdot) - \mu_{t_0}) (u_\varepsilon(t, x) dx - d\mu_s(x))$$

controls $\|I \star (u_\varepsilon(t, \cdot) - \mu_{t_0})\|_{L^2}^2$ since $\hat{I}^2 \leq C\hat{I}$.

So we deduce since $D_\varepsilon \geq 0$

$$\begin{aligned} \frac{1}{\varepsilon} \int_{t_0}^t \|I \star (u_\varepsilon(s, \cdot) - \mu_{t_0})\|_{L^2}^2 ds &\leq \int_{\mathbb{R}} \log \frac{u_\varepsilon(t, x)}{u_\varepsilon(t_0, x)} d\mu_{t_0} \\ &\quad - \int_{\mathbb{R}} (u_\varepsilon(t, x) - u_\varepsilon(t_0, x)) dx + \int_{t_0}^t \frac{A(s)}{\varepsilon} ds - \frac{1}{\varepsilon} \int_{t_0}^t \int_{\mathbb{R}} H_\varepsilon(\varphi_\varepsilon(t_0)) d\mu_{t_0}. \end{aligned} \tag{51}$$

Step 3: Easy bounds.

Lemma 4.0.1 tells that

$$-H_\varepsilon(\varphi_\varepsilon) \leq C \frac{\varepsilon}{\varphi_\varepsilon}.$$

The total mass stays bounded in time so

$$- \int_{\mathbb{R}} (u_\varepsilon(t, x) - u_\varepsilon(t_0, x)) dx \leq \int_{\mathbb{R}} (u_\varepsilon(t, x) + u_\varepsilon(t_0, x)) dx \leq C.$$

And

$$\begin{aligned} \int_{\mathbb{R}} \log \frac{u_\varepsilon(t, x)}{u_\varepsilon(t_0, x)} d\mu_{t_0} &= \frac{1}{\varepsilon} \int_{\mathbb{R}} (\varphi_\varepsilon(t, x) - \varphi_\varepsilon(t_0, x)) d\mu_{t_0} \\ &\leq \frac{1}{\varepsilon} \int_{\mathbb{R}} (\varphi(t, x) - \varphi(t_0, x)) d\mu_{t_0} + \frac{2}{\varepsilon} \|\varphi_\varepsilon - \varphi\|_{L^\infty(\Omega)}. \end{aligned}$$

Note that by Prop. 1, μ_{t_0} is supported on $\{\varphi(s, \cdot) = 0\}$. Since in addition we know that $\varphi \leq 0$,

$$\int_{\mathbb{R}} \log \frac{u_\varepsilon(t, x)}{u_\varepsilon(t_0, x)} d\mu_{t_0} \leq \frac{2}{\varepsilon} \|\varphi_\varepsilon - \varphi\|_{L^\infty(\Omega)}.$$

Consequently we deduce from (51) the bound

$$\frac{1}{\varepsilon} \int_{t_0}^t \|I \star (u_\varepsilon(s, \cdot) - \mu_{t_0})\|_{L^2}^2 ds \leq \frac{C}{\bar{\varphi}_\varepsilon} + \frac{2}{\varepsilon} \|\varphi_\varepsilon - \varphi\|_{L^\infty(\Omega)} + \int_{t_0}^t \frac{A(s)}{\varepsilon} ds. \quad (52)$$

Step 4: Control on A.

The key point here is that $\text{supp } \mu_{t_0} = \{\varphi(t_0, \cdot) = 0\}$ is very close to $\{\varphi_\varepsilon \geq -\bar{\varphi}_\varepsilon\}$ as implied by Lemma 4.0.10.

First of all $D_\varepsilon(s, \cdot)$ is Lipschitz in x and vanishes on $\{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}$, so one has that

$$\int_{t_0}^t \int_{\mathbb{R}} D_\varepsilon(s, x) d\mu_{t_0} ds \leq C \int_{t_0}^t \delta(\text{supp } \mu_{t_0}, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) ds.$$

By the definition of μ , its support is included in $\{\varphi = 0\}$ and by Lemma 4.0.2 we finally deduce that

$$\begin{aligned} \int_{t_0}^t \int_{\mathbb{R}} D_\varepsilon(s, x) d\mu_{t_0} ds &\leq C \tau(t - t_0) (t - t_0) \\ &+ C \int_{t_0}^t \delta(\{\varphi(s, \cdot) = 0\}, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) ds. \end{aligned} \quad (53)$$

This was the only additional term with respect to [11]. The other term is treated the same by decomposing

$$\begin{aligned} \int_{t_0}^t \int_{\mathbb{R}} (r - I \star \mu_{t_0}) u_\varepsilon(s, x) dx ds &= \int_{t_0}^t \int_{\mathbb{R}} (r - I \star \mu_{t_0}) u_\varepsilon(s, x) \mathbb{I}_{\varphi_\varepsilon(s, x) \leq -\bar{\varphi}_\varepsilon} dx ds \\ &+ \int_{t_0}^t \int_{\mathbb{R}} (r - I \star \mu_{t_0}) u_\varepsilon(s, x) \mathbb{I}_{\varphi_\varepsilon(s, x) \geq -\bar{\varphi}_\varepsilon} dx ds. \end{aligned}$$

For the first part, note again that by (18), there exists R s.t. $\forall |x| > R, r - I \star \mu_{t_0} < 0$.

Therefore, we may simply dominate

$$\int_{t_0}^t \int_{\mathbb{R}} (r - I \star \mu_{t_0}) u_\varepsilon(s, x) \mathbb{I}_{\varphi_\varepsilon(s, x) \leq -\bar{\varphi}_\varepsilon} dx ds \leq C (t - s) e^{-\bar{\varphi}_\varepsilon/\varepsilon}.$$

Concerning the second part, just note that $\{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\} \subset \{\varphi(s, \cdot) \geq -\alpha_\varepsilon\}$ with $\alpha_\varepsilon = \bar{\varphi}_\varepsilon + \|\varphi - \varphi_\varepsilon\|_{L^\infty}$ and that $r - I \star \mu_{t_0}$ is non positive on $\{\varphi(t_0, \cdot) = 0\}$. Hence

$$\begin{aligned} \int_{t_0}^t \int_{\mathbb{R}} (r - I \star \mu_{t_0}) u_\varepsilon(s, x) \mathbb{I}_{\varphi_\varepsilon(s, x) \geq -\bar{\varphi}_\varepsilon} dx ds \\ \leq C \int_{t_0}^t \delta(\{\varphi(s, \cdot) \geq -\alpha_\varepsilon\}, \{\varphi(t_0, \cdot) = 0\}) ds. \end{aligned}$$

By Lemma 4.0.8, this implies that

$$\begin{aligned} \int_{t_0}^t \int_{\mathbb{R}} (r - I \star \mu_{t_0}) u_\varepsilon(s, x) \mathbb{I}_{\varphi_\varepsilon(s, x) \geq -\bar{\varphi}_\varepsilon} dx ds &\leq \tau(t - t_0) (t - t_0) \\ &+ C \int_{t_0}^t \delta(\{\varphi(s, \cdot) \geq -\alpha_\varepsilon\}, \{\varphi(s, \cdot) = 0\}) ds. \end{aligned}$$

Inequality (52) now becomes

$$\begin{aligned} \int_{t_0}^t \|I \star (u_\varepsilon(s, \cdot) - \mu_{t_0})\|_{L^2}^2 ds &\leq \frac{C\varepsilon}{\bar{\varphi}_\varepsilon} + 2\|\varphi_\varepsilon - \varphi\|_{L^\infty(\Omega)} + C(t-s)e^{-\bar{\varphi}_\varepsilon/\varepsilon} \\ &+ C\tau(t-t_0)(t-t_0) + C \int_{t_0}^t \delta(\{\varphi(s, \cdot) \geq -\alpha_\varepsilon\}, \{\varphi(s, \cdot) = 0\}) ds \\ &+ C \int_{t_0}^t \delta(\{\varphi(s, \cdot) = 0\}, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) ds. \end{aligned} \tag{54}$$

Conclusion. Eq. (51) indeed gives Lemma 4.0.11 if one defines

$$\begin{aligned} \tilde{\sigma}(\varepsilon) &= \frac{C\varepsilon}{\bar{\varphi}_\varepsilon} + 2\|\varphi_\varepsilon - \varphi\|_{L^\infty(\Omega)} + CT e^{-\bar{\varphi}_\varepsilon/\varepsilon} \\ &+ C \int_0^T \delta(\{\varphi(s, \cdot) \geq -\alpha_\varepsilon\}, \{\varphi(s, \cdot) = 0\}) ds \\ &+ C \int_0^T \delta(\{\varphi(s, \cdot) = 0\}, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) ds. \end{aligned}$$

Thus, in order to complete the proof of Lemma 4.0.11, we only have to check that $\tilde{\sigma}(\varepsilon) \rightarrow 0$ when $\varepsilon \rightarrow 0$.

It is obvious for the first three terms. For the second note that

$$C \int_0^T \delta(\{\varphi(s, \cdot) \geq -\alpha_\varepsilon\}, \{\varphi(s, \cdot) = 0\}) ds \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0,$$

since by dominated convergence it is enough that for any s

$$\delta(\{\varphi(s, \cdot) \geq -\alpha_\varepsilon\}, \{\varphi(s, \cdot) = 0\}) \rightarrow 0.$$

For the last term, we use Lemma 4.0.10 to show that it converges to 0 as $\varepsilon \rightarrow 0$. Indeed for any δ one has

$$\int_0^T \delta(\{\varphi(s, \cdot) = 0\}, \{\varphi_\varepsilon(s, \cdot) \geq -\bar{\varphi}_\varepsilon\}) ds \leq |J_\varepsilon| + T\delta,$$

with $|J_\varepsilon| \rightarrow 0$. □

4.7. Obtaining the limit equation in the sense of viscosity solutions. This is the final step in the proof of Theorem 2.1, which we postponed until now. With the many properties that we have been able to show, we are now able to obtain φ as a true viscosity solution (and not in the modified sense of [20]) to

$$\partial_t \varphi = r - I \star u(t, \cdot) - D + H(\partial_x \phi). \tag{55}$$

First of all let us summarize the important properties on D and $I \star u$ that we need

Lemma 4.0.12. *The function D is continuous in x and left-continuous in time. The function $I \star u$ is continuous on $[0, T] \times \mathbb{R}$. Moreover as $\varepsilon \rightarrow 0$*

$$\int_0^T (\|D(\varphi(t, \cdot)) - D_\varepsilon(\varphi_\varepsilon(t, \cdot))\|_{L^\infty} + \|I \star u(t, \cdot) - I \star u_\varepsilon(t, \cdot)\|_{L^\infty}) dt \rightarrow 0.$$

Proof of Lemma 4.0.12. It is simple and uses previous estimates. First of all by Lemma 4.0.2 and property (37), $\{\varphi(t, \cdot) = 0\}$ is continuous on the left in time. The formula (50) for D then immediately implies the continuity of D in x and on the left in t . Notice in addition that any jump on D in time is decreasing.

We have seen that $I \star \mu(t, \cdot) = I \star u(t, \cdot)$. As I is smooth then $I \star u$ is of course continuous (and even more) in x . By the stability estimates in [26] for instance, we know that the equilibrium measure μ defined by Prop. 1 is continuous in the set Ω . As again $\{\varphi(t, \cdot) = 0\}$ is continuous on the left in time, $\mu(t, \cdot)$ is continuous on the left in time with value in M_+^1 for some weak topology of measures.

From the continuity on the left and Corollary 2, we finally deduce that $I \star \mu$ is continuous on both sides in time.

Let us now turn to the convergence estimates, first for $D_\varepsilon(\phi_\varepsilon)$. For any η , there exists $\bar{\delta}$ s.t. if

$$\begin{aligned} \delta(\{\varphi(t, \cdot) = 0\}, \{\varphi_\varepsilon(t, \cdot) \geq -\bar{\varphi}_\varepsilon\}) &< \delta, \\ \delta(\{\varphi_\varepsilon(t, \cdot) \geq -\bar{\varphi}_\varepsilon\}, \{\varphi(t, \cdot) = 0\}) &< \delta, \end{aligned}$$

then $|D_\varepsilon(\varphi_\varepsilon) - D(\varphi)| < \eta$. By Lemma 4.0.10 the first is true for any time $t \in [0, T] \setminus J_\varepsilon$. By estimate (43), the second is true for ε small enough and for any time $t \in [0, T] \setminus I_\varepsilon$. Therefore

$$\int_0^T \|D_\varepsilon(\varphi_\varepsilon(t, \cdot)) - D(\varphi(t, \cdot))\|_{L^\infty} dt \leq T \bar{\delta} + |J_\varepsilon| + |I_\varepsilon|.$$

As $|I_\varepsilon|$ and $|J_\varepsilon|$ converge to 0 with ε , this proves the convergence on D_ε .

Concerning $I \star u_\varepsilon$, Lemma 4.0.11 implies that $I \star u_\varepsilon$ converges strongly to $I \star u$ in $L^2([0, T] \times \mathbb{R})$ (see [11] for the complete argument for instance). As I is smooth, this then implies that $I \star u_\varepsilon$ converges strongly to $I \star u$ in $L^2([0, T], L^\infty(\mathbb{R}))$ and hence *a fortiori* in $L_t^1(L_x^\infty)$. This concludes the proof of the lemma. \square

With the estimates of Lemma 4.0.12, it is now straightforward to show that φ satisfies (55) in the viscosity sense and we only sketch the main steps. For instance, one may define $\tilde{\varphi}_\varepsilon$ as the solution to

$$\partial_t \tilde{\varphi}_\varepsilon = r - I \star u - D(\varphi) + H_\varepsilon(\tilde{\varphi}_\varepsilon). \quad (56)$$

Note that in this equation we have taken the limit for $I \star u$ and D but not on the hamiltonian. Applying the maximum principle, one deduces that

$$\begin{aligned} \frac{d}{dt} \|\tilde{\varphi}_\varepsilon(t, \cdot) - \varphi_\varepsilon(t, \cdot)\|_{L^\infty} &\leq \|I \star u_\varepsilon(t, \cdot) - I \star u(t, \cdot)\|_{L^\infty} \\ &\quad + \|D(\varphi(t, \cdot)) - D_\varepsilon(\varphi_\varepsilon(t, \cdot))\|_{L^\infty}. \end{aligned}$$

So by Lemma 4.0.12, $\tilde{\varphi}_\varepsilon$ also converges uniformly to φ . The standard theory of viscosity solutions may now be applied (see [1], [2] or [14]) since the source terms in the righthand side of (56) are now continuous (or semi-continuous for D). We conclude that $\tilde{\varphi}_\varepsilon$ converges to a viscosity solution of (55) and since the limit of $\tilde{\varphi}_\varepsilon$ is φ that φ is this viscosity solution.

4.8. Concluding remarks: Finite speed of propagation of the support.

It is now easy to obtain additional properties for the limit φ , for instance the finite speed of propagation of the support. We just sketch quickly here which properties one should be able to obtain without giving proofs.

First by the definition of μ , one has that

$$r - I \star \mu(\{\varphi(t, \cdot)\}) \leq 0, \quad \text{on } \{\varphi(t, \cdot) = 0\}.$$

One then deduces immediately that for any x

$$r(x) - I \star \mu(\{\varphi(t, \cdot)\}) - D(t, x) \leq 0,$$

since $D = \min(Ld(x, \{\varphi(t, \cdot) = 0\}), D_0)$. That means by Eq. (55)

$$\partial_t \varphi \leq H(\partial_x \varphi).$$

Just by the following the characteristics of the Hamilton-Jacobi equation, one obtains that the set $\{\varphi = 0\}$ can only propagate at finite speed V .

We mentioned in the introduction that, for the non corrected model, branching occurs at infinite speed. Therefore one may wonder whether branching is possible in the corrected model. First of all numerical simulations suggest without doubt that the correction do not prevent branching.

Second the formal analysis in [20] that showed the infinite speed at branching cannot be carried anymore with the correction. This analysis required the solution to be C^2 near the branching point but the correction actually ensures that this is no more possible. Indeed as soon as branching occurs and the set $\{\varphi(t, \cdot) = 0\}$ includes at least two points $x_1(t), x_2(t)$ then the function $d(x, \{\varphi(t, \cdot) = 0\})$ necessarily has a singularity between x_1 and x_2 .

Of course a conclusive argument would be to exhibit an explicit solution that branches. This is already unavailable for the uncorrected model and hence even more so here.

Finally let us mention that some time regularity should also be available on μ . For instance the set $\{\varphi = 0\}$ is continuous in time by Lemma 4.0.2 and we have already seen how to deduce from it that $\mu(\{\varphi(t, \cdot)\})$ is continuous in time. But as a matter of fact the finite speed of propagation of $\{\varphi = 0\}$ should even imply Hölder continuity in time for μ . This is certainly more delicate and it might require the reworking of the stability estimates in [26].

REFERENCES

- [1] M. Bardi and I. Capuzzo Dolcetta, “Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations”.
- [2] G. Barles, “Solutions de Viscosité et Équations de Hamilton-Jacobi,” Collec. SMAI, Springer-Verlag, 2002.
- [3] G. Barles, S. Mirrahimi and B. Perthame, *Concentration in Lotka-Volterra parabolic or integral equations: a general convergence result*, Methods Appl. Anal., **16** (2009), 321–340.
- [4] G. Barles and B. Perthame, *Concentrations and constrained Hamilton-Jacobi equations arising in adaptive dynamics*, Recent Developments in Nonlinear Partial Differential Equations, 57–68, Contemp. Math., **439**, Amer. Math. Soc., Providence, RI, (2007).
- [5] R. Bürger and I. M. Bomze, *Stationary distributions under mutation-selection balance: structure and properties*, Adv. Appl. Prob., **28** (1996), 227–251.
- [6] A. Calsina and S. Cuadrado, *Small mutation rate and evolutionarily stable strategies in infinite dimensional adaptive dynamics*, J. Math. Biol., **48** (2004), 135–159.
- [7] J. A. Carrillo, S. Cuadrado and B. Perthame, *Adaptive dynamics via Hamilton-Jacobi approach and entropy methods for a juvenile-adult model*, Math. Biosci., **205** (2007), 137–161.
- [8] N. Champagnat, *A microscopic interpretation for adaptive dynamics trait substitution sequence models*, Stoch. Proc. Appl., **116** (2006), 1127–1160.
- [9] N. Champagnat, R. Ferrière and G. Ben Arous, *The canonical equation of adaptive dynamics: A mathematical view*, Selection, **2** (2001), 71–81.
- [10] N. Champagnat, R. Ferrière and S. Méléard, *From individual stochastic processes to macroscopic models in adaptive evolution*, Stoch. Models, **24** (2008), 2–44.
- [11] N. Champagnat and P.-E. Jabin, *The evolutionary limit for models of populations interacting competitively via several resources*, J. Differential Equations, **251** (2011), 176–195.
- [12] N. Champagnat, P.-E. Jabin and G. Raoul, *Convergence to equilibrium in competitive Lotka-Volterra and chemostat systems*, C. R. Math. Acad. Sci. Paris, **348** (2010), 1267–1272.
- [13] N. Champagnat and S. Méléard, *Polymorphic evolution sequence and evolutionary branching*, To appear in Probab. Theory Relat. Fields (published online, 2010).

- [14] M. G. Crandall and P.-L. Lions, *Users guide to viscosity solutions of second order partial differential equations*, Bull. Amer. Math. Soc., **27** (1992), 167.
- [15] R. Cressman and J. Hofbauer, *Measure dynamics on a one-dimensional continuous trait space: theoretical foundations for adaptive dynamics*, Theor. Pop. Biol., **67** (2005), 47–59.
- [16] L. Desvillettes, P.-E. Jabin, S. Mischler and G. Raoul, *On selection dynamics for continuous structured populations*, Commun. Math. Sci., **6** (2008), 729–747.
- [17] U. Dieckmann and R. Law, *The dynamical theory of coevolution: A derivation from stochastic ecological processes*, J. Math. Biol., **34** (1996), 579–612.
- [18] O. Diekmann, *A beginner's guide to adaptive dynamics. In Mathematical modelling of population dynamics*, Banach Center Publ., **63**, 47–86, Polish Acad. Sci., Warsaw, (2004).
- [19] O. Diekmann, M. Gyllenberg, H. Huang, M. Kirkilionis, J. A. J. Metz and H. R. Thieme, *On the formulation and analysis of general deterministic structured population models. II. Nonlinear theory*, J. Math. Biol., **43** (2001), 157–189.
- [20] O. Diekmann, P. E. Jabin, S. Mischler and B. Perthame, *The dynamics of adaptation: An illuminating example and a Hamilton-Jacobi approach*, Theor. Popul. Biol., **67** (2005), 257–271.
- [21] S. Génieys, N. Bessonov and V. Volpert, *Mathematical model of evolutionary branching*, Math. Comput. Modelling, **49** (2009), 2109–2115.
- [22] S. A. H. Geritz, J. A. J. Metz, E. Kisdi and G. Meszéna, *Dynamics of adaptation and evolutionary branching*, Phys. Rev. Lett., **78** (1997), 2024–2027.
- [23] S. A. H. Geritz, E. Kisdi, G. Meszéna and J. A. J. Metz, *Evolutionary singular strategies and the adaptive growth and branching of the evolutionary tree*, Evol. Ecol., **12** (1998), 35–57.
- [24] M. Gyllenberg and G. Meszéna, *On the impossibility of coexistence of infinitely many strategies*, J. Math. Biol., **50** (2005), 133–160.
- [25] J. Hofbauer and R. Sigmund, *Adaptive dynamics and evolutionary stability*, Applied Math. Letters, **3** (1990), 75–79.
- [26] P. E. Jabin and G. Raoul, *Selection dynamics with competition*, J. Math. Biol., **63** (2011), 493–517.
- [27] A. Lorz, S. Mirrahimi and B. Perthame, *Dirac mass dynamics in multidimensional nonlocal parabolic equations*, Comm. Partial Differential Equations, **36** (2011), 1071–1098.
- [28] S. Méléard, *Introduction to stochastic models for evolution*, Markov Process. Related Fields, **15** (2009), 259–264.
- [29] S. Méléard and V. C. Tran, *Trait substitution sequence process and canonical equation for age-structured populations*, J. Math. Biol., **58** (2009), 881–921.
- [30] J. A. J. Metz, R. M. Nisbet and S. A. H. Geritz, *How should we define 'fitness' for general ecological scenarios?*, Trends in Ecology and Evolution, **7** (1992), 198–202.
- [31] J. A. J. Metz, S. A. H. Geritz, G. Meszéna, F. A. J. Jacobs and J. S. van Heerwaarden, *Adaptive Dynamics, a geometrical study of the consequences of nearly faithful reproduction*, in “Stochastic and Spatial Structures of Dynamical Systems” (eds. S. J. van Strien & S. M. Verduyn Lunel), North Holland, Amsterdam, (1996), 183–231.
- [32] S. Mirrahimi, G. Barles, B. Perthame and P. E. Souganidis, *Singular Hamilton-Jacobi equation for the tail problem*, Submitted.
- [33] B. Perthame and M. Gauduchon, *Survival thresholds and mortality rates in adaptive dynamics: Conciliating deterministic and stochastic simulations*, IMA Journal of Mathematical Medicine and Biology, to appear (published online, 2009).
- [34] B. Perthame and S. Génieys, *Concentration in the nonlocal Fisher equation: the Hamilton-Jacobi limit*, Math. Model. Nat. Phenom., **2** (2007), 135–151.
- [35] G. Raoul, *Long time evolution of populations under selection and rare mutations*, Acta Applicandae Mathematica, **114** (2011), 114.
- [36] F. Yu, *Stationary distributions of a model of sympatric speciation*, Ann. Appl. Probab., **17** (2007), 840–874.

Received March 2012; revised August 2012.

E-mail address: pjabin@cscamm.umd.edu