

STABILITY ESTIMATES FOR SCALAR CONSERVATION LAWS WITH MOVING FLUX CONSTRAINTS

MARIA LAURA DELLE MONACHE

Department of mathematical sciences
Rutgers University - Camden
311 N. 5th Street, Camden, NJ 08102, USA
and

Inria, Univ. Grenoble Alpes, CNRS, GIPSA-lab
F-38000 Grenoble, France

PAOLA GOATIN*

Inria Sophia Antipolis - Méditerranée
Université Côte d’Azur, Inria, CNRS, LJAD
2004, route des Lucioles - BP 93, 06902 Sophia Antipolis Cedex, France

ABSTRACT. We study well-posedness of scalar conservation laws with moving flux constraints. In particular, we show the Lipschitz continuous dependence of BV solutions with respect to the initial data and the constraint trajectory. Applications to traffic flow theory are detailed.

1. Introduction. Motivated by the modeling of moving bottlenecks in traffic flow, which can be caused by a large, slow moving vehicle, we consider the Cauchy problem for a scalar conservation law with moving flux constraint

$$\partial_t \rho + \partial_x f(\rho) = 0, \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \quad (1a)$$

$$\rho(0, x) = \rho_0(x), \quad x \in \mathbb{R}, \quad (1b)$$

$$f(\rho(t, y(t))) - \dot{y}(t)\rho(t, y(t)) \leq \tilde{F}_\alpha(\dot{y}(t)), \quad t \in \mathbb{R}^+, \quad (1c)$$

where $t \mapsto y(t)$ is a given trajectory, starting from $y(0) = y_0$, and

$$\tilde{F}_\alpha(\dot{y}) := F_\alpha(\rho_\alpha(\dot{y})) = f_\alpha(\rho_\alpha(\dot{y})) - \dot{y}\rho_\alpha(\dot{y}), \quad (2)$$

with $\rho_\alpha(\dot{y})$ such that $f'(\rho_\alpha(\dot{y})/\alpha) = \dot{y}$. Systems of the form (1) arise in the modeling of moving bottlenecks in vehicular traffic [12, 16]: $\rho = \rho(t, x) \in [0, R]$ is the scalar conserved quantity and represents the traffic density, whose maximum attainable value is R . The flux function $f : [0, R] \rightarrow \mathbb{R}^+$ is assumed to be strictly concave, Lipschitz continuous and such that $f(0) = f(R) = 0$. The time-dependent variable y denotes the constraint position. In the present paper we consider a weakly coupled PDE-ODE system, in the sense that we assume that the constraint trajectory is given, and it does not depend on the solution of (1a).

2010 *Mathematics Subject Classification.* Primary: 35L65; Secondary: 35B35.

Key words and phrases. Scalar conservation laws, moving flux constraints, PDE-ODE system.

This research was supported by the Inria Associated Team “Optimal REroute Strategies for Traffic managEment” (ORESTE).

* Corresponding author: Paola Goatin.

Let us detail the meaning of inequality (1c). A moving flux constraint located at $x = y(t)$ acts as an obstacle, thus hindering the flow as expressed by the unilateral constraint (1c). There, $\alpha \in]0, 1[$ is the dimensionless reduction rate of the road capacity (the maximal allowed density) at the bottleneck position. The inequality (1c) is derived by studying the problem in the constraint reference frame, i.e., setting $\tilde{\rho}(t, x) := \rho(t, x + y(t))$ and rewriting the conservation law (1a) as

$$\partial_t \tilde{\rho} + \partial_x F(\tilde{\rho}) = 0, \quad F(\tilde{\rho}) = f(\tilde{\rho}) - \dot{y}\tilde{\rho}. \quad (3)$$

In fact, let $f_\alpha : [0, \alpha R] \rightarrow \mathbb{R}^+$ be the rescaled flux function describing the reduced flow at $x = y(t)$, defined by $f_\alpha(\rho) = \alpha f(\rho/\alpha)$, and let $\rho_\alpha(\dot{y}) \in]0, \alpha R[$ such that

$$F'_\alpha(\rho_\alpha(\dot{y})) = f'_\alpha(\rho_\alpha(\dot{y})) - \dot{y} = 0 \quad \Leftrightarrow \quad f'_\alpha(\rho_\alpha(\dot{y})) = \dot{y}, \quad (4)$$

with $F_\alpha(\rho) = f_\alpha(\rho) - \dot{y}\rho$, see Figure 1. Notice that, for (4) to have a solution in $]0, \alpha R[$, we will need to assume that $\dot{y}(t) < f'(0)$ for $t \geq 0$. Then setting $\tilde{F}_\alpha(\dot{y}) := F_\alpha(\rho_\alpha(\dot{y}))$ we recover (2). Imposing that in the obstacle reference frame the flux F is less than the maximum value of the flux of the reduced flow, one gets (1c). Notice that the inequality (1c) is always satisfied if

$$\dot{y}(t) = \frac{f(\rho(t, y(t) \pm))}{\rho(t, y(t) \pm)},$$

since its left-hand side is 0. Moreover, it is well defined even if ρ has a jump at $y(t)$, because of the Rankine-Hugoniot conditions.




FIGURE 1. Graphical representation of the constraint action in the fixed (left) and moving (right) reference frames.

Problem (3), (1b), (1c), can therefore be recast in the framework of conservation laws with fix local constraint, first introduced in [10], then developed in [2, 4] for scalar equations and extended in [3, 14, 13] to systems. Following [12, Definition 4.1] and [6, Definition 1 and 2], solutions of (1) are defined as follows.

Definition 1.1. Let $y \in \mathbf{W}^{1,\infty}(\mathbb{R}^+; \mathbb{R})$ with $0 \leq \dot{y} < f'(0)$ and $\rho_0 \in \mathbf{L}^1 \cap \mathbf{L}^\infty(\mathbb{R}; [0, R])$ be given. A function $\rho \in \mathcal{C}^0(\mathbb{R}^+; \mathbf{L}^1(\mathbb{R}; [0, R]))$ is a solution to (1) if

1. ρ satisfies Kruřkov entropy conditions [15] on $(\mathbb{R}^+ \times \mathbb{R}) \setminus \{(t, y(t)) : t \in \mathbb{R}^+\}$, i.e. for every $k \in [0, R]$ and for all $\varphi \in \mathcal{C}_c^1(\mathbb{R}^2; \mathbb{R}^+)$ and $\varphi(t, y(t)) = 0, t > 0$,

$$\int_{\mathbb{R}^+} \int_{\mathbb{R}} (|\rho - k| \partial_t \varphi + \text{sgn}(\rho - k) (f(\rho) - f(k)) \partial_x \varphi) dx dt + \int_{\mathbb{R}} |\rho_0 - k| \varphi(0, x) dx \geq 0 ; \tag{5a}$$

2. for a. e. $t > 0$ the left and right traces of ρ at $x = y(t)$ satisfy

$$(\rho(t, y(t)-), \rho(t, y(t)+)) \in \mathcal{G}_\alpha(\dot{y}(t)). \tag{5b}$$

Notice that the left and right traces in (5b) do exist, see [4, Section 2].

The set $\mathcal{G}_\alpha(\dot{y})$ in (5b) is defined as follows, see [4, 5, 6].

Definition 1.2. The *admissibility germ* $\mathcal{G}_\alpha(\dot{y}) \subset [0, R]^2$ for (1a), (1c) is the union $\mathcal{G}_\alpha(\dot{y}) := \mathcal{G}_1(\dot{y}) \cup \mathcal{G}_2(\dot{y}) \cup \mathcal{G}_3(\dot{y})$, where

$$\mathcal{G}_1(\dot{y}) := \{(c_L, c_R) \in [0, R]^2 : c_L > c_R, f(c_L) - \dot{y}c_L = f(c_R) - \dot{y}c_R = F_\alpha(\dot{y})\},$$

$$\mathcal{G}_2(\dot{y}) := \{(c, c) \in [0, R]^2 : f(c) - \dot{y}c \leq F_\alpha(\dot{y})\},$$

$$\mathcal{G}_3(\dot{y}) := \{(c_L, c_R) \in [0, R]^2 : c_L < c_R, f(c_L) - \dot{y}c_L = f(c_R) - \dot{y}c_R \leq F_\alpha(\dot{y})\}.$$

We refer the reader to Figure 2 for a graphical representation of $\mathcal{G}_\alpha(\dot{y})$




FIGURE 2. The set $\mathcal{G}_\alpha(\dot{y})$ (thick lines) in the case of a flux function of the form $f(\rho) = V\rho(1 - \rho/R)$, as in Section 3.

The equivalence between Definition 1.1 and [12, Definition 4.1] can be proved as in [4, Proposition 2.6].

Systems of the type (1) arise in the modeling of moving bottlenecks in traffic flows, see [12, 16] and Section 3 below, where they are coupled with an ODE depending on the downstream traffic velocity and describing the trajectory of a slow moving vehicle (a bus or a truck) acting as a bottleneck.

This paper is a first step towards establishing well-posedness for the strongly coupled models [12, 16]. Section 2 presents the main result, stating the \mathbf{L}^1 Lipschitz continuous dependence of solutions of (1) from the initial data and the constraint trajectory. Section 3 describes in details the related traffic flow model with moving bottleneck. Technical proof details are deferred to Appendix A.

2. Lipschitz continuous dependence of BV solutions with respect to \dot{y} .

Let us fix the constraint trajectory $y \in \mathbf{W}^{1,\infty}(\mathbb{R}^+; \mathbb{R})$ with $0 \leq \dot{y} < f'(0)$ and the initial datum $\rho_0 \in \text{BV}(\mathbb{R}; [0, R])$, and let $\rho \in \mathcal{C}^0(\mathbb{R}^+; \mathbf{L}^1(\mathbb{R}; [0, R]))$ be a solution of (1) in the sense of Definition 1.1. Applying the coordinate transformation $\tilde{\rho}(t, x) := \rho(t, x + y(t))$, $\tilde{\rho}$ is a weak entropy solution of the problem

$$\partial_t \tilde{\rho} + \partial_x F(\tilde{\rho}) = 0, \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \tag{6a}$$

$$\tilde{\rho}(0, x) = \rho_0(x + y_0), \quad x \in \mathbb{R}, \tag{6b}$$

$$F(\tilde{\rho}(t, 0)) \leq \tilde{F}_\alpha(\dot{y}(t)), \quad t \in \mathbb{R}^+, \tag{6c}$$

in the sense of [4, Proposition 2.6(A)] where we have set $F(\tilde{\rho}) := f(\tilde{\rho}) - \dot{y} \tilde{\rho}$. Existence and uniqueness for (6) are proved in [4, 10]. In particular, we have that for every $k \in [0, R]$ and for all $\varphi \in \mathcal{C}_c^1(\mathbb{R}^2; \mathbb{R}^+)$ such that $\varphi(t, 0) = 0, t > 0$,

$$\int_{\mathbb{R}^+} \int_{\mathbb{R}} (|\tilde{\rho} - k| \partial_t \varphi + \Phi(\dot{y}; \tilde{\rho}, k) \partial_x \varphi) dx dt + \int_{\mathbb{R}} |\rho_0(x + y_0) - k| \varphi(0, x) dx \geq 0,$$

where we have set

$$\Phi(\dot{y}; a, b) := \text{sgn}(a - b) (f(a) - f(b)) - \dot{y}|a - b|, \quad \text{for } a, b \in \mathbb{R},$$

and

$$(\tilde{\rho}(t, 0-), \tilde{\rho}(t, 0+)) \in \mathcal{G}_\alpha(\dot{y}(t)) \quad \text{for a. e. } t > 0,$$

see [4, Proposition 2.6]. Moreover, we note that, since $\alpha < 1$ and f is strictly concave, we have

$$\tilde{F}_\alpha(\dot{y}) = f_\alpha(\rho_\alpha(\dot{y})) - \dot{y} \rho_\alpha(\dot{y}) < f(\rho_\alpha(\dot{y})) - \dot{y} \rho_\alpha(\dot{y}) = \max_{\rho \in [0, R]} F(\rho). \tag{7}$$

Remark 1. Solution to (6) are not in BV in general (see [1, 10]), but thanks to the strict inequality in (7) and since $y \in \mathbf{W}^{1,\infty}(\mathbb{R}^+; \mathbb{R})$ implies $\dot{y} \in \text{BV}_{loc}(\mathbb{R}^+)$, we can conclude that $\tilde{\rho}(t, \cdot) \in \text{BV}(\mathbb{R}; [0, R])$ for any $t > 0$, as in [12].

We compare solutions of (6) corresponding to different constraint trajectories y and z .

Theorem 2.1. *Assume $y, z \in \mathbf{W}^{1,\infty}(\mathbb{R}^+; \mathbb{R})$, with $0 \leq \dot{y}, \dot{z} < f'(0)$, $\tilde{\rho}_0, \tilde{\sigma}_0 \in \mathbf{L}^\infty(\mathbb{R}, [0, R])$ and $\tilde{\rho}_0 - \tilde{\sigma}_0 \in \mathbf{L}^1(\mathbb{R})$. Let $\tilde{\rho}, \tilde{\sigma} \in \mathcal{C}^0(\mathbb{R}^+; \mathbf{L}^1(\mathbb{R}; [0, R]))$ be solutions of (6) corresponding respectively to $y, \tilde{\rho}_0$ and $z, \tilde{\sigma}_0$. Moreover, let $C_t = \sup_{s \in [0, t]} \text{TV}(\tilde{\rho}(s, \cdot))$ be finite. Then we have*

$$\|\tilde{\rho}(t, \cdot) - \tilde{\sigma}(t, \cdot)\|_{\mathbf{L}^1(\mathbb{R})} \leq \|\tilde{\rho}_0 - \tilde{\sigma}_0\|_{\mathbf{L}^1(\mathbb{R})} + (C_t + 2R) \|\dot{y} - \dot{z}\|_{\mathbf{L}^1([0, t])}. \tag{8}$$

Proof. The two solutions $\tilde{\rho}, \tilde{\sigma}$ satisfy

$$\partial_t |\tilde{\rho} - k| + \partial_x \Phi(\dot{y}; \tilde{\rho}, k) \leq 0, \tag{9}$$

$$\partial_t |\tilde{\sigma} - k| + \partial_x \Phi(\dot{z}; \tilde{\sigma}, k) \leq 0, \tag{10}$$

in $\mathcal{D}'(\mathbb{R}^+ \times \mathbb{R}^*)$ (where we have noted $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$). Following the proofs of [6, Lemma 15] and [7, Theorem 3.1] we observe that

$$\partial_t |\tilde{\rho} - k| + \partial_x \Phi(\dot{y}; \tilde{\rho}, k) = \partial_t |\tilde{\rho} - k| + \partial_x \Phi(\dot{z}; \tilde{\rho}, k) + \partial_x \Phi(\dot{y}; \tilde{\rho}, k) - \partial_x \Phi(\dot{z}; \tilde{\rho}, k)$$

therefore

$$\begin{aligned} \partial_t |\tilde{\rho} - k| + \partial_x \Phi(\dot{z}; \tilde{\rho}, k) &\leq \partial_x \Phi(\dot{z}; \tilde{\rho}, k) - \partial_x \Phi(\dot{y}; \tilde{\rho}, k) \\ &\leq |\dot{y} - \dot{z}| |\partial_x \tilde{\rho}|. \end{aligned} \tag{11}$$

Applying the classical Kruřkov doubling of variables technique [15], with a test function $\psi \in C_c^1(\mathbb{R}^2; \mathbb{R}^+)$ such that $\psi(t, 0) = 0$, we deduce the following Kato inequality

$$\int_{\mathbb{R}^+} \int_{\mathbb{R}} (|\tilde{\rho} - \tilde{\sigma}| \partial_t \psi + \Phi(\dot{z}; \tilde{\rho}, \tilde{\sigma}) \partial_x \psi) dx dt + \int_{\mathbb{R}} |\tilde{\rho}_0 - \tilde{\sigma}_0| \psi(0, x) dx + C_t \|\psi\|_{\infty} \int_{\mathbb{R}^+} |\dot{y} - \dot{z}| dt \geq 0,$$

We now choose the test function $\psi(t, x) = \theta_{\varepsilon}(x) \xi(t, x)$, where $\xi \in C_c^1(\mathbb{R}^2; \mathbb{R}^+)$ is an approximation of the characteristic function of the trapezoid

$$\{(s, x) : |x| \leq M + L(t - s), 0 \leq s \leq t\}$$

(where $L \geq \sup_{\rho \in [0, R], s \in [0, t]} |f'(\rho) - \dot{z}(s)|$ and $M \in \mathbb{R}, M > 0$) and θ_{ε} a smooth approximation of $x \rightarrow \min\{|x|/\varepsilon, 1\}$. Following the proof of [10, Proposition 4.4] and letting $\varepsilon \searrow 0$ we get

$$\begin{aligned} & \int_{-M}^M |\tilde{\rho}(t, x) - \tilde{\sigma}(t, x)| dx \\ & \leq \int_{-M-Lt}^{M+Lt} |\tilde{\rho}_0(x) - \tilde{\sigma}_0(x)| dx + C \int_0^t |\dot{y}(s) - \dot{z}(s)| ds \\ & \quad + \int_0^t (\Phi(\dot{z}; \tilde{\rho}(t, 0+), \tilde{\sigma}(t, 0+)) - \Phi(\dot{z}; \tilde{\rho}(t, 0-), \tilde{\sigma}(t, 0-))) ds. \end{aligned}$$

By Lemma A.1, the last integrand can be bounded by

$$\int_0^t (\Phi(\dot{z}; \tilde{\rho}(t, 0+), \tilde{\sigma}(t, 0+)) - \Phi(\dot{z}; \tilde{\rho}(t, 0-), \tilde{\sigma}(t, 0-))) ds \leq 2R \int_0^t |\dot{y}(s) - \dot{z}(s)| ds. \tag{12}$$

Letting $M \rightarrow \infty$, we recover (8). □

We are now able to state the well-posedness of problem (1).

Corollary 1. *Assume $y, z \in \mathbf{W}^{1,\infty}(\mathbb{R}^+; \mathbb{R})$, with $0 \leq \dot{y}, \dot{z} < f'(0)$, $\rho_0, \sigma_0 \in \mathbf{L}^{\infty}(\mathbb{R}, [0, R])$ and $\rho_0 - \sigma_0 \in \mathbf{L}^1(\mathbb{R})$. Let $\rho, \sigma \in C^0(\mathbb{R}^+; \mathbf{L}^1(\mathbb{R}; [0, R]))$ be solutions of (1) corresponding respectively to y, ρ_0 and z, σ_0 . Moreover, let $C_t = \sup_{s \in [0, t]} TV(\rho(s, \cdot))$ be finite. Then we have*

$$\begin{aligned} & \|\rho(t, \cdot) - \sigma(t, \cdot)\|_{\mathbf{L}^1(\mathbb{R})} \\ & \leq \|\rho_0 - \sigma_0\|_{\mathbf{L}^1(\mathbb{R})} + 2C_t |y(0) - z(0)| + (2C_t + 2R) \|\dot{y} - \dot{z}\|_{\mathbf{L}^1([0, t])}. \end{aligned} \tag{13}$$

Proof. Setting

$$\tilde{\rho}(t, x) = \rho(t, x + y(t)) \quad \text{and} \quad \tilde{\sigma}(t, x) = \sigma(t, x + z(t)),$$

for any $t > 0$ we get

$$\begin{aligned} \int_{\mathbb{R}} |\rho(t, x) - \sigma(t, x)| dx &= \int_{\mathbb{R}} |\rho(t, x + z(t)) - \sigma(t, x + z(t))| dx \\ &= \int_{\mathbb{R}} |\rho(t, x + z(t)) \mp \rho(t, x + y(t)) - \sigma(t, x + z(t))| dx \\ &\leq |y(t) - z(t)| TV(\rho(t, \cdot)) + \int_{\mathbb{R}} |\tilde{\rho}(t, x) - \tilde{\sigma}(t, x)| dx. \end{aligned}$$

Therefore, by (8), we get

$$\begin{aligned} & \int_{\mathbb{R}} |\rho(t, x) - \sigma(t, x)| dx \\ & \leq C_t |y(t) - z(t)| + \int_{\mathbb{R}} |\tilde{\rho}_0(x) - \tilde{\sigma}_0(x)| dx + (C_t + 2R) \int_0^t |\dot{y}(s) - \dot{z}(s)| ds \\ & \leq \int_{\mathbb{R}} |\rho_0(x) - \sigma_0(x)| dx + |y(0) - z(0)| \text{TV}(\rho_0) \\ & \quad + C_t \left(|y(0) - z(0)| + \int_0^t |\dot{y}(s) - \dot{z}(s)| ds \right) + (C_t + 2R) \int_0^t |\dot{y}(s) - \dot{z}(s)| ds \\ & = \int_{\mathbb{R}} |\rho_0(x) - \sigma_0(x)| dx + 2C_t |y(0) - z(0)| + (2C_t + 2R) \int_0^t |\dot{y}(s) - \dot{z}(s)| ds, \end{aligned}$$

where we have used the estimate $|y(t) - z(t)| \leq |y(0) - z(0)| + \int_0^t |\dot{y}(s) - \dot{z}(s)| ds$. \square

3. Application to traffic modeling. Setting

$$f(\rho) := \rho v(\rho),$$

where $v(\rho) = V(1 - \rho/R)$ is the mean traffic speed, V being the maximal velocity allowed on the road, problem (1) can be used to describe the situation of a moving bottleneck along a road, see [12]. In this case, we get $f_\alpha(\rho) = V\rho(1 - \frac{\rho}{\alpha R})$ and $\rho_\alpha(\dot{y}) = \frac{\alpha R}{2} \left(1 - \frac{\dot{y}}{V}\right)$, so that

$$\tilde{F}_\alpha(\dot{y}) = \frac{\alpha R}{4V} (V - \dot{y})^2.$$

Let us suppose that a slow and large vehicle, like for example a bus or a truck moves on the road. The slow vehicle, that in the following we will refer as “the bus”, reduces the road capacity and moves with a trajectory given by the following ODE:

$$\begin{cases} \dot{y}(t) = \omega(\rho(t, y(t)+)), & t \in \mathbb{R}^+, \\ y(0) = y_0, \end{cases} \tag{14}$$

where the velocity of the bus is given by the following traffic density dependent function (see Figure 3)

$$\omega(\rho) = \begin{cases} V_b & \text{if } \rho \leq \rho^* \doteq R(1 - V_b/V), \\ v(\rho) & \text{otherwise.} \end{cases} \tag{15}$$

This means that if the traffic is not too congested, the bus moves at its own maximal speed $V_b < V$. When the surrounding traffic density becomes too high, the bus adapts its velocity accordingly. In particular, it is not possible for the bus to overtake the cars.

Solutions of the coupled system (1), (14) for general initial data are defined as follows.

Definition 3.1. A couple $(\rho, y) \in \mathcal{C}^0(\mathbb{R}^+; \mathbf{L}^1(\mathbb{R}; [0, R])) \times \mathbf{W}^{1,1}(\mathbb{R}^+; \mathbb{R})$ is a solution to (1) if




FIGURE 3. Bus and cars speed.

1. ρ satisfies Kruřkov entropy conditions [15] on $(\mathbb{R}^+ \times \mathbb{R}) \setminus \{(t, y(t)) : t \in \mathbb{R}^+\}$, i.e. for every $k \in [0, R]$ and for all $\varphi \in C_c^1(\mathbb{R}^2; \mathbb{R}^+)$ and $\varphi(t, y(t)) = 0, t > 0$,

$$\int_{\mathbb{R}^+} \int_{\mathbb{R}} (|\rho - k| \partial_t \varphi + \text{sgn}(\rho - k) (f(\rho) - f(k)) \partial_x \varphi) dx dt + \int_{\mathbb{R}} |\rho_o - k| \varphi(0, x) dx \geq 0 ; \tag{16a}$$

2. for a. e. $t > 0$ the one-sided traces of ρ at $x = y(t)$ satisfy

$$(\rho(t, y(t)-), \rho(t, y(t)+)) \in \mathcal{G}_\alpha(\dot{y}(t)); \tag{16b}$$

3. y is a Carathéodory solution of (14), i.e. for a.e. $t \in \mathbb{R}^+$

$$y(t) = y_o + \int_0^t \omega(\rho(s, y(s)+)) ds . \tag{16c}$$

The proof of existence of solutions for the general Cauchy problem (1) strongly coupled with the bus trajectory (14) with BV initial data can be found in [12]. For completeness, we recall the definition of the solution of a Riemann problem, as given in [12]. Let us consider a Riemann type initial datum

$$\rho_0(x) = \begin{cases} \rho_L & \text{if } x < 0, \\ \rho_R & \text{if } x > 0, \end{cases} \quad y_0 = 0. \tag{17}$$

Denote by \mathcal{R} the standard (i.e., without the constraint (1c)) Riemann solver for (1a)-(1b)-(17), i.e., the (right continuous) map $(t, x) \mapsto \mathcal{R}(\rho_L, \rho_R)(x/t)$ given by the standard weak entropy solution, see for instance [8, Chapter 5]. Moreover, assume that \dot{y} is constant and let $\check{\rho} = \check{\rho}(\dot{y})$ and $\hat{\rho} = \hat{\rho}(\dot{y})$, with $\check{\rho} \leq \hat{\rho}$, be the intersection points of the flux function $f(\rho)$ with the line $f_\alpha(\rho_\alpha) + \dot{y}(\rho - \rho_\alpha)$ (see Figure 1(a)):

$$\begin{aligned} \check{\rho}(\dot{y}) &= \frac{R}{2V} (1 - \sqrt{1 - \alpha}) (V - \dot{y}), \\ \hat{\rho}(\dot{y}) &= \frac{R}{2V} (1 + \sqrt{1 - \alpha}) (V - \dot{y}). \end{aligned} \tag{18}$$

Definition 3.2. The constrained Riemann solver $\mathcal{R}^\alpha : [0, R]^2 \rightarrow \mathbf{L}_{\text{loc}}^1(\mathbb{R}; [0, R])$ corresponding to (1), (14), (17) is defined as follows.

1. If $f(\mathcal{R}(\rho_L, \rho_R)(V_b)) > F_\alpha + V_b \mathcal{R}(\rho_L, \rho_R)(V_b)$, then

$$\mathcal{R}^\alpha(\rho_L, \rho_R)(x/t) = \begin{cases} \mathcal{R}(\rho_L, \hat{\rho}(V_b))(x/t) & \text{if } x < V_b t, \\ \mathcal{R}(\check{\rho}(V_b), \rho_R)(x/t) & \text{if } x \geq V_b t, \end{cases} \quad \text{and } y(t) = V_b t.$$

2. If $V_b \mathcal{R}(\rho_L, \rho_R)(V_b) \leq f(\mathcal{R}(\rho_L, \rho_R)(V_b)) \leq F_\alpha + V_b \mathcal{R}(\rho_L, \rho_R)(V_b)$, then

$$\mathcal{R}^\alpha(\rho_L, \rho_R) = \mathcal{R}(\rho_L, \rho_R) \quad \text{and} \quad y(t) = V_b t.$$

3. If $f(\mathcal{R}(\rho_L, \rho_R)(V_b)) < V_b \mathcal{R}(\rho_L, \rho_R)(V_b)$, i.e., $v(\mathcal{R}(\rho_L, \rho_R)(V_b)) < V_b$ then

$$\mathcal{R}^\alpha(\rho_L, \rho_R) = \mathcal{R}(\rho_L, \rho_R) \quad \text{and} \quad y(t) = v(\rho_R)t.$$

Note that, when the constraint is enforced (point 1. in the above definition), a non-classical shock arises between $\hat{\rho}(V_b)$ and $\check{\rho}(V_b)$, which satisfies the Rankine-Hugoniot condition but violates the Lax entropy condition, see Figure 4 for an example.

Remark 2. Unfortunately, no result about the Lipschitz continuous dependence of the solution $y = y(t)$ of (14) from the solution $\rho = \rho(t, x)$ of (1) is known at present. Related results [9, 11] concerning uniqueness and continuous dependence for ODEs of the form (14) hold under hypothesis on the speed function ω that are not satisfied by (15).

Appendix A. Proof of (12).

Lemma A.1. For any $(\rho_-, \rho_+) \in \mathcal{G}_\alpha(\dot{y})$ and $(\sigma_-, \sigma_+) \in \mathcal{G}_\alpha(\dot{z})$, it holds

$$\Phi(\dot{z}; \rho_+, \sigma_+) - \Phi(\dot{z}; \rho_-, \sigma_-) \leq 2R|\dot{y} - \dot{z}|.$$

Proof. First of all, let us remark that

$$\Phi(\dot{z}; \rho_+, \sigma_+) - \Phi(\dot{z}; \rho_-, \sigma_-) \tag{19}$$

$$= \text{sgn}(\rho_+ - \sigma_+) (f(\rho_+) - \dot{z}\rho_+ - f(\sigma_+) + \dot{z}\sigma_+) \\ - \text{sgn}(\rho_- - \sigma_-) (f(\rho_-) - \dot{z}\rho_- - f(\sigma_-) + \dot{z}\sigma_-)$$

$$= (\lambda(\rho_+, \sigma_+) - \dot{z})|\rho_+ - \sigma_+| - (\lambda(\rho_-, \sigma_-) - \dot{z})|\rho_- - \sigma_-|, \tag{20}$$

where

$$\lambda(\rho, \sigma) = \frac{f(\rho) - f(\sigma)}{\rho - \sigma}.$$

Without loss of generality, we can assume $\dot{y} < \dot{z}$. Therefore we get

$$F_\alpha(\dot{y}) - F_\alpha(\dot{z}) \geq \rho_\alpha(\dot{z})(\dot{z} - \dot{y}) > 0. \tag{21}$$

We distinguish the following cases:

1. $(\rho_-, \rho_+) \in \mathcal{G}_1(\dot{y})$: we observe that

$$\rho_- = \hat{\rho}, \quad \rho_+ = \check{\rho} \quad \text{and} \quad f(\rho_-) - \dot{y}\rho_- = f(\rho_+) - \dot{y}\rho_+ = F_\alpha(\dot{y}).$$

Depending on the values of σ_-, σ_+ , different situations can occur as shown in Figure 5:


1.1 $(\sigma_-, \sigma_+) \in \mathcal{G}_1(\dot{z})$: in this case $\sigma_- = \hat{\sigma}$, $\sigma_+ = \check{\sigma}$, as shown in Figure 5a and $f(\sigma_-) - \dot{z}\sigma_- = f(\sigma_+) - \dot{z}\sigma_+ = F_\alpha(\dot{z})$, therefore

$$(19) = (F_\alpha(\dot{y}) + \dot{y}\rho_+ - \dot{z}\rho_+ - F_\alpha(\dot{z})) - (F_\alpha(\dot{y}) + \dot{y}\rho_- - \dot{z}\rho_- - F_\alpha(\dot{z})) \\ = (\rho_- - \rho_+)(\dot{z} - \dot{y}) \\ \leq R|\dot{y} - \dot{z}|.$$


1.2 $(\sigma_-, \sigma_+) \in \mathcal{G}_2(\dot{z})$: we set

$$\sigma := \sigma_- = \sigma_+ \quad \text{and} \quad f(\sigma) - \dot{z}\sigma =: F(\sigma) \leq F_\alpha(\dot{z}).$$


The following cases can occur:



(a) Example of solution corresponding to case 1. of Definition 3.2 for $\rho_L = \rho_R = \bar{\rho}$, $\hat{\rho}_\alpha < \bar{\rho} < \check{\rho}_\alpha$.



(b) Example of solution corresponding to case 2. of Definition 3.2 for $0 < \rho_L < \hat{\rho}_\alpha$ and $\hat{\rho}_\alpha < \rho_R < \rho^*$.



(c) Example of solution corresponding to case 3. of Definition 3.2 for $\rho_L, \rho_R > \rho^*$.

FIGURE 4. Different solutions of the Riemann problem (17). Each subfigure illustrates a point of the Definition 3.2: fundamental diagram representation (left) and space-time diagram (right).




FIGURE 5. Case 1

- $0 \leq \sigma \leq \check{\sigma}$:

$$\begin{aligned}
 (19) &= (F_\alpha(\check{y}) + \check{y}\rho_+ - \check{z}\rho_+ - F(\sigma)) - (F_\alpha(\check{y}) + \check{y}\rho_- - \check{z}\rho_- - F(\sigma)) \\
 &= (\rho_- - \rho_+)(\check{z} - \check{y}) \\
 &\leq R|\check{y} - \check{z}|.
 \end{aligned}$$

- $\hat{\sigma} \leq \sigma \leq \hat{\rho}$, see Figure 5b:

$$\begin{aligned}
 (19) &= -(F_\alpha(\check{y}) + \check{y}\rho_+ - \check{z}\rho_+ - F(\sigma)) - (F_\alpha(\check{y}) + \check{y}\rho_- - \check{z}\rho_- - F(\sigma)) \\
 &= 2F(\sigma) - 2F_\alpha(\check{y}) + (\rho_+ + \rho_-)(\check{z} - \check{y}) \\
 &\leq 2F_\alpha(\check{z}) - 2F_\alpha(\check{y}) + 2R(\check{z} - \check{y}) \\
 &= -2\rho_\alpha(\check{z})(\check{z} - \check{y}) + 2R(\check{z} - \check{y}) \\
 &\leq 2R|\check{y} - \check{z}|.
 \end{aligned}$$

- $\hat{\rho} \leq \sigma \leq R$:

$$\begin{aligned}
 (19) &= -(F_\alpha(\check{y}) + \check{y}\rho_+ - \check{z}\rho_+ - F(\sigma)) + (F_\alpha(\check{y}) + \check{y}\rho_- - \check{z}\rho_- - F(\sigma)) \\
 &= (\rho_+ - \rho_-)(\check{z} - \check{y}) \leq 0.
 \end{aligned}$$

1.3 $(\sigma_-, \sigma_+) \in \mathcal{G}_3(\dot{z})$: we set

$$f(\sigma_-) - \dot{z}\sigma_- = f(\sigma_+) - \dot{z}\sigma_+ =: F(\sigma) \leq F_\alpha(\dot{z}).$$

We observe that $\text{sgn}(\rho_+ - \sigma_+) < 0$ and $\text{sgn}(\rho_- - \sigma_-) > 0$, see Figure 5c. Therefore

$$\begin{aligned} (19) &= -(F_\alpha(\dot{y}) + \dot{y}\rho_+ - \dot{z}\rho_+ - F(\sigma)) - (F_\alpha(\dot{y}) + \dot{y}\rho_- - \dot{z}\rho_- - F(\sigma)) \\ &= 2F(\sigma) - 2F_\alpha(\dot{y}) + (\rho_+ + \rho_-)(\dot{z} - \dot{y}) \\ &\leq 2F_\alpha(\dot{z}) - 2F_\alpha(\dot{y}) + 2R(\dot{z} - \dot{y}) \\ &= -2\rho_\alpha(\dot{z})(\dot{z} - \dot{y}) + 2R(\dot{z} - \dot{y}) \\ &\leq 2R|\dot{y} - \dot{z}|. \end{aligned}$$

2. $(\rho_-, \rho_+) \in \mathcal{G}_2(\dot{y})$: we set

$$\rho := \rho_- = \rho_+ \quad \text{and} \quad f(\rho) - \dot{y}\rho =: F(\rho) \leq F_\alpha(\dot{y}).$$

as illustrated in Figure 6.




FIGURE 6. Case 2

2.1 $(\sigma_-, \sigma_+) \in \mathcal{G}_1(\dot{z})$:

- $0 \leq \rho \leq \check{\sigma}$:

$$(19) = -(F(\rho) + \dot{y}\rho - \dot{z}\rho - F_\alpha(\dot{z})) + (F(\rho) + \dot{y}\rho - \dot{z}\rho - F_\alpha(\dot{z})) = 0.$$

- $\check{\sigma} \leq \rho \leq \check{\rho}$, this case is displayed in Figure 6a:

$$\begin{aligned} (19) &= (F(\rho) + \dot{y}\rho - \dot{z}\rho - F_\alpha(\dot{z})) + (F(\rho) + \dot{y}\rho - \dot{z}\rho - F_\alpha(\dot{z})) \\ &= 2F(\rho) - 2F_\alpha(\dot{z}) + 2\rho(\dot{y} - \dot{z}) \\ &\leq 2(F_\alpha(\dot{y}) - F_\alpha(\dot{z})) \\ &\leq 2\rho_\alpha(\dot{y})(\dot{z} - \dot{y}) \\ &\leq 2\alpha R|\dot{y} - \dot{z}|. \end{aligned}$$

- $\hat{\rho} \leq \rho \leq R$:

$$(19) = (F(\rho) + \dot{y}\rho - \dot{z}\rho - F_\alpha(\dot{z})) - (F(\rho) + \dot{y}\rho - \dot{z}\rho - F_\alpha(\dot{z})) = 0.$$

2.2 $(\sigma_-, \sigma_+) \in \mathcal{G}_2(\dot{z})$, illustrated in Figure 6b:

We observe that $\text{sgn}(\rho_+ - \sigma_+) = \text{sgn}(\rho_- - \sigma_-) = \text{sgn}(\rho - \sigma)$, therefore

$$(19) = \text{sgn}(\rho - \sigma) [(F(\rho) + \dot{y}\rho - \dot{z}\rho - F(\sigma)) - (F(\rho) + \dot{y}\rho - \dot{z}\rho - F(\sigma))] = 0.$$

2.3 $(\sigma_-, \sigma_+) \in \mathcal{G}_3(\dot{z})$ shown in Figure 6c: If $\text{sgn}(\rho - \sigma_+) = \text{sgn}(\rho - \sigma_-)$, we get

$$(19) = \text{sgn}(\rho - \sigma_+) [(F(\rho) + \dot{y}\rho - \dot{z}\rho - F(\sigma)) - (F(\rho) + \dot{y}\rho - \dot{z}\rho - F(\sigma))] = 0.$$

Otherwise, we have that $\sigma_- \leq \rho \leq \check{\rho}$ or $\hat{\rho} \leq \rho \leq \sigma_+$. In this case

$$\begin{aligned} \lambda(\rho_+, \sigma_+) - \dot{z} &= \frac{f(\sigma_+) - f(\rho)}{\sigma_+ - \rho} - \frac{f(\sigma_+) - f(\sigma_-)}{\sigma_+ - \sigma_-} \leq 0, \\ \lambda(\rho_-, \sigma_-) - \dot{z} &= \frac{f(\sigma_-) - f(\rho)}{\sigma_- - \rho} - \frac{f(\sigma_+) - f(\sigma_-)}{\sigma_+ - \sigma_-} \geq 0, \end{aligned}$$

by the concavity of f . Therefore, (19) ≤ 0 by (20).

3. $(\rho_-, \rho_+) \in \mathcal{G}_3(\dot{y})$: we set

$$f(\rho_-) - \dot{y}\rho_- = f(\rho_+) - \dot{y}\rho_+ =: F(\rho) \leq F_\alpha(\dot{y}).$$

See Figure 7, for a graphical representation.

3.1 $(\sigma_-, \sigma_+) \in \mathcal{G}_1(\dot{z})$, see Figure 7a.

$$\begin{aligned} (19) &= (F(\rho) + \dot{y}\rho_+ - \dot{z}\rho_+ - F_\alpha(\dot{z})) + (F(\rho) + \dot{y}\rho_- - \dot{z}\rho_- - F_\alpha(\dot{z})) \\ &= 2F(\rho) - 2F_\alpha(\dot{z}) + (\rho_+ + \rho_-)(\dot{y} - \dot{z}) \\ &\leq 2(F_\alpha(\dot{y}) - F_\alpha(\dot{z})) \\ &\leq 2\rho_\alpha(\dot{y})(\dot{z} - \dot{y}) \\ &\leq 2\alpha R|\dot{y} - \dot{z}|. \end{aligned}$$

3.2 $(\sigma_-, \sigma_+) \in \mathcal{G}_2(\dot{z})$: If $\text{sgn}(\rho_+ - \sigma) = \text{sgn}(\rho_- - \sigma)$, we get

$$\begin{aligned} (19) &= \text{sgn}(\rho_+ - \sigma) [(F(\rho) + \dot{y}\rho_+ - \dot{z}\rho_+ - F(\sigma)) - (F(\rho) + \dot{y}\rho_- - \dot{z}\rho_- - F(\sigma))] \\ &\leq (\rho_+ - \rho_-)(\dot{z} - \dot{y}) \\ &\leq R|\dot{y} - \dot{z}|. \end{aligned}$$




FIGURE 7. Case 3

Otherwise as shown in Figure 7b, we have that $\rho_- \leq \sigma \leq \hat{\sigma}$ or $\hat{\sigma} \leq \sigma \leq \rho_+$. In this case

$$\begin{aligned} \lambda(\rho_+, \sigma_+) - \dot{z} &\leq \lambda(\rho_+, \sigma_+) - \dot{y} \\ &= \frac{f(\rho_+) - f(\sigma)}{\rho_+ - \sigma} - \frac{f(\rho_+) - f(\rho_-)}{\rho_+ - \rho_-} \leq 0. \end{aligned}$$

Moreover, we observe that $\lambda(\rho_-, \sigma_-) > \dot{y}$. Indeed

$$\lambda(\rho_-, \sigma_-) - \dot{y} = \frac{f(\rho_-) - f(\sigma)}{\rho_- - \sigma} - \frac{f(\rho_+) - f(\rho_-)}{\rho_+ - \rho_-} \geq 0.$$

Therefore, by (20)

$$(19) \leq R|\dot{y} - \dot{z}|.$$

3.3 $(\sigma_-, \sigma_+) \in \mathcal{G}_3(\dot{z})$:

We observe that one of the following relations must hold

$$\rho_- \leq \sigma_- < \sigma_+ \leq \rho_+, \quad \sigma_- \leq \rho_- < \sigma_+ \leq \rho_+, \quad \sigma_- \leq \rho_- < \rho_+ \leq \sigma_+.$$

For an example see Figure 7c. Therefore

$$\lambda(\rho_+, \sigma_+) - \dot{z} = \frac{f(\rho_+) - f(\sigma_+)}{\rho_+ - \sigma_+} - \frac{f(\sigma_+) - f(\sigma_-)}{\sigma_+ - \sigma_-} \leq 0,$$

$$\lambda(\rho_-, \sigma_-) - \dot{z} = \frac{f(\rho_-) - f(\sigma_-)}{\rho_- - \sigma_-} - \frac{f(\sigma_+) - f(\sigma_-)}{\sigma_+ - \sigma_-} \geq 0,$$

again by concavity of f . Hence (19) ≤ 0 by (20).

□

Acknowledgments. The authors are grateful to Boris Andreianov for suggesting links with the results in [6]. The authors also thank the two anonymous referees for the careful revision and the insightful comments, which were of great help in improving the paper.

REFERENCES

- [1] Adimurthi, R. Dutta, S. S. Ghoshal and G. D. Veerappa Gowda, Existence and nonexistence of TV bounds for scalar conservation laws with discontinuous flux, *Comm. Pure Appl. Math.*, **64** (2011), 84–115.
- [2] B. Andreianov, C. Donadello, U. Razafison and M. D. Rosini, Riemann problems with non-local point constraints and capacity drop, *Math. Biosci. Eng.*, **12** (2015), 259–278.
- [3] B. Andreianov, C. Donadello and M. D. Rosini, A second-order model for vehicular traffics with local point constraints on the flow, *Math. Models Methods Appl. Sci.*, **26** (2016), 751–802.
- [4] B. Andreianov, P. Goatin and N. Seguin, Finite volume schemes for locally constrained conservation laws, *Numer. Math.*, **115** (2010), 609–645, With supplementary material available online.
- [5] B. Andreianov, K. H. Karlsen and N. H. Risebro, A theory of L^1 -dissipative solvers for scalar conservation laws with discontinuous flux, *Arch. Ration. Mech. Anal.*, **201** (2011), 27–86.
- [6] B. Andreianov, F. Lagoutière, N. Seguin and T. Takahashi, Well-posedness for a one-dimensional fluid-particle interaction model, *SIAM J. Math. Anal.*, **46** (2014), 1030–1052.
- [7] F. Bouchut and B. Perthame, Kruřkov’s estimates for scalar conservation laws revisited, *Trans. Amer. Math. Soc.*, **350** (1998), 2847–2870.
- [8] A. Bressan, *Hyperbolic Systems of Conservation Laws*, vol. 20 of Oxford Lecture Series in Mathematics and its Applications, Oxford University Press, Oxford, 2000, The one-dimensional Cauchy problem.
- [9] A. Bressan and W. Shen, Uniqueness for discontinuous ODE and conservation laws, *Nonlinear Anal.*, **34** (1998), 637–652.
- [10] R. M. Colombo and P. Goatin, A well posed conservation law with a variable unilateral constraint, *J. Differential Equations*, **234** (2007), 654–675.
- [11] R. M. Colombo and A. Marson, A Hölder continuous ODE related to traffic flow, *Proc. Roy. Soc. Edinburgh Sect. A*, **133** (2003), 759–772.
- [12] M. L. Delle Monache and P. Goatin, Scalar conservation laws with moving constraints arising in traffic flow modeling: an existence result, *J. Differential Equations*, **257** (2014), 4015–4029.
- [13] M. Garavello and P. Goatin, The Aw-Rascle traffic model with locally constrained flow, *J. Math. Anal. Appl.*, **378** (2011), 634–648.
- [14] M. Garavello and S. Villa, The Cauchy problem for the Aw-Rascle-Zhang traffic model with locally constrained flow, Preprint, 2016.
- [15] S. N. Kruřkov, First order quasilinear equations with several independent variables, *Mat. Sb. (N.S.)*, **81** (1970), 228–255.
- [16] S. Villa, P. Goatin and C. Chalons, Moving bottlenecks for the Aw-Rascle-Zhang traffic flow model, 2016, URL <https://hal.archives-ouvertes.fr/hal-01347925>, preprint.

Received October 2016; revised January 2017.

E-mail address: ml.dellemonache@inria.fr

E-mail address: paola.goatin@inria.fr