

## REDUCTION OF A MODEL FOR SODIUM EXCHANGES IN KIDNEY NEPHRON

MARTA MARULLI

University of Bologna, Department of Mathematics  
Piazza di Porta S. Donato 5, 40126 Bologna, Italy  
Université Sorbonne Paris Nord  
Laboratoire Analyse Géométrie et Applications (LAGA), CNRS UMR 7539  
F-93430, Villetaneuse, France

VUK MILIŠIĆ\* AND NICOLAS VAUCHELET

Université Sorbonne Paris Nord  
Laboratoire Analyse Géométrie et Applications (LAGA), CNRS UMR 7539  
F-93430, Villetaneuse, France

(Communicated by Pierre-Emmanuel Jabin)

**ABSTRACT.** This work deals with a mathematical analysis of sodium's transport in a tubular architecture of a kidney nephron. The nephron is modelled by two counter-current tubules. Ionic exchange occurs at the interface between the tubules and the epithelium and between the epithelium and the surrounding environment (interstitium). From a mathematical point of view, this model consists of a  $5 \times 5$  semi-linear hyperbolic system. In literature similar models neglect the epithelial layers. In this paper, we show rigorously that such models may be obtained by assuming that the permeabilities between lumen and epithelium are large. We show that when these permeabilities grow, solutions of the  $5 \times 5$  system converge to those of a reduced  $3 \times 3$  system without epithelial layers. The problem is defined on a bounded spacial domain with initial and boundary data. In order to show convergence, we use BV compactness, which leads to introduce initial layers and to handle carefully the presence of lateral boundaries. We then discretize both  $5 \times 5$  and  $3 \times 3$  systems, and show numerically the same asymptotic result, for a fixed meshsize.

**1. Introduction.** In this study, we consider a mathematical model for a particular component of the nephron, the functional unit of kidney [1]. It describes the ionic exchanges through the nephron tubules in the Henle's loop. The main function of kidneys is to filtrate the blood. Through filtration, secretion and excretion of filtered metabolic wastes and toxins, the kidneys are able to maintain a certain homeostatic balance within cells. The first models of nephrons and kidneys were developed in the 1950s with the purpose of explaining the concentration gradient, [10] (or [16] in Chap.3). In their attempt, the authors describe for the first time the urinary concentration mechanism as a consequence of countercurrent transport in the tubules. At a later time the point of view starts to change and a simple mathematical model

---

2020 *Mathematics Subject Classification.* Primary: 35L04, 35L60, 35L81; Secondary: 35Q92, 92C42.

*Key words and phrases.* Hyperbolic systems, relaxation limit, characteristics method, boundary layers, ionic exchanges.

\* Corresponding author.

for a single nephron has been introduced and described, [17]. Furthermore, thanks to the work of J.L. Stephenson [26] it turns out that the formation of a large axial concentration gradient in the kidney depends mainly on the different permeabilities in the tubules and on their counterflow arrangement. Most mathematical models were stationary systems of coupled differential equations. It is possible to find an historical excursus and how these types of models have been developed, referring to [26]. Sophisticated models have been developed to describe the transport of water and electrolyte in the kidney in the recent literature. At the microscopic level of description, cell-based models have been proposed in [23, 30, 14, 31, 4] to model small populations of nephrons at equilibrium. Macroscopic models have been also considered in [28, 27, 21, 5, 2, 9] without accounting for cell-specific transport mechanisms. Despite the development of such sophisticated models, some aspects of the fundamental functions of the kidney remain yet to be fully explained [15]. For example, how a concentrated urine can be produced by the mammalian kidney when the animal is deprived of water remains not entirely clear.

The loop of Henle and its architecture play an important role in the concentrated or diluted urine formation. In order to explain the regulation of urine's concentration, we analyse the counter-current transport in the 'ascending' and 'descending' tubules. There the ionic exchanges between the cell membrane and the environment where tubules are immersed, take place.

We consider a simplified model for sodium exchange in the kidney nephron : the nephron is modelled by an ascending (resp. descending) tubule, of length denoted  $L$ . Ionic exchanges and transport occur at the interface between the lumen and the epithelial layer (cell membrane) and at the interface between the cells and the interstitium (the surrounding tissue between tubules and blood vessels) c.f. Figure 1. Denoting by  $t \geq 0$  (resp.  $x \in (0, L)$ ) the time (resp. space) variable, the dynamics of ionic concentrations is modelled by the semi-linear hyperbolic system [19, 27, 28, 29] :

$$\begin{cases} a_1 \partial_t u_1 + \alpha \partial_x u_1 = J_1 = 2\pi r_1 P_1(q_1 - u_1) \\ a_2 \partial_t u_2 - \alpha \partial_x u_2 = J_2 = 2\pi r_2 P_2(q_2 - u_2) \\ a_3 \partial_t q_1 = J_{1,e} = 2\pi r_1 P_1(u_1 - q_1) + 2\pi r_{1,e} P_{1,e}(u_0 - q_1) \\ a_4 \partial_t q_2 = J_{2,e} = 2\pi r_2 P_2(u_2 - q_2) + 2\pi r_{2,e} P_{2,e}(u_0 - q_2) - G(q_2) \\ a_0 \partial_t u_0 = J_0 = 2\pi r_{1,e} P_{1,e}(q_1 - u_0) + 2\pi r_{2,e} P_{2,e}(q_2 - u_0) + G(q_2), \end{cases} \quad (1)$$

complemented with the boundary and initial conditions

$$\begin{aligned} u_1(t, 0) &= u_b(t), \quad u_1(t, L) = u_2(t, L), \quad \forall t > 0 \\ u_1(0, x) &= u_1^0(x), \quad u_2(0, x) = u_2^0(x), \quad u_0(0, x) = u_0^0(x), \\ q_1(0, x) &= q_1^0(x), \quad q_2(0, x) = q_2^0(x), \quad \forall x \in (0, L). \end{aligned} \quad (2)$$

In this model, we have used the following notations :

- $r_i$  : denote the radius for the lumen  $i$  ( $[m]$ ).
- $r_{i,e}$  : denote the radius for the tubule  $i$  with epithelium layer.
- Sodium's concentrations ( $[mol/m^3]$ ) :
  - $u_i(t, x)$  : in the lumen  $i$ ,
  - $q_i(t, x)$  : in the epithelial layer of lumen  $i$ ,
  - $u_0(t, x)$  : in the interstitium.

- Permeabilities ( $[m/s]$ ):  
 $P_i$  : between the lumen and the epithelium,  
 $P_{i,e}$  : between the epithelium and the interstitium.
- Section areas  $a_i$  ( $[m^2]$ ),  $i \in \{0, \dots, 4\}$  are defined as

$$a_1 = \pi r_1^2, \quad a_2 = \pi r_2^2, \quad a_3 = \pi(r_{1,e}^2 - r_1^2), \quad a_4 = \pi(r_{2,e}^2 - r_2^2), \quad a_0 = \pi \left( \frac{r_{1,e}^2 + r_{2,e}^2}{2} \right).$$

In this work we indicate as lumen the limb under consideration and as tubule the segment together with its epithelial layer. In physiological common language, the term ‘tubule’ refers to the cavity of lumen together with its related epithelial layer (membrane) as part of it, [16].

In the ascending tubule, the transport of solutes both by passive diffusion and active re-absorption uses  $Na^+/K^+$ -ATPases pumps, which exchange 3  $Na^+$  ions for 2  $K^+$  ions. This active transport is taken into account in (1) by a non-linear term given by Michaelis-Menten’s kinetics :

$$G(q_2) = V_{m,2} \left( \frac{q_2}{k_{M,2} + q_2} \right)^3 \quad (3)$$

where  $k_{M,2}$  and  $V_{m,2}$  are real positive constants. In each tubule, the fluid (mostly water) is assumed to flow at constant rate  $\alpha$  and we only consider one generic uncharged solute in two tubules as depicted in Figure 1.




FIGURE 1. Simplified model of the loop of Henle.  $q_1$ ,  $q_2$ ,  $u_1$  and  $u_2$  denote solute concentration in the epithelial layer and lumen of the descending/ascending limb, respectively.

In a recent paper [19], the authors have studied, from the modelling and biological perspective, the role of the epithelial layer in the ionic transport. The aim of this work is to clarify from the mathematical point of view the link between model (1) taking into account the epithelial layer and models neglecting it. In particular, when the permeability between the epithelium and the lumen is large it is expected that these two regions merge, allowing to reduce system (1) to a model with no epithelial layer. More precisely, as the permeabilities  $P_1$  and  $P_2$  grow large, we show

rigorously that solutions of (1) with boundary conditions (2) relax to solutions of a reduced system with no epithelial layer. From a mathematical and numerical point of view, system (1) may be seen as a hyperbolic system with a stiff source term. The source term is in some sense a relaxation of another hyperbolic system of smaller dimension. Such an approach has been widely studied in the literature, see e.g. [13, 22, 12, 25]. Since the initial data of the starting system for fixed  $\varepsilon$  has no reason to be compatible with the limit system, the mathematical analysis of this relaxation procedure should account for initial layers. For relaxed convection-diffusion systems, but in the Cauchy case, for *ill-prepared* data, (out of equilibrium with respect to the reduced manifold), the construction of initial layers and the corresponding error analysis in Sobolev spaces can be found in [6]. The proof of our convergence result is obtained thanks to a BV compactness argument in space and in time. This framework is better suited for purely hyperbolic problems. Another difficulty is due to the presence of the boundary, which must be carefully handled in the *a priori* estimates, in order to be uniform with respect to  $\varepsilon$ , the relaxation parameter depending on the permeabilities.

The outline of the paper is the following. In the next section, we provide the mathematical model and state our main result. Section 3 is devoted to the proof of existence of solution of our model. Then, we focus on the convergence of this solution when permeabilities go to infinity. To this aim, we first establish in section 4 *a priori* estimates. Using compactness results, we prove convergence in section 5. Numerical illustration of this convergence result is proposed in section 6. Finally an appendix is devoted to a formal computation of the convergence rate at stationary state.

**2. Main results.** Before presenting our main result, we list some assumptions which will be used throughout this paper.

**Assumption 2.1.** *We assume that the initial solute concentrations are non-negative and uniformly bounded in  $L^\infty(0, L)$  and with respect to the total variation :*

$$0 \leq u_1^0, u_2^0, q_1^0, q_2^0, u_0^0 \in BV(0, L) \cap L^\infty(0, L). \quad (4)$$

For detailed definitions of the BV setting we refer to the standard text-books [7, 32]. A more recent overview gives a global picture in an extensive way [11], it unifies also the diversity of definitions found in the literature dealing with the BV spaces in either the probabilistic or the deterministic context.

**Assumption 2.2.** *Boundary conditions are such that*

$$0 \leq u_b \in BV(0, T) \cap L^\infty(0, T). \quad (5)$$

**Assumption 2.3.** *Regularity and boundedness of  $G$ .*

*We assume that the non-linear function modelling active transport in the ascending limb is an odd and  $W^{2,\infty}(\mathbb{R})$  function :*

$$\forall x \geq 0, \quad G(-x) = -G(x), \quad 0 \leq G(x) \leq \|G\|_\infty, \quad 0 \leq G'(x) \leq \|G'\|_\infty. \quad (6)$$

*We notice that the function  $G$  defined on  $\mathbb{R}^+$  by the expression in (3) may be straightforwardly extended by symmetry on  $\mathbb{R}$  by a function satisfying (6).*

To simplify our notations in (1), we set  $2\pi r_{i,e} P_{i,e} = K_i$ ,  $i = 1, 2$  and  $2\pi r_i P_i = k_i$ ,  $i = 1, 2$ . The orders of magnitude of  $k_1$  and  $k_2$  are the same even if their values are not definitely equal, we may assume to further simplify the analysis that  $k_1 = k_2$ . We consider the case where permeability between the lumen and the epithelium is

large and we set,  $k_1 = k_2 = \frac{1}{\varepsilon}$  for  $\varepsilon \ll 1$ . Then, we investigate the limit  $\varepsilon$  goes to zero of the solutions of the following system :

$$a_1 \partial_t u_1^\varepsilon + \alpha \partial_x u_1^\varepsilon = \frac{1}{\varepsilon} (q_1^\varepsilon - u_1^\varepsilon) \quad (7a)$$

$$a_2 \partial_t u_2^\varepsilon - \alpha \partial_x u_2^\varepsilon = \frac{1}{\varepsilon} (q_2^\varepsilon - u_2^\varepsilon) \quad (7b)$$

$$a_3 \partial_t q_1^\varepsilon = \frac{1}{\varepsilon} (u_1^\varepsilon - q_1^\varepsilon) + K_1 (u_0^\varepsilon - q_1^\varepsilon) \quad (7c)$$

$$a_4 \partial_t q_2^\varepsilon = \frac{1}{\varepsilon} (u_2^\varepsilon - q_2^\varepsilon) + K_2 (u_0^\varepsilon - q_2^\varepsilon) - G(q_2^\varepsilon) \quad (7d)$$

$$a_0 \partial_t u_0^\varepsilon = K_1 (q_1^\varepsilon - u_0^\varepsilon) + K_2 (q_2^\varepsilon - u_0^\varepsilon) + G(q_2^\varepsilon) \quad (7e)$$

Formally, when  $\varepsilon \rightarrow 0$ , we expect the concentrations  $u_1^\varepsilon$  and  $q_1^\varepsilon$  to converge to the same function. The same happens for  $u_2^\varepsilon$  and  $q_2^\varepsilon$ . We denote  $u_1$ , respectively  $u_2$ , these limits. Adding (7a) to (7c) and (7b) to (7d), we end up with the system

$$\begin{aligned} a_1 \partial_t u_1^\varepsilon + a_3 \partial_t q_1^\varepsilon + \alpha \partial_x u_1^\varepsilon &= K_1 (u_0^\varepsilon - q_1^\varepsilon) \\ a_2 \partial_t u_2^\varepsilon + a_4 \partial_t q_2^\varepsilon - \alpha \partial_x u_2^\varepsilon &= K_2 (u_0^\varepsilon - q_2^\varepsilon) - G(q_2^\varepsilon). \end{aligned}$$

Passing formally to the limit when  $\varepsilon$  goes to 0, we arrive at

$$(a_1 + a_3) \partial_t u_1 + \alpha \partial_x u_1 = K_1 (u_0 - u_1) \quad (8)$$

$$(a_2 + a_4) \partial_t u_2 - \alpha \partial_x u_2 = K_2 (u_0 - u_2) - G(u_2), \quad (9)$$

coupled to the equation for the concentration in the interstitium obtained by passing into the limit in equation (7e)

$$a_0 \partial_t u_0 = K_1 (u_1 - u_0) + K_2 (u_2 - u_0) + G(u_2). \quad (10)$$

This system is complemented with the initial and boundary conditions

$$u_1(0, x) = \frac{(a_1 u_1^0(x) + a_3 q_1^0(x))}{(a_1 + a_3)}, \quad u_2(0, x) = \frac{(a_2 u_2^0(x) + a_4 q_2^0(x))}{(a_2 + a_4)}, \quad (11)$$

$$u_0(0, x) = u_0^0(x), \quad (12)$$

$$u_1(t, 0) = u_b(t), \quad u_2(t, L) = u_1(t, L). \quad (13)$$

Finally, we recover a simplified system for only three unknowns. From a physical point of view this means fusing the epithelial layer with the lumen. It turns out to merge the lumen and the epithelium into a single domain when we consider the limit of infinite permeability. For  $\varepsilon$  fixed, the actual speed at which  $u_1$  (resp.  $u_2$ ) is convected in (7a) is  $\alpha/a_1$  (resp  $-\alpha/a_2$  in (7b)). When  $\varepsilon$  goes to zero, since the permeability increases, the lumen and the epithelium layer are merged. Thus, as  $q_1$  (resp.  $q_2$ ) are not transported, this results in a slower convection at speed  $\frac{\alpha}{a_1+a_3}$  in the descending tubule (resp.  $\frac{\alpha}{a_2+a_4}$  in the ascending tubule).

The aim of this paper is to make these formal computations rigorous. For this sake, we define weak solutions associated to the limit system (8)-(10) :

**Definition 2.4.** Let  $u_1^0(x)$ ,  $u_2^0(x)$ ,  $u_0^0(x) \in L^1(0, L) \cap L^\infty(0, L)$  and  $u_b(t) \in L^1(0, T) \cap L^\infty(0, T)$ . We say that  $U(t, x) = (u_1(t, x), u_2(t, x), u_0(t, x))^\top \in L^\infty((0, T); L^1(0, L) \cap L^\infty(0, L))^3$  is a weak solution of system (8)-(10) if for all  $\phi \in \mathcal{S}_3$ , with

$$\mathcal{S}_3 := \{ \phi \in C^1([0, T] \times [0, L])^3, \quad \phi(T, x) = 0, \phi_1(t, L) = \phi_2(t, L), \text{ and } \phi_2(t, 0) = 0 \},$$

we have

$$\begin{aligned}
& \int_0^T \int_0^L u_1((a_1 + a_3)\partial_t \phi_1 + \alpha \partial_x \phi_1) dx dt + \alpha \int_0^T u_b(t) \phi_1(t, 0) dt \\
& + \int_0^L (a_1 + a_3) u_1(0, x) \phi_1(0, x) dx \\
& + \int_0^T \int_0^L u_2((a_2 + a_3)\partial_t \phi_2 - \alpha \partial_x \phi_2) dx dt + \int_0^L (a_2 + a_4) u_2(0, x) \phi_2(0, x) dx \quad (14) \\
& + \int_0^T \int_0^L \{a_0 u_0 \partial_t \phi_3 + K_1(u_1 - u_0)(\phi_3 - \phi_1) + K_2(u_2 - u_0)(\phi_3 - \phi_2) \\
& + G(u_2)(\phi_3 - \phi_2)\} dx dt + \int_0^L a_0 u_0^0(x) \phi_3(0, x) dx = 0.
\end{aligned}$$

More precisely, the main result reads

**Theorem 2.5.** *Let  $T > 0$  and  $L > 0$ . We assume that initial data and boundary conditions satisfy (4), (5), (6). Then, the weak solution  $(u_1^\varepsilon, u_2^\varepsilon, q_1^\varepsilon, q_2^\varepsilon, u_0^\varepsilon)$  of system (7) with boundary and initial conditions (2) converges, as  $\varepsilon$  goes to zero, to the weak solution of reduced (or limit) problem (8)–(10) complemented with (11)–(13). More precisely,*

$$\begin{aligned}
u_i^\varepsilon & \xrightarrow{\varepsilon \rightarrow 0} u_i \quad i = 0, 1, 2, \quad \text{strongly in } L^1([0, T] \times [0, L]), \\
q_j^\varepsilon & \xrightarrow{\varepsilon \rightarrow 0} u_j \quad j = 1, 2, \quad \text{strongly in } L^1([0, T] \times [0, L]),
\end{aligned}$$

where  $(u_1, u_2, u_0)$  is the unique weak solution of the limit problem (8)–(10) in the sense of Definition 2.4.

The system (7) can be seen as a particular case of the model without epithelial layer introduced and studied in [28] and [27].

A priori estimates uniform with respect to the parameter  $\varepsilon$  (accounting for permeability) are obtained in Section 4. We emphasize that estimates on time derivatives are more subtle due to specific boundary conditions of system and because one has to take care of singular initial layers. Concerning existence and uniqueness of a solution, in previous works [28] and [27], authors proposed a semi-discrete scheme in space in order to show existence. In this work we propose a fixed point theorem giving the same result for any fixed  $\varepsilon > 0$  in Section 3. The advantage of our approach is that we directly work with weak solutions associated to (7). After recalling the definition of weak solution for problem (1), we report below the statement of Theorem 2.7, and we refer to Section 3 for the proof.

**Definition 2.6.** Let  $(u_1^0(x), u_2^0(x), q_1^0(x), q_2^0(x), u_0^0(x)) \in (L^1(0, L) \cap L^\infty(0, L))^5$  and  $u_b(t) \in L^1(0, T) \cap L^\infty(0, T)$ . Let  $\varepsilon > 0$  be fixed. We say that  $U^\varepsilon(t, x) = (u_1^\varepsilon(t, x), u_2^\varepsilon(t, x), q_1^\varepsilon(t, x), q_2^\varepsilon(t, x), u_0^\varepsilon(t, x)) \in L^\infty((0, T); L^1(0, L) \cap L^\infty(0, L))^5$  is a weak solution of system (7) if for all  $\phi = (\phi_1, \phi_2, \phi_3, \phi_4, \phi_5) \in \mathcal{S}_5$ , with

$$\mathcal{S}_5 := \{\phi \in C^1([0, T] \times [0, L])^5, \quad \phi(T, x) = 0, \phi_1(t, L) = \phi_2(t, L), \text{ and } \phi_2(t, 0) = 0\}$$

we have

$$\begin{aligned}
& \int_0^T \int_0^L \left( u_1^\varepsilon (a_1 \partial_t \phi_1 + \alpha \partial_x \phi_1) + \frac{1}{\varepsilon} (q_1^\varepsilon - u_1^\varepsilon) \phi_1 \right) dx dt + \alpha \int_0^T u_b^\varepsilon(t) \phi_1(t, 0) dt \\
& + \int_0^L a_1 u_1^0(x) \phi_1(0, x) dx \\
& + \int_0^T \int_0^L \left( u_2^\varepsilon (a_2 \partial_t \phi_2 - \alpha \partial_x \phi_2) + \frac{1}{\varepsilon} (q_2^\varepsilon - u_2^\varepsilon) \phi_2 \right) dx dt + \int_0^L a_1 u_2^0(x) \phi_2(0, x) dx \\
& + \int_0^T \int_0^L \left( a_3 q_1^\varepsilon (\partial_t \phi_3) + K_1 (u_0^\varepsilon - q_1^\varepsilon) \phi_3 - \frac{1}{\varepsilon} (q_1^\varepsilon - u_1^\varepsilon) \phi_3 \right) dx dt + \int_0^L a_3 q_1^0(x) \phi_3(0, x) dx \\
& + \int_0^T \int_0^L \left( a_4 q_2^\varepsilon (\partial_t \phi_4) + K_2 (u_0^\varepsilon - q_2^\varepsilon) \phi_4 - \frac{1}{\varepsilon} (q_2^\varepsilon - u_2^\varepsilon) \phi_4 - G(q_2^\varepsilon) \phi_4 \right) dx dt \\
& + \int_0^L a_4 q_2^0(x) \phi_4(0, x) dx \\
& + \int_0^T \int_0^L \left( a_0 u_0^\varepsilon (\partial_t \phi_5) + K_1 (q_1^\varepsilon - u_0^\varepsilon) \phi_5 + K_2 (q_2^\varepsilon - u_0^\varepsilon) \phi_5 + G(q_2^\varepsilon) \phi_5 \right) dx dt \\
& + \int_0^L a_0 u_0^0(x) \phi_5(0, x) dx = 0.
\end{aligned} \tag{15}$$

**Theorem 2.7** (Existence). *Under assumptions (4), (5), (6) and for every fixed  $\varepsilon > 0$ , there exists a unique weak solution  $U^\varepsilon$  of the problem (7).*

**3. Proof of the existence result.** We define the Banach space  $\mathbf{B} := (L^1(0, L) \cap L^\infty(0, L))^5$ . We prove existence using the Banach-Picard fixed point theorem (see e.g., [24] for various examples of its application). We consider a time  $T > 0$  (to be chosen later) and the map  $\mathcal{T} : X_T \rightarrow X_T$  with the Banach space  $X_T = L^\infty([0, T]; \mathbf{B})$ , and we denote  $\|\cdot\|_{X_T} = \sup_{t \in (0, T)} \|\cdot\|_{\mathbf{B}}$ . For a given function  $\tilde{U} \in X_T$ , with  $\tilde{U} = (\tilde{u}_1, \tilde{u}_2, \tilde{q}_1, \tilde{q}_2, \tilde{u}_0)$ , we define  $U := \mathcal{T}(\tilde{U})$  solution to the problem :

$$\begin{cases} a_1 \partial_t u_1 + \alpha \partial_x u_1 = \frac{(\tilde{q}_1 - u_1)}{\varepsilon}, \\ a_2 \partial_t u_2 - \alpha \partial_x u_2 = \frac{(\tilde{q}_2 - u_2)}{\varepsilon}, \\ a_3 \partial_t q_1 = \frac{(\tilde{u}_1 - q_1)}{\varepsilon} + K_1(\tilde{u}_0 - q_1), \\ a_4 \partial_t q_2 = \frac{(\tilde{u}_2 - q_2)}{\varepsilon} + K_2(\tilde{u}_0 - q_2) - G(\tilde{q}_2), \\ a_0 \partial_t u_0 = K_1(\tilde{q}_1 - u_0) + K_2(\tilde{q}_2 - u_0) + G(\tilde{q}_2), \end{cases} \tag{16}$$

with initial data  $u_1^0, u_2^0, q_1^0, q_2^0, u_0^0$  in  $L^1(0, L) \cap L^\infty(0, L)$  and with boundary conditions

$$u_1(t, 0) = u_b(t) \geq 0, \quad u_2(t, L) = u_1(t, L), \quad \text{for } t > 0,$$

where  $u_b \in L^1(0, T) \cap L^\infty(0, T)$ .

First we define the solutions of (16) using Duhamel's formula. Under these hypothesis, we may compute  $u_1$  and  $u_2$  with the method of characteristics

$$u_1(t, x) = \begin{cases} u_1^0(x - \frac{\alpha}{a_1}t) e^{-\frac{t}{a_1\varepsilon}} + \frac{1}{a_1\varepsilon} \int_0^t e^{-\frac{t-s}{a_1\varepsilon}} \tilde{q}_1(x - \frac{\alpha}{a_1}(t-s), s) ds, & \text{if } x > \frac{\alpha}{a_1}t, \\ u_b(t - \frac{a_1x}{\alpha}) e^{-\frac{x}{\varepsilon\alpha}} + \frac{1}{\alpha\varepsilon} \int_0^x e^{-\frac{x-y}{\varepsilon\alpha}} \tilde{q}_1(t - \frac{a_1(x-y)}{\alpha}, y) dy, & \text{if } x < \frac{\alpha}{a_1}t, \end{cases} \quad (17)$$

with  $u_1^0(x)$ , and  $u_b(t)$  initial and boundary condition, respectively. We have a similar expression for  $u_2(t, x)$  with  $u_2^0$  instead of  $u_1^0$  and the boundary condition  $u_2(t, L) = u_1(t, L)$  which is well-defined thanks to (17). It reads :

$$u_2(t, x) = \begin{cases} u_2^0(x + \frac{\alpha}{a_2}t) e^{-\frac{t}{a_2\varepsilon}} + \frac{1}{a_2\varepsilon} \int_0^t e^{-\frac{t-s}{a_2\varepsilon}} \tilde{q}_2(s, x + \frac{\alpha}{a_2}(t-s)) ds, & \text{if } x < L - \frac{\alpha}{a_2}t, \\ u_1(t + \frac{a_2(x-L)}{\alpha}) e^{-\frac{x-L}{\alpha\varepsilon}} + \frac{1}{\alpha\varepsilon} \int_x^L e^{-\frac{x-y}{\alpha\varepsilon}} \tilde{q}_2(t + \frac{a_2(x-y)}{\alpha}, y) dy & \text{if } x > L - \frac{\alpha}{a_2}t. \end{cases} \quad (18)$$

Then, for the other unknowns, one simply solves a system of uncoupled ordinary differential equations leading to :

$$\begin{cases} q_1(t, x) = q_1^0(x) e^{-(\frac{1}{\varepsilon} + K_1)\frac{t}{a_3}} + \frac{1}{a_3} \int_0^t e^{-(\frac{1}{\varepsilon} + K_1)\frac{(t-s)}{a_3}} \left( \frac{1}{\varepsilon} \tilde{u}_1 + K_1 \tilde{u}_0 \right)(s, x) ds, \\ q_2(t, x) = q_2^0(x) e^{-(\frac{1}{\varepsilon} + K_2)\frac{t}{a_4}} + \frac{1}{a_4} \int_0^t e^{-(\frac{1}{\varepsilon} + K_2)\frac{(t-s)}{a_4}} \left( \frac{1}{\varepsilon} \tilde{u}_2 + K_1 \tilde{u}_0 - G(\tilde{q}_2) \right)(s, x) ds, \\ u_0(t, x) = u_0^0(x) e^{-(K_1 + K_2)\frac{t}{a_0}} + \frac{1}{a_0} \int_0^t e^{-(K_1 + K_2)\frac{(t-s)}{a_0}} \left( K_1 \tilde{q}_1 + K_2 \tilde{q}_2 + G(\tilde{q}_2) \right)(s, x) ds. \end{cases} \quad (19)$$

Using Theorem B.1, the previous unknowns solve the weak formulation reading :

$$\begin{cases} \int_0^T \int_0^L \left( -u_1(a_1\partial_t + \alpha\partial_x)\varphi_1(t, x) + \frac{1}{\varepsilon}(u_1 - \tilde{q}_1)\varphi_1(t, x) \right) dxdt \\ \quad + a_1 \left[ \int_0^L u_1(t, x)\varphi_1(t, x)dt \right]_{t=0}^{t=T} + \alpha \left[ \int_0^T u_1(t, x)\varphi_1(t, x) \right]_{x=0}^{x=L} = 0, \\ \int_0^T \int_0^L \left( -u_2(a_2\partial_t - \alpha\partial_x)\varphi_2(t, x) + \frac{1}{\varepsilon}(u_2 - \tilde{q}_2)\varphi_2(t, x) \right) dxdt \\ \quad + a_2 \left[ \int_0^L u_2(t, x)\varphi_2(t, x)dt \right]_{t=0}^{t=T} + \alpha \left[ \int_0^T u_2(t, x)\varphi_2(t, x) \right]_{x=0}^{x=L} = 0, \end{cases} \quad (20)$$

for any  $(\varphi_1, \varphi_2) \in C^1([0, T] \times [0, L])^2$ . Using again Theorem B.1 with  $\Phi = |\cdot|$  show that the same holds true for  $|u_1|$  (resp.  $|u_2|$ ) :

$$\begin{aligned} & \int_0^T \int_0^L -|u_1|(a_1\partial_t + \alpha\partial_x)\varphi_1(t, x) + \frac{1}{\varepsilon}(|u_1| - \text{sgn}(u_1)\tilde{q}_1)\varphi_1(t, x) dxdt \\ & \quad + a_1 \left[ \int_0^L |u_1|(t, x)\varphi_1(t, x)dt \right]_{t=0}^{t=T} + \alpha \left[ \int_0^T |u_1|(t, x)\varphi_1(t, x) \right]_{x=0}^{x=L} = 0. \end{aligned} \quad (21)$$

The same holds also for the other unknowns  $(q_i)_{i \in \{1, 2\}}$  and  $u_0$ , since for the ODE part of the system (19) provides directly similar results. In the rest of the paper, each time that we mention that we are multiplying formally by  $\text{sgn}$  each function of system (16) in order to get :

$$\begin{cases} a_1\partial_t|u_1| + \alpha\partial_x|u_1| = \frac{1}{\varepsilon}(\text{sgn}(u_1)\tilde{q}_1 - |u_1|) \\ a_2\partial_t|u_2| - \alpha\partial_x|u_2| = \frac{1}{\varepsilon}(\text{sgn}(u_2)\tilde{q}_2 - |u_2|), \end{cases}$$



we actually mean that these inequalities hold in the previous sense, *i.e.* in the sense of (21). The reader should notice that the stronger regularity of the integrated forms (17), (18) and (19) allows to define these solutions on the boundaries of the domain  $\partial((0, T) \times (0, L))$ . If these would only belong to  $L^\infty((0, T); L^1(0, L))$  they would not make sense. Now the meaning of the formal setting is well defined, we then can proceed by writing that one has :

$$\begin{cases} a_1 \partial_t |u_1| + \alpha \partial_x |u_1| \leq \frac{1}{\varepsilon} (|\tilde{q}_1| - |u_1|) \\ a_2 \partial_t |u_2| - \alpha \partial_x |u_2| \leq \frac{1}{\varepsilon} (|\tilde{q}_2| - |u_2|) \\ a_3 \partial_t |q_1| \leq \frac{1}{\varepsilon} (|\tilde{u}_1| - |q_1|) + K_1 (|\tilde{u}_0| - |q_1|) \\ a_4 \partial_t |q_2| \leq \frac{1}{\varepsilon} (|\tilde{u}_2| - |q_2|) + K_2 (|\tilde{u}_0| - |q_2|) - |G(\tilde{q}_2)| \\ a_0 \partial_t |u_0| \leq K_1 (|\tilde{q}_1| - |u_0|) + K_2 (|\tilde{q}_2| - |u_0|) + |G(\tilde{q}_2)|. \end{cases} \quad (22)$$

We have used the fact that  $\text{sgn}(G(\tilde{q}_2)) = \text{sgn}(\tilde{q}_2)$  from (6), which implies in particular  $-G(\tilde{q}_2) \cdot \text{sgn}(\tilde{q}_2) = -|G(\tilde{q}_2)|$  and  $G(\tilde{q}_2) \cdot \text{sgn}(u_0) \leq |G(\tilde{q}_2)|$ . In order to obtain inequalities in the weak formulation associated to the latter system it is enough to choose non-negative test functions in  $C^1([0, T] \times [0, L])$ .

Adding all equations and integrating on  $[0, L]$ , we obtain formally

$$\begin{aligned} \frac{d}{dt} \int_0^L (a_1 |u_1| + a_2 |u_2| + a_0 |u_0| + a_3 |q_1| + a_4 |q_2|) dx &\leq \alpha |u_1(t, 0)| \\ &+ \frac{1}{\varepsilon} \int_0^L (|\tilde{u}_1| + |\tilde{u}_2| + |\tilde{q}_1| + |\tilde{q}_2|) dx + (K_1 + K_2) \int_0^L |\tilde{u}_0| dx, \end{aligned}$$

where we use the boundary condition  $u_1(t, L) = u_2(t, L)$  and (6). Let us define  $\|U(t, \cdot)\|_{L^1(0, L)^5} := \int_0^L (|u_1| + |u_2| + |q_1| + |q_2| + |u_0|)(t, x) dx$  and integrating with respect to time, we obtain:

$$\begin{aligned} \|U(t, x)\|_{L^1(0, L)^5} &\leq \|U(0, x)\|_{L^1(0, L)^5} + \frac{\alpha}{\min_i a_i} \int_0^T |u_b(s)| ds \\ &+ \eta \int_0^T \|\tilde{U}(t, x)\|_{L^1(0, L)^5} dt, \end{aligned} \quad (23)$$

with  $\eta = \frac{1}{\min_i a_i} (K_1 + K_2 + \frac{1}{\varepsilon}) > 0$ . Here the formal computations are to be understood in the following manner : in the weak formulation associated to (22) we choose the test function  $\varphi = (1, 1, 1, 1, 1)$ , and the result (23) comes in a straightforward way when neglecting the out-coming characteristic at  $x = 0$ .

On the other hand, using (17), (18) and (19), one quickly checks that

$$\|U\|_{L^\infty((0, T) \times (0, L))^5} \leq \max \left( \|U^0\|_{L^\infty(0, L)^5}, \|u_b\|_{L^\infty(0, T)} \right) + \frac{CT}{\varepsilon} \|\tilde{U}\|_{L^\infty((0, T) \times (0, L))^5} \quad (24)$$

where the generic constant  $C$  depends only on  $(a_i)_{i \in \{0, \dots, 4\}}$ ,  $(K_i)_{i \in \{1, 2\}}$  and  $\|G'\|_{L^\infty(\mathbb{R})}$  but not on  $\tilde{U}$  nor on the data  $U^0$ . At this step,  $\mathcal{T}$  maps  $X_T$  into itself.

Let us now prove that  $\mathcal{T}$  is a contraction. Let  $(\tilde{U}, \tilde{W}) \in X_T^2$ , we define  $U := \mathcal{T}(\tilde{U})$ ,  $W := \mathcal{T}(\tilde{W})$ . Then, by the same token as obtaining (23), we have

$$\begin{aligned} \|\mathcal{T}(\tilde{U}) - \mathcal{T}(\tilde{W})\|_{L^1(0,L)^5} &= \|U - W\|_{L^1(0,L)^5} \leq \eta \int_0^T \|\tilde{U} - \tilde{W}\|_{L^1(0,L)} \\ &\leq \eta T \|\tilde{U} - \tilde{W}\|_{X_T}. \end{aligned}$$

Again similar computations as in (24), show that

$$\|U - W\|_{L^\infty((0,T) \times (0,L))^5} \leq \frac{CT}{\varepsilon} \|\tilde{U} - \tilde{W}\|_{L^\infty((0,T) \times (0,L))^5}.$$

Therefore, as soon as  $T < \min(1/\eta, \varepsilon/C)$ ,  $\mathcal{T}$  is a contraction in  $X_T$ . It allows to construct a solution on  $[0, T]$  for  $T$  small enough. The fixed point solves (17) and (19) in an implicit way. Along characteristics solutions have enough regularity to satisfy (7) in a weak sense (20). Choosing then the test functions  $\varphi := (\varphi_i)_{i \in \{1, \dots, 5\}}$  to belong to  $\mathcal{S}_5$  shows that the fixed point is a weak solution in the sense of Definition 2.6. Since the solution  $U(t, x) = (u_1(t, x), u_2(t, x), q_1(t, x), q_2(t, x), u_0(t, x))$  is well defined as on  $\{T\} \times (0, L)$  thanks to regularity arguments stated above,  $U(T, x)$  becomes the initial condition of a new initial boundary problem. Thus, we may iterate this process on  $[T, 2T]$ ,  $[2T, 3T]$ ,  $\dots$ , since the condition on  $T$  does not depend on the iteration.

As a result of above computations, we have also that if  $U^1$  (resp.  $U^2$ ) is a solution with initial data  $U^{1,0}$  (resp.  $U^{2,0}$ ) and boundary data  $u_b^1$  (resp.  $u_b^2$ ). Then we have the comparison principle :

$$\|U^1 - U^2\|_{L^1((0,T) \times (0,L))} \leq \|U^{0,1} - U^{0,2}\|_{L^1(0,L)} + \frac{\alpha}{\min_i a_i} \|u_b^1 - u_b^2\|_{L^1(0,T)}. \quad (25)$$

which shows and implies uniqueness as well.

**4. Uniform *a priori* estimates.** In order to prove our convergence result, we first establish some uniform *a priori* estimates. The strategy of the proof of Theorem 2.5 relies on a compactness argument. In this Section we will omit the superscript  $\varepsilon$  in order to simplify the notation.

**4.1. Non-negativity and  $L^1 \cap L^\infty$  estimates.** The following lemma establishes that all concentrations of system are non-negative and this is consistent with the biological framework.

**Lemma 4.1** (Non-negativity). *Let  $U(t, x)$  be a weak solution of system (1) such that the assumptions (4), (5), (6) hold. Then for almost every  $(t, x) \in (0, T) \times (0, L)$ ,  $U(t, x)$  is non-negative, i.e.:  $u_1(t, x), u_2(t, x), q_1(t, x), q_2(t, x), u_0(t, x) \geq 0$ .*

*Proof.* We prove that the negative part of our functions vanishes. Using Stampacchia's method, we formally multiply each equation of system (7) by corresponding indicator function as follows:

$$\begin{cases} (a_1 \partial_t u_1 + \alpha \partial_x u_1) \mathbf{1}_{\{u_1 < 0\}} = \frac{1}{\varepsilon} (q_1 - u_1) \mathbf{1}_{\{u_1 < 0\}} \\ (a_2 \partial_t u_2 - \alpha \partial_x u_2) \mathbf{1}_{\{u_2 < 0\}} = \frac{1}{\varepsilon} (q_2 - u_2) \mathbf{1}_{\{u_2 < 0\}} \\ (a_3 \partial_t q_1) \mathbf{1}_{\{q_1 < 0\}} = \frac{1}{\varepsilon} (u_1 - q_1) \mathbf{1}_{\{q_1 < 0\}} + K_1 (u_0 - q_1) \mathbf{1}_{\{q_1 < 0\}} \\ (a_4 \partial_t q_2) \mathbf{1}_{\{q_2 < 0\}} = \frac{1}{\varepsilon} (u_2 - q_2) \mathbf{1}_{\{q_2 < 0\}} + K_2 (u_0 - q_2) \mathbf{1}_{\{q_2 < 0\}} - G(q_2) \mathbf{1}_{\{q_2 < 0\}} \\ (a_0 \partial_t u_0) \mathbf{1}_{\{u_0 < 0\}} = K_1 (q_1 - u_0) \mathbf{1}_{\{u_0 < 0\}} + K_2 (q_2 - u_0) \mathbf{1}_{\{u_0 < 0\}} + G(q_2) \mathbf{1}_{\{u_0 < 0\}}. \end{cases}$$

Again as in the proof of existence in Section 3, these computations can be made rigorously using the extra regularity provided along characteristics in the spirit of Lemma 3.1, [20] summarized in Theorem B.1.

Using again Theorem B.1, one writes formally :

$$\begin{cases} a_1 \partial_t u_1^- + \alpha \partial_x u_1^- \leq \frac{1}{\varepsilon} (q_1^- - u_1^-) \\ a_2 \partial_t u_2^- - \alpha \partial_x u_2^- \leq \frac{1}{\varepsilon} (q_2^- - u_2^-) \\ a_3 \partial_t q_1^- \leq \frac{1}{\varepsilon} (u_1^- - q_1^-) + K_1(u_0^- - q_1^-) \\ a_4 \partial_t q_2^- \leq \frac{1}{\varepsilon} (u_2^- - q_2^-) + K_2(u_0^- - q_2^-) + G(q_2) \mathbf{1}_{\{q_2 < 0\}} \\ a_0 \partial_t u_0^- \leq K_1(q_1^- - u_0^-) + K_2(q_2^- - u_0^-) - G(q_2) \mathbf{1}_{\{u_0 < 0\}}. \end{cases}$$

Adding the previous expressions, one recovers a single inequality reading

$$\partial_t (a_1 u_1^- + a_2 u_2^- + a_3 q_1^- + a_4 q_2^- + a_0 u_0^-) + \alpha \partial_x (u_1^- - u_2^-) \leq G(q_2) (\mathbf{1}_{\{q_2 < 0\}} - \mathbf{1}_{\{u_0 < 0\}}).$$

By Assumption 2.3, we have that  $\text{sgn}(G(q_2)) = \text{sgn}(q_2)$ . Thus  $G(q_2)(\mathbf{1}_{\{q_2 < 0\}} - \mathbf{1}_{\{u_0 < 0\}}) = G(q_2) (\mathbf{1}_{\{G(q_2) < 0\}} - \mathbf{1}_{\{u_0 < 0\}}) \leq 0$ . Then integrating on the interval  $[0, L]$ , we get :

$$\begin{aligned} & \frac{d}{dt} \int_0^L (a_1 u_1^- + a_2 u_2^- + a_3 q_1^- + a_4 q_2^- + a_0 u_0^-)(t, x) dx \\ & \leq \alpha (u_2^-(t, L) - u_2^-(t, 0) - u_1^-(t, L) + u_1^-(t, 0)). \end{aligned}$$

Since  $u_1^-(t, L) = u_2^-(t, L)$  thanks to condition (5), it follows:

$$\frac{d}{dt} \int_0^L (a_1 u_1^- + a_2 u_2^- + a_3 q_1^- + a_4 q_2^- + a_0 u_0^-)(t, x) dx \leq \alpha u_1^-(t, 0) = \alpha u_b^-(t).$$

From Assumptions 2.2 and 2.1, the initial and boundary data are all non-negative. Thus  $u_1^-(0, x), q_1^-(0, x), q_2^-(0, x), u_2^-(0, x), u_0^-(0, x)$  are necessarily zero. This proves solutions' non-negativity and concludes the proof.  $\square$

**Lemma 4.2** ( $L^\infty$  bound). *Let  $(u_1, u_2, q_1, q_2, u_0)$  be the unique weak solution of problem (7). Assume that (4), (5), (6) hold, then it is bounded i.e. for a.e.  $(t, x) \in (0, T) \times (0, L)$ ,*

$$0 \leq u_0(t, x) \leq \kappa(1+t), \quad 0 \leq u_i(t, x) \leq \kappa(1+t), \quad 0 \leq q_i(t, x) \leq \kappa(1+t), \quad i = 1, 2,$$

$$0 \leq u_2(t, 0) \leq \kappa(1+t), \quad 0 \leq u_1(t, L) \leq \kappa(1+t),$$

where the constant  $\kappa \geq \max \{ \|G\|_\infty, \|u_b\|_\infty, \|u_0^0\|_\infty, \|u_i^0\|_\infty, \|q_i^0\|_\infty, i \in \{1, 2\} \}$ .

*Proof.* We use the same method as in the previous lemma for the functions

$$w_i = (u_i - \kappa(1+t)), \quad i = 0, 1, 2, \quad z_j = (q_j - \kappa(1+t)), \quad j = 1, 2.$$

From system (7) and using the fact that

$$z_j \mathbf{1}_{\{w_i \geq 0\}} = z_j^+ \mathbf{1}_{\{w_i \geq 0\}} - z_j^- \mathbf{1}_{\{w_i \geq 0\}} \leq z_j^+, \quad w_i \mathbf{1}_{\{z_j \geq 0\}} \leq w_i^+,$$

we get

$$\begin{cases} a_1 \partial_t w_1^+ + a_1 \kappa \mathbf{1}_{\{w_1 \geq 0\}} + \alpha \partial_x w_1^+ \leq \frac{1}{\varepsilon} (z_1^+ - w_1^+) \\ a_2 \partial_t w_2^+ + a_2 \kappa \mathbf{1}_{\{w_2 \geq 0\}} - \alpha \partial_x w_2^+ \leq \frac{1}{\varepsilon} (z_2^+ - w_2^+) \\ a_3 \partial_t z_1^+ + a_3 \kappa \mathbf{1}_{\{z_1 \geq 0\}} \leq \frac{1}{\varepsilon} (w_1^+ - z_1^+) + K_1 (w_0^+ - z_1^+) \\ a_4 \partial_t z_2^+ + a_4 \kappa \mathbf{1}_{\{z_2 \geq 0\}} \leq \frac{1}{\varepsilon} (w_2^+ - z_2^+) + K_2 (w_0^+ - z_2^+) - G(q_2) \mathbf{1}_{\{z_2 \geq 0\}} \\ a_0 \partial_t w_0^+ + a_0 \kappa \mathbf{1}_{\{w_0 \geq 0\}} \leq K_1 (z_1^+ - w_0^+) + K_2 (z_2^+ - w_0^+) + G(q_2) \mathbf{1}_{\{w_0 \geq 0\}}. \end{cases} \quad (26)$$

Adding expressions above gives

$$\begin{aligned} & \partial_t (a_1 w_1^+ + a_2 w_2^+ + a_3 z_1^+ + a_4 z_2^+ + a_0 w_0^+) + \alpha \partial_x (w_1^+ - w_2^+) \\ & \leq -\kappa \mathbf{1}_{\{w_0 \geq 0\}} + G(q_2) (\mathbf{1}_{\{w_0 \geq 0\}} - \mathbf{1}_{\{z_2 \geq 0\}}). \end{aligned}$$

Integrating with respect to  $x$  yields

$$\begin{aligned} & \frac{d}{dt} \int_0^L (a_1 w_1^+ + a_2 w_2^+ + a_3 z_1^+ + a_4 z_2^+ + a_0 w_0^+) (t, x) dx \\ & \leq \alpha (w_2^+(t, L) - w_2^+(t, 0) - w_1^+(t, L) + w_1^+(t, 0)) + \int_0^L (G(q_2) - \kappa) \mathbf{1}_{\{w_0 \geq 0\}} dx, \end{aligned}$$

where we use the fact that  $G(q_2) \geq 0$  from assumption (6) since  $q_2 \geq 0$  thanks to the previous lemma. From the boundary conditions in (1), we have for all  $t \geq 0$ ,  $w_2^+(t, L) = [u_2(t, L) - \kappa(1+t)]^+ = [u_1(t, L) - \kappa(1+t)]^+ = w_1^+(t, L)$ . Then,

$$\begin{aligned} & \frac{d}{dt} \int_0^L (a_1 w_1^+ + a_2 w_2^+ + a_3 z_1^+ + a_4 z_2^+ + a_0 w_0^+) (t, x) dx + \alpha w_2^+(t, 0) \\ & \leq \alpha (u_b(t) - \kappa(1+t))^+ + (\|G\|_\infty - \kappa) \int_0^L \mathbf{1}_{\{w_0 \geq 0\}} dx. \end{aligned}$$

If we adjust the constant  $\kappa$  such that  $\kappa \geq \max \{\|G\|_\infty, \|u_b\|_\infty\}$ , it implies that :

$$\frac{d}{dt} \int_0^L (a_1 w_1^+ + a_2 w_2^+ + a_3 z_1^+ + a_4 z_2^+ + a_0 w_0^+) (t, x) dx + \alpha w_2^+(t, 0) \leq 0,$$

which shows the claim.

For the last estimate on  $u_1(t, L)$ , we sum the first and the third inequalities of the system (26) and integrate on  $(0, L)$ ,

$$\begin{aligned} & \frac{d}{dt} \int_0^L (a_1 w_1^+ + a_3 z_1^+) dx + \alpha w_1^+(t, L) \leq \alpha w_1^+(t, 0) + K_1 \int_0^L (w_0^+ - z_1^+) dx \\ & \quad - \kappa \int_0^L (\mathbf{1}_{\{w_1 \geq 0\}} + \mathbf{1}_{\{z_1 \geq 0\}}) dx. \end{aligned}$$

Integrating on  $(0, T)$  and since we have proved above that  $w_0^+ = 0$  and  $z_1^+ = 0$ , we arrive at

$$\alpha \int_0^T w_1^+(t, L) dt \leq \alpha \int_0^T w_1^+(t, 0) dt = 0,$$

for  $\kappa \geq \|u_b\|_\infty$ . □

**Lemma 4.3** ( $L_t^\infty L_x^1$  estimates). *Let  $T > 0$  and let  $(u_1, u_2, q_1, q_2, u_0)$  be a weak solution of system (1) in  $(L^\infty([0, T]; (L^1 \cap L^\infty)(0, L)))^5$ . We define:*

$$\mathcal{H}(t) = \int_0^L (a_1|u_1| + a_2|u_2| + a_0|u_0| + a_3|q_1| + a_4|q_2|)(t, x) dx.$$

*Then, under hypothesis (4), (5), (6) the following a priori estimate, uniform in  $\varepsilon > 0$ , holds:*

$$\mathcal{H}(t) \leq \alpha \|u_b\|_{L^1(0, T)} + \mathcal{H}(0), \quad \forall t > 0.$$

*Moreover the following inequalities hold:*

$$\int_0^T |u_2(t, 0)| dt \leq \|u_b\|_{L^1(0, T)} + \frac{1}{\alpha} \mathcal{H}(0),$$

*and*

$$\int_0^T |u_1(t, L)| dt \leq \int_0^L (|u_1^0(x)| + |q_1^0(x)|) dx + CT$$

*with  $C > 0$  constant.*

*Proof.* Since from Lemma 4.1 all concentrations are non-negative, we may write from system (7)

$$\begin{cases} a_1 \partial_t |u_1| + \alpha \partial_x |u_1| = \frac{1}{\varepsilon} (|q_1| - |u_1|) \\ a_2 \partial_t |u_2| - \alpha \partial_x |u_2| = \frac{1}{\varepsilon} (|q_2| - |u_2|) \\ a_3 \partial_t |q_1| = \frac{1}{\varepsilon} (|u_1| - |q_1|) + K_1 (|u_0| - |q_1|) \\ a_4 \partial_t |q_2| = \frac{1}{\varepsilon} (|u_2| - |q_2|) + K_2 (|u_0| - |q_2|) - |G(q_2)| \\ a_0 \partial_t |u_0| = K_1 (|q_1| - |u_0|) + K_2 (|q_2| - |u_0|) + |G(q_2)|. \end{cases} \quad (27)$$

Adding all equations and integrating on  $(0, L)$ , we get, recalling the boundary condition  $u_1(t, L) = u_2(t, L)$ ,

$$\frac{d}{dt} \mathcal{H}(t) + \alpha |u_2(t, 0)| = \alpha |u_1(t, 0)| = \alpha |u_b(t)|. \quad (28)$$

Integrating now with respect to time, we obtain:

$$\mathcal{H}(t) + \alpha \int_0^t |u_2(s, 0)| ds \leq \alpha \int_0^t |u_b(s)| ds + \mathcal{H}(0). \quad (29)$$

with  $\mathcal{H}(t)$  previously defined. It gives the first two estimates of the Lemma. Finally, to obtain the last inequality, we add equations (7a) and (7c) and integrate on  $(0, L)$  to get

$$\frac{d}{dt} \int_0^L (a_1|u_1| + a_3|q_1|) dx + \alpha |u_1(t, L)| \leq \alpha |u_b(t)| + K_1 \int_0^L |u_0| dx.$$

Since we have shown that  $\int_0^L |u_0| dx \leq \frac{1}{a_0} \mathcal{H}(t) < \infty$ , we can conclude after integrating with respect to time.  $\square$

## 4.2. Estimates on the derivatives.

4.2.1. *Data regularization.* Here we detail the notion of regularization for BV functions. The  $C^\infty((0, L))$ -regularization denoted  $f_\delta$  for a  $BV(0, L)$  function  $f$  follows the steps below [32, Theorem 5.3.3], *i.e.*

$$f_\delta(x) = \sum_{k=0}^{\infty} \eta_{\delta_k} * (f \zeta_k)$$

where  $\{\zeta_k\}_{k=1}^{\infty}$  is a sequence of regular cut-off functions s.t.

$$\begin{cases} \zeta_k \in C_c^\infty(V_k), \quad 0 \leq \zeta_k \leq 1, \quad (k = 1, 2, \dots) \\ \sum_{k=1}^{\infty} \zeta_k \equiv 1 \text{ on } (0, L) \end{cases}$$

where

$$\begin{cases} \Theta_0 = \emptyset, \\ \Theta_k := \left\{ x \in (0, L) ; \text{dist}(x, \{0, L\}) > \frac{1}{k} \right\}, \quad (k = 1, 2, \dots) \\ V_k := \Theta_{k+1} \setminus \overline{\Theta}_{k-1}, \quad (k = 1, 2, \dots) \end{cases}$$

and there exists  $\delta_k > 0$  such that

$$\begin{cases} \text{Supp}(\eta_{\delta_k} * (\zeta_k f)) \subseteq V_k, \\ \int_{(0, L)} |\eta_{\delta_k} * (\zeta_k f) - \zeta_k f| dx \leq \delta 2^{-k}, \\ \int_{(0, L)} |\eta_{\delta_k} * (\zeta'_k f) - \zeta'_k f| dx \leq \delta 2^{-k}, \end{cases}$$

Although  $f_\delta$  converges strongly to  $f$  in  $L^1(0, L)$ , one has only that

$$\lim_{\delta \rightarrow 0} \|\partial_x f_\delta\|_{L^1(0, L)} = \|\lambda_f((0, L))\|,$$

where the right hand side is the total variation of the Radon measure associated to the derivative of  $f$  [32, Theorem 5.3.3]. This result comes partly from the estimates from above :

$$\|\partial_x f_\delta\|_{L^1(0, L)} \leq \|f\|_{BV(0, L)}. \quad (30)$$

Then we set

$$f_c^\delta(x) := (1 - \chi_\delta(x))f_\delta(x) + c\chi_\delta(x), \quad \forall x \in (0, L) \quad (31)$$

where  $c$  is a given real constant and

$$\chi_\delta(x) := \chi\left(\frac{x}{\delta}\right), \quad \chi(x) := \begin{cases} 1 & \text{if } |x| < 1 \\ 0 & \text{if } |x| > 2 \end{cases},$$

$\chi \in C^\infty(\mathbb{R})$  being a positive monotone function.

**Lemma 4.4.** *If  $f \in BV(0, L)$ ,  $c \in \mathbb{R}$  and  $f_c^\delta$  defined as above, one has :*

$$\|\partial_x f_c^\delta\|_{L^1(0, L)} \leq C \left(1 + \|f\|_{BV(0, L)}\right)$$

where the generic constant  $C(c, \chi)$  is independent of  $\delta$ .

*Proof.* Differentiating (31) and integrating in space gives :

$$\begin{aligned}
\|\partial_x f_c^\delta\|_{L^1(0,L)} &\leq 2 \|\partial_x f^\delta\|_{L^1(0,L)} + \frac{1}{\delta} \int_0^\delta |f^\delta(x) - c| |\chi'(x/\delta)| dx \\
&\leq 2 \|\partial_x f^\delta\|_{L^1(0,L)} + \|\chi'\|_{L^\infty(\mathbb{R})} \left( |f^\delta(0^+) - c| + \frac{1}{\delta} \int_0^\delta |f^\delta(x) - f^\delta(0^+)| dx \right) \\
&\leq C \left( \|f\|_{\text{BV}(0,L)} + |f^\delta(0^+) - c| \right) \\
&\leq C \left( \|f\|_{\text{BV}(0,L)} + c + |f^\delta(0^+)| \right) \leq C \left( \|f\|_{\text{BV}(0,L)} + c + \|f^\delta\|_{W^{1,1}(0,L)} \right) \\
&\leq C \left( 1 + \|f\|_{\text{BV}(0,L)} \right)
\end{aligned}$$

where we used (30).  $\square$

**Definition 4.5.** If  $(u_1^0, u_2^0, q_1^0, q_2^0, u_0^0)$  and  $u_b$  are respectively the initial and boundary data associated to the problem (7), under hypotheses 4 and 5, we define as **regular data** their regularization in the following manner : we set

$$\begin{cases}
u_1^{0,\delta}(x) := (1 - \chi_\delta(x) - \chi_\delta(L - x))u_{1,\delta}^0 + c_1\chi_\delta(x) + c_2\chi_\delta(L - x), & \forall x \in [0, L] \\
u_b^\delta(t) := (1 - \chi_\delta(t))u_{b,\delta} + c_1\chi_\delta(t), & \forall t \in [0, T] \\
u_2^{0,\delta}(x) := (1 - \chi_\delta(L - x))u_{2,\delta}^0 + c_2\chi_\delta(L - x), & \forall x \in [0, L] \\
q_1^{0,\delta}(x) := q_{1,\delta}^0, & \forall x \in [0, L] \\
q_2^{0,\delta}(x) := q_{2,\delta}^0, & \forall x \in [0, L] \\
u_0^{0,\delta}(x) := u_{0,\delta}^0, & \forall x \in [0, L]
\end{cases}$$

where  $u_{1,\delta}^0$  is the approximation introduced above, the same notation holding for the rest of the initial data.

The matching is  $C^\infty$  in the neighborhood of  $(0,0)$  in  $[0, L] \times [0, T]$ . Indeed,  $u_1^{0,\delta}(x) = c_1$  when  $x$  is close enough to 0 and in the same way  $u_b^\delta(t) = c_1$  when  $t$  is near 0, whereas for the derivatives  $(u_1^{0,\delta})^{(k)}(x) = 0$  when  $x$  is close to zero for any derivative of order  $k$ , and the same holds for  $(u_b^\delta)^{(k)}(t)$  when  $t$  is sufficiently small. The same holds true in the neighborhood of the point  $(t, x) = (L, 0)$ .

This regularization procedure allows then to obtain

**Lemma 4.6.** Assume hypotheses 2.3, and let  $U^\delta$  be the solution associated to problem (7) with initial data  $U^{0,\delta} = (u_1^{0,\delta}, u_2^{0,\delta}, q_1^{0,\delta}, q_2^{0,\delta}, u_0^{0,\delta})$  and the boundary condition  $u_b^\delta$ . Then  $\partial_t U^\delta$  belongs to  $X_T := L^\infty((0, T); (L^1(0, L) \cap L^\infty(0, L)))^5$ . and solves the problem

$$\begin{cases}
a_1 \partial_t + \alpha \partial_x u_{1,t}^\delta = \frac{1}{\varepsilon} (q_{1,t}^\delta - u_{1,t}^\delta) \\
a_2 \partial_t - \alpha \partial_x u_{2,t}^\delta = \frac{1}{\varepsilon} (q_{2,t}^\delta - u_{2,t}^\delta) \\
a_3 \partial_t q_{1,t}^\delta = -\frac{1}{\varepsilon} (q_{1,t}^\delta - u_{1,t}^\delta) + K_1(u_{0,t}^\delta - q_{1,t}^\delta) \\
a_4 \partial_t q_{2,t}^\delta = -\frac{1}{\varepsilon} (q_{2,t}^\delta - u_{2,t}^\delta) + K_2(u_{0,t}^\delta - q_{2,t}^\delta) - G'(q_2^\delta) q_{2,t}^\delta \\
a_0 \partial_t u_{0,t}^\delta = K_1(q_{1,t}^\delta - u_{0,t}^\delta) + K_2(q_{2,t}^\delta - u_{0,t}^\delta) - G'(q_2^\delta) q_{2,t}^\delta
\end{cases} \quad (32)$$

where  $u_{i,t}^\delta = \partial_t u_i^\delta$  and so on.

$$\begin{cases} u_{1,t}^\delta(t, 0) = \partial_t u_b^\delta(t), \\ a_1 u_{1,t}^\delta(0, x) = -\alpha \partial_x u_1^{0,\delta} + \frac{1}{\varepsilon} (q_1^{0,\delta} - u_1^{0,\delta}) \\ a_2 u_{2,t}^\delta(0, x) = \alpha \partial_x u_2^{0,\delta} + \frac{1}{\varepsilon} (q_2^{0,\delta} - u_2^{0,\delta}) \\ a_3 q_{1,t}^\delta(0, x) = -\frac{1}{\varepsilon} (q_1^{0,\delta} - u_1^{0,\delta}) + K_1(u_0^{0,\delta} - q_1^{0,\delta}) \\ a_4 q_{2,t}^\delta(0, x) = -\frac{1}{\varepsilon} (q_2^{0,\delta} - u_2^{0,\delta}) + K_2(u_0^{0,\delta} - q_2^{0,\delta}) - G(q_2^{0,\delta}) \\ a_0 u_{0,t}^\delta(0, x) = K_1(q_1^{0,\delta} - u_0^{0,\delta}) + K_2(q_2^{0,\delta} - u_0^{0,\delta}) + G(q_2^{0,\delta}) \end{cases} \quad (33)$$

*Proof.* The Duhamel's formula obtained by the fixed point method in the proof of Theorem 2.7 provides a solution  $U^\delta \in X_T$ . Deriving  $U^\delta$  with respect to  $t$ , one can show that  $\partial_t U^\delta$  solves (32) with initial and boundary conditions (33). Applying then the existence result again proves that actually  $\partial_t U^\delta$  belongs to  $X_T$ .  $\square$

**Remark 1.** *A priori* estimates from previous sections, when applied to the problem (32) complemented with initial-boundary data (33), do not provide a control of  $\partial_t U^\delta$  which is uniform with respect to  $\varepsilon$ .

This remark motivates next paragraphs.

4.2.2. *The initial layer.* At this step we introduce initial layer correctors. For this sake, on the microscopic scale we define for  $t \in \mathbb{R}_+$ , functions  $(\tilde{u}_1, \tilde{u}_2, \tilde{q}_1, \tilde{q}_2)$  solving

$$\begin{cases} a_1 \partial_t \tilde{u}_1 = \tilde{q}_1 - \tilde{u}_1 \\ a_2 \partial_t \tilde{u}_2 = \tilde{q}_2 - \tilde{u}_2 \\ a_3 \partial_t \tilde{q}_1 = \tilde{u}_1 - \tilde{q}_1 \\ a_4 \partial_t \tilde{q}_2 = \tilde{u}_2 - \tilde{q}_2, \end{cases} \quad (34)$$

with initial conditions

$$\tilde{u}_1(0, x) = q_1^{0,\delta} - u_1^{0,\delta}, \quad \tilde{u}_2(0, x) = q_2^{0,\delta} - u_2^{0,\delta}, \quad \tilde{q}_1(0, x) = 0, \quad \tilde{q}_2(0, x) = 0.$$

Actually, this system may be solved explicitly and we obtain for  $i \in \{1, 2\}$ ,

$$\begin{cases} \tilde{u}_i(t, x) = \frac{a_i + a_{i+2} \exp\left(-t\left(\frac{1}{a_i} + \frac{1}{a_{i+2}}\right)\right)}{a_i + a_{i+2}} (q_i^{0,\delta}(x) - u_i^{0,\delta}(x)) \\ \tilde{q}_i(t, x) = \frac{a_i \left(1 - \exp\left(-t\left(\frac{1}{a_i} + \frac{1}{a_{i+2}}\right)\right)\right)}{a_i + a_{i+2}} (q_i^{0,\delta}(x) - u_i^{0,\delta}(x)) \end{cases} \quad (35)$$

4.2.3. *Uniform  $L^1$  bounds of the time derivatives.* We introduce the following quantities on the macroscopic time scale  $t \in [0, T]$  :

$$\begin{cases} v_1^\delta(t, x) = u_1^\delta(t, x) + \tilde{u}_1\left(\frac{t}{\varepsilon}, x\right) \\ v_2^\delta(t, x) = u_2^\delta(t, x) + \tilde{u}_2\left(\frac{t}{\varepsilon}, x\right) \\ s_1^\delta(t, x) = q_1^\delta(t, x) + \tilde{q}_1\left(\frac{t}{\varepsilon}, x\right) \\ s_2^\delta(t, x) = q_2^\delta(t, x) + \tilde{q}_2\left(\frac{t}{\varepsilon}, x\right) \end{cases} \quad (36)$$

Next, we prove uniform bounds on the time derivatives :



**Proposition 1.** *Let  $T > 0$ . If the data is regular in the sense of Definition 4.5, setting :*

$$\tilde{\mathcal{H}}_t(t) = \int_0^L (a_1|\partial_t v_1^\delta| + a_2|\partial_t v_2^\delta| + a_3|\partial_t s_1^\delta| + a_4|\partial_t s_2^\delta| + a_0|\partial_t u_0^\delta|)(t, x) dx,$$

with functions  $v_1, v_2, s_1^\delta, s_2^\delta$  defined in (36), one has :

$$\begin{aligned} \tilde{\mathcal{H}}_t(t) + \alpha \int_0^t |\partial_t v_2^\delta(\tau, 0)| d\tau + \alpha \int_0^t |\partial_t v_1^\delta(\tau, L)| d\tau \\ \leq C \left( \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)} + \|u_b^\delta\|_{W^{1,1}(0,T)} \right), \quad \text{for a.e. } t \in (0, T), \end{aligned} \quad (37)$$

where  $\mathbf{W}^{1,1}(0, L)$  denotes the vector-space  $W^{1,1}(0, L)^5$ .

*Proof.* From system (7) we deduce

$$\begin{cases} a_1 \partial_t v_1^\delta + \alpha \partial_x v_1^\delta = \frac{1}{\varepsilon} (s_1^\delta - v_1^\delta) + \alpha \partial_x \tilde{u}_1(\frac{t}{\varepsilon}, x) \\ a_2 \partial_t v_2^\delta - \alpha \partial_x v_2^\delta = \frac{1}{\varepsilon} (s_2^\delta - v_2^\delta) - \alpha \partial_x \tilde{u}_2(\frac{t}{\varepsilon}, x) \\ a_3 \partial_t s_1^\delta = \frac{1}{\varepsilon} (v_1^\delta - s_1^\delta) + K_1(u_0^\delta - s_1^\delta) + K_1 \tilde{q}_1(\frac{t}{\varepsilon}, x) \\ a_4 \partial_t s_2^\delta = \frac{1}{\varepsilon} (v_2^\delta - s_2^\delta) + K_2(u_0^\delta - s_2^\delta) + K_2 \tilde{q}_2(\frac{t}{\varepsilon}, x) - G(q_2^\delta) \\ a_0 \partial_t u_0^\delta = K_1(s_1^\delta - u_0^\delta) + K_2(s_2^\delta - u_0^\delta) - K_1 \tilde{q}_1(\frac{t}{\varepsilon}, x) - K_2 \tilde{q}_2(\frac{t}{\varepsilon}, x) + G(q_2^\delta) \end{cases} \quad (38)$$

with following initial and boundary conditions:

$$\begin{aligned} v_1^\delta(t, 0) &= u_1(t, 0) + \tilde{u}_1(t, 0) = u_b(t) + \tilde{u}_1(\frac{t}{\varepsilon}, 0), \quad t \in (0, T), \\ v_2^\delta(t, L) &= v_2^\delta(t, L) + \tilde{u}_2(\frac{t}{\varepsilon}, L), \quad t \in (0, T), \\ v_1^\delta(0, x) &= u_1^\delta(0, x) + \tilde{u}_1(0, x) = q_1^0(x), \quad x \in (0, L), \\ v_2^\delta(0, x) &= u_2^\delta(0, x) + \tilde{u}_2(0, x) = q_2^0(x), \\ s_1^\delta(0, x) &= q_1^\delta(0, x) + \tilde{q}_1(0, x) = q_1^0(x), \\ s_2^\delta(0, x) &= q_2^\delta(0, x) + \tilde{q}_2(0, x) = q_2^0(x). \end{aligned} \quad (39)$$

As  $G \in C^2(\mathbb{R})$ , thanks to Lemma 4.6,  $\partial_t U^\delta \in L^\infty((0, T); (L^1(0, T) \cap L^\infty(0, L)))^5$ , taking the derivative with respect to  $t$  in system (38),  $\partial_t V^\delta = \partial_t(v_1^\delta, v_2^\delta, s_1^\delta, s_2^\delta, u_0^\delta)$  solves

$$\begin{cases} a_1 \partial_t v_{1,t}^\delta + \alpha \partial_x v_{1,t}^\delta = \frac{1}{\varepsilon} (s_{1,t}^\delta - v_{1,t}^\delta) + \frac{1}{\varepsilon} \partial_x \tilde{u}_{1,t} \\ a_2 \partial_t v_{2,t}^\delta - \alpha \partial_x v_{2,t}^\delta = \frac{1}{\varepsilon} (s_{2,t}^\delta - v_{2,t}^\delta) - \frac{1}{\varepsilon} \partial_x \tilde{u}_{2,t} \\ a_3 \partial_t s_{1,t}^\delta = \frac{1}{\varepsilon} (v_{1,t}^\delta - s_{1,t}^\delta) + K_1(u_{0,t}^\delta - s_{1,t}^\delta) + \frac{1}{\varepsilon} K_1 \tilde{q}_{1,t} \\ a_4 \partial_t s_{2,t}^\delta = \frac{1}{\varepsilon} (v_{2,t}^\delta - s_{2,t}^\delta) + K_2(u_{0,t}^\delta - s_{2,t}^\delta) + \frac{1}{\varepsilon} K_2 \tilde{q}_{2,t} - G'(q_2^\delta) q_{2,t} \\ a_0 \partial_t u_{0,t}^\delta = K_1(s_{1,t}^\delta - u_{0,t}^\delta) + K_2(s_{2,t}^\delta - u_{0,t}^\delta) - \frac{1}{\varepsilon} K_1 \tilde{q}_{1,t} - \frac{1}{\varepsilon} K_2 \tilde{q}_{2,t} + G'(q_2^\delta) q_{2,t} \end{cases}$$

in the sense of Definition (2.6). Again formally, we multiply each equation respectively by  $\text{sgn}(v_{i,t}^\delta)$  with  $i = 1, 2$ , and  $\text{sgn}(s_{j,t}^\delta)$ , for  $j = 1, 2$ , and by  $\text{sgn}(u_{0,t}^\delta)$  in the sense explained in the proof of Theorem 2.7. It gives

$$\begin{cases} a_1 \partial_t |v_{1,t}^\delta| + \alpha \partial_x |v_{1,t}^\delta| \leq \frac{1}{\varepsilon} (|s_{1,t}^\delta| - |v_{1,t}^\delta|) + \frac{1}{\varepsilon} \partial_x |\tilde{u}_{1,t}| \\ a_2 \partial_t |v_{2,t}^\delta| - \alpha \partial_x |v_{2,t}^\delta| \leq \frac{1}{\varepsilon} (|s_{2,t}^\delta| - |v_{2,t}^\delta|) + \frac{1}{\varepsilon} \partial_x |\tilde{u}_{2,t}| \\ a_3 \partial_t |s_{1,t}^\delta| \leq \frac{1}{\varepsilon} (|v_{1,t}^\delta| - |s_{1,t}^\delta|) + K_1(|u_{0,t}^\delta| - |s_{1,t}^\delta|) + \frac{1}{\varepsilon} K_1 |\tilde{q}_{1,t}| \\ a_4 \partial_t |s_{2,t}^\delta| \leq \frac{1}{\varepsilon} (|v_{2,t}^\delta| - |s_{2,t}^\delta|) + K_2(|u_{0,t}^\delta| - |s_{2,t}^\delta|) + \frac{1}{\varepsilon} K_2 |\tilde{q}_{2,t}| \\ \quad + |G'(q_2^\delta) \frac{1}{\varepsilon} \tilde{q}_{2,t}| - G'(q_2^\delta) |s_{2,t}^\delta| \\ a_0 \partial_t |u_{0,t}^\delta| \leq K_1(|s_{1,t}^\delta| - |u_{0,t}^\delta|) + K_2(|s_{2,t}^\delta| - |u_{0,t}^\delta|) + \frac{1}{\varepsilon} K_1 |\tilde{q}_{1,t}| \\ \quad + \frac{1}{\varepsilon} K_2 |\tilde{q}_{2,t}| + |G'(q_2^\delta) \frac{1}{\varepsilon} \tilde{q}_{2,t}| + G'(q_2^\delta) |s_{2,t}^\delta|. \end{cases} \quad (40)$$

Indeed, the right hand side of the 4th and 5th inequalities can be obtained as follows. On the one hand, we have

$$\begin{aligned} -G'(q_2^\delta)q_{2,t}\operatorname{sgn}(s_{2,t}^\delta) &= -G'(q_2^\delta)\left(s_{2,t}^\delta(t,x) - \frac{1}{\varepsilon}\tilde{q}_{2,t}\left(\frac{t}{\varepsilon},x\right)\right)\operatorname{sgn}(s_{2,t}^\delta) \\ &\leq -G'(q_2^\delta)|s_{2,t}^\delta| + \frac{1}{\varepsilon}|G'(q_2^\delta)\tilde{q}_{2,t}|. \end{aligned}$$

On the other hand

$$\begin{aligned} -G'(q_2^\delta)q_{2,t}\operatorname{sgn}(u_{0,t}^\delta) &= -G'(q_2^\delta)\left(s_{2,t}^\delta(t,x) - \frac{1}{\varepsilon}\tilde{q}_{2,t}\left(\frac{t}{\varepsilon},x\right)\right)\operatorname{sgn}(u_{0,t}^\delta) \\ &\leq G'(q_2^\delta)|s_{2,t}^\delta| + \frac{1}{\varepsilon}|G'(q_2^\delta)\tilde{q}_{2,t}|, \end{aligned}$$

since  $G$  is non-decreasing by assumption (6). Summing all inequalities in (40) and integrating with respect to space on  $(0, L)$ , we obtain formally

$$\frac{d}{dt}\tilde{\mathcal{H}}_t(t) + \alpha|v_{2,t}^\delta(t,0)| \leq F_1(t) + F_2(t) + F_3(t) + F_4(t), \quad (41)$$

where

$$\begin{aligned} F_1(t) &:= \alpha(|v_{2,t}^\delta(t,L)| - |v_{1,t}^\delta(t,L)|); \quad F_2(t) := \alpha|v_{1,t}^\delta(t,0)|, \\ F_3(t) &:= \frac{1}{\varepsilon} \int_0^L \left| \partial_x \tilde{u}_{2,t}\left(\frac{t}{\varepsilon},x\right) \right| dx + \frac{1}{\varepsilon} \int_0^L \left| \partial_x \tilde{u}_{1,t}\left(\frac{t}{\varepsilon},x\right) \right| dx, \\ F_4(t) &:= \frac{2K_1}{\varepsilon} \int_0^L \left| \tilde{q}_{1,t}\left(\frac{t}{\varepsilon},x\right) \right| dx + \frac{2}{\varepsilon}(\|G'\|_\infty + K_2) \int_0^L \left| \tilde{q}_{2,t}\left(\frac{t}{\varepsilon},x\right) \right| dx. \end{aligned}$$

Integrating (41) in time, we get

$$\tilde{\mathcal{H}}_t(t) + \alpha \int_0^T |v_{2,t}^\delta(t,0)| dt \leq \int_0^T (F_1(t) + F_2(t) + F_3(t) + F_4(t)) dt + \tilde{\mathcal{H}}_t(0). \quad (42)$$

Let us consider each term of the right hand side of (42) separately:

- $F_1$ : On the right boundary  $x = L$ , one has

$$\begin{aligned} \int_0^T (|v_{2,t}^\delta(t,L)| - |v_{1,t}^\delta(t,L)|) dt &\leq \frac{1}{\varepsilon} \int_0^T \left| (\tilde{u}_{2,t} - \tilde{u}_{1,t})\left(\frac{t}{\varepsilon},L\right) \right| dt \\ &\leq \int_0^{\frac{T}{\varepsilon}} |(\tilde{u}_{2,t} - \tilde{u}_{1,t})(\tau,L)| d\tau \leq \frac{1}{2} \left\{ |u_1^{0,\delta}(0) - q_1^{0,\delta}(0)| + |u_2^{0,\delta}(0) - q_2^{0,\delta}(0)| \right\} \\ &\leq C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)} \end{aligned}$$

where we used trace operator's continuity for  $W^{1,1}(0, L)$  functions (using integration by parts, one shows as well that the constant  $C$  does not depend on  $\delta$ ).

- $F_2$ : On the other hand at  $x = 0$ , the boundary condition can be estimated as

$$\begin{aligned} \int_0^T |v_{1,t}^\delta(t,0)| dt &\leq \int_0^T |(u_b^\delta)'(s)| ds + \int_0^{\frac{T}{\varepsilon}} |(u_1^{0,\delta}(0) - q_1^{0,\delta}(0))e^{-2\tau}| d\tau \\ &\leq C \left( \|u_b^\delta\|_{W^{1,1}(0,T)} + \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)} \right) \end{aligned}$$

as above.

- $F_3$ : With the change of variable  $\tau = \frac{t}{\varepsilon}$ , we have, using again (35),

$$\int_0^T \int_0^L \left| \frac{1}{\varepsilon} \partial_x \tilde{u}_{i,t} \left( \frac{t}{\varepsilon}, x \right) \right| dx dt = \int_0^{\frac{T}{\varepsilon}} \int_0^L |\partial_x \tilde{u}_{i,t}(\tau, x)| dx d\tau \leq C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)},$$

which is uniformly bounded with respect to  $\varepsilon$ .

- $F_4$ : similarly, we have

$$\int_0^T \int_0^L \left| \frac{1}{\varepsilon} \tilde{q}_{i,t} \left( \frac{t}{\varepsilon}, x \right) \right| dx dt = \int_0^{\frac{T}{\varepsilon}} \int_0^L |\tilde{q}_{i,t}(\tau, x)| dx d\tau \leq C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)}$$

thanks to the fact that  $\partial_t \tilde{q}_i(\tau, x) = (q_i^0(x) - u_i^0(x))e^{-2\tau}$  with  $i = 1, 2$ .

It remains to estimate  $\tilde{\mathcal{H}}_t(0)$  in (42). Indeed, using (38) at  $t = 0$  in order to convert time derivatives into expressions involving only the data and its space derivatives, one obtains

$$\begin{aligned} \tilde{\mathcal{H}}_t(0) &= \int_0^L \left( a_1 |v_{1,t}^\delta(0, x)| + a_2 |v_{2,t}^\delta(0, x)| + a_3 |s_{1,t}^\delta(0, x)| + a_4 |s_{2,t}^\delta(0, x)| + a_0 |u_{0,t}^\delta(0, x)| \right) dx \\ &\leq C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)} \end{aligned}$$

So, for instance, for the first term of the sum, we use the first equation in (38) and we write

$$a_1 \partial_t v_1^\delta(0, x) = \frac{1}{\varepsilon} (s_1^\delta(0, x) - v_1^\delta(0, x)) + \alpha \partial_x \tilde{u}_1(0, x) - \alpha \partial_x v_1^\delta(0, x).$$

Recalling that  $s_1^\delta(0, x) = q_1^0(x)$  and  $v_1^\delta(0, x) = q_1^0(x)$  as defined in (39) we get:  $a_1 \partial_t v_1^\delta(0, x) = \alpha \partial_x (q_1^0(x) - u_1^0(x)) - \alpha \partial_x v_1^\delta(0, x) = -\alpha \partial_x u_1^0(x)$ , then

$$\int_0^L |\partial_t v_1^\delta(0, x)| dx \leq \frac{\alpha}{a_1} \int_0^L |\partial_x u_1^0(x)| dx < C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)},$$

The rest follows exactly the same way. We conclude from (42) and the above calculations that

$$\begin{aligned} \tilde{\mathcal{H}}_t(t) + \alpha \int_0^T |v_{2,t}^\delta(t, 0)| dt &\leq \tilde{\mathcal{H}}_t(0) + \int_0^t (F_1 + F_2 + F_3 + F_4)(s) ds \\ &\leq C \left( \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)} + \|u_b^\delta\|_{W^{1,1}(0,T)} \right). \end{aligned}$$

Finally, in order to recover (37), we add the first and third inequalities in (40) and integrate on  $(0, T) \times (0, L)$ , we get

$$\begin{aligned} &\int_0^L (a_1 |v_{1,t}^\delta(T, x)| + a_3 |s_{1,t}^\delta(T, x)|) dx + \alpha \int_0^T |v_{1,t}^\delta(t, L)| dt \\ &\leq \alpha \int_0^T |v_{1,t}^\delta(t, 0)| dt + \int_0^L (a_1 |v_{1,t}^\delta(0, x)| + a_3 |s_{1,t}^\delta(0, x)|) dx \\ &\quad + \int_0^T \int_0^L \left( K_1 |u_{0,t}^\delta(t, x)| + \frac{K_1}{\varepsilon} \left| \tilde{q}_{1,t} \left( \frac{t}{\varepsilon}, x \right) \right| + \frac{1}{\varepsilon} \left| \partial_x \tilde{u}_{1,t} \left( \frac{t}{\varepsilon}, x \right) \right| \right) dx dt. \end{aligned}$$

We have already proved that the second term of the right hand side is bounded. We have also proved above that  $u_{0,t}^\delta$  is uniformly bounded in  $L^\infty((0, T); L^1(0, L))$ . From (39), we have  $v_{1,t}^\delta(t, 0) = u_b'(t) + \frac{1}{\varepsilon} \tilde{u}_{1,t}(\frac{t}{\varepsilon}, 0)$ . As above, we use the expressions of  $\tilde{u}_1$  and  $\tilde{q}_1$  and a change of variable to bound each term of the right hand side.  $\square$

As a consequence, we deduce the following estimates on the time derivatives of the original unknowns  $(u_1^\delta, u_2^\delta, q_1^\delta, q_2^\delta, u_0^\delta)$ :

**Corollary 1.** *Let  $T > 0$ , under the same assumptions, there exists a constant  $C_T > 0$  depending only on the  $W^{1,1}$  norm of the data but independent on  $\varepsilon$ , such that :*

$$\begin{aligned} \int_0^T \int_0^L (|\partial_t u_1^\delta| + |\partial_t u_2^\delta| + |\partial_t u_0^\delta| + |\partial_t q_1^\delta| + |\partial_t q_2^\delta|)(t, x) dx dt &\leq C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)}, \\ \int_0^T |\partial_t u_2^\delta(t, 0)| dt &\leq C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)}, \quad \int_0^T |\partial_t u_1^\delta(t, L)| dt \leq C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)}. \end{aligned} \quad (43)$$

*Proof.* We recall the expressions

$$v_1^\delta = u_1^\delta + \tilde{u}_1, \quad v_2^\delta = u_2^\delta + \tilde{u}_2, \quad s_1^\delta = q_1^\delta + \tilde{q}_1, \quad s_2^\delta = q_2^\delta + \tilde{q}_2.$$

By the triangle inequality, we have for  $i \in \{1, 2\}$ ,

$$\begin{aligned} \|\partial_t u_i^\delta\|_{L^1([0,T] \times [0,L])} &\leq \|\partial_t v_i^\delta\|_{L^1([0,T] \times [0,L])} + \frac{1}{\varepsilon} \|\partial_t \tilde{u}_i(t/\varepsilon, x)\|_{L^1([0,T] \times [0,L])}, \\ \|\partial_t q_i^\delta\|_{L^1([0,T] \times [0,L])} &\leq \|\partial_t s_i^\delta\|_{L^1([0,T] \times [0,L])} + \frac{1}{\varepsilon} \|\partial_t \tilde{q}_i(t/\varepsilon, x)\|_{L^1([0,T] \times [0,L])}. \end{aligned}$$

The first terms of the latter right hand side are bounded from Proposition 1. For the second terms, we have, as above,

$$\begin{aligned} \int_0^T \int_0^L \frac{1}{\varepsilon} \left| \partial_t \tilde{q}_i \left( \frac{t}{\varepsilon}, x \right) \right| dx dt &= \frac{1}{\varepsilon} \int_0^T \int_0^L \left| (q_i^0(x) - u_i^0(x)) e^{-2\frac{t}{\varepsilon}} \right| dx dt \\ &< C \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)}, \\ \int_0^T \int_0^L \frac{1}{\varepsilon} \left| \partial_t \tilde{u}_i \left( \frac{t}{\varepsilon}, x \right) \right| dx dt &= \frac{1}{\varepsilon} \int_0^T \int_0^L \left| (u_i^0(x) - q_i^0(x)) e^{-2\frac{t}{\varepsilon}} \right| dx dt \\ &< C' \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)}. \end{aligned}$$

Furthermore, from (42), we get

$$\int_0^T |v_{2,t}^\delta(t, 0)| dt \leq C_T \|U^{0,\delta}\|_{\mathbf{W}^{1,1}(0,L)}.$$

By the triangle inequality, it implies the second estimate in (43)

To recover the third claim in (43), we notice that by definition of  $v_1^\delta$  and a triangle inequality, we have

$$|u_{1,t}^\delta(t, L)| \leq |v_{1,t}^\delta(t, L)| + \frac{1}{\varepsilon} \left| \tilde{u}_{1,t} \left( \frac{t}{\varepsilon}, L \right) \right| \leq |v_{1,t}^\delta(t, L)| + \frac{1}{\varepsilon} \|q_1^{0,\delta} - u_1^{0,\delta}\|_{W^{1,1}(0,L)} e^{-2\frac{t}{\varepsilon}}.$$

where again we use the continuity of the trace operator on  $W^{1,1}(0, L)$  functions in order to recover the dependence between the values at  $x = L$  and the  $W^{1,1}(0, L)$ -norm of the initial data. Integrating in time and using (37) allows to conclude.  $\square$

#### 4.2.4. Uniform bounds on the space derivatives.

**Lemma 4.7.** *Let  $T > 0$ . If the data is regular in the sense of Definition 4.5, then, the space derivatives of functions  $u_1^\delta, u_2^\delta$  satisfy the following estimates :*

$$\int_0^T \int_0^L \left( \sum_{i=0}^2 |\partial_x u_i^\delta(t, x)| + \sum_{i=1}^2 |\partial_x q_i^\delta(t, x)| \right) dx dt \leq C_T \|U^{0,\delta}\|_{W^{1,1}((0,L))},$$

for some non-negative constant  $C_T$  uniformly bounded with respect to  $\varepsilon$ .

*Proof.* Adding equation (7a) with (7c) and also (7b) with (7d) we get

$$\begin{aligned}\alpha \partial_x u_1^\delta &= K_1(u_0^\delta - q_1^\delta) - a_1 \partial_t u_1^\delta - a_3 \partial_t q_1^\delta, \\ -\alpha \partial_x u_2^\delta &= K_2(u_0^\delta - q_2^\delta) - a_2 \partial_t u_2^\delta - a_4 \partial_t q_2^\delta - G(q_2^\delta).\end{aligned}$$

Using Corollary 1 and (6), the right hand sides are uniformly bounded in  $L^1((0, T) \times (0, L))$ . Deriving the ODE part of (7) with respect to the space variable and using the latter estimates provides the results for  $u_0^\delta$ ,  $q_1^\delta$  and  $q_2^\delta$ .  $\square$

**4.3. Extension to BV data.** We show here how to use Corollary 1 and Lemma 4.7 in order to obtain BV compactness.

**Theorem 4.8.** *Under hypotheses (2.1)-(2.3), there exists a uniform bound such that the  $\varepsilon$ -dependent solutions of system (7) satisfy*

$$\sum_{i=0}^2 \|u_i\|_{BV((0,T) \times (0,L))} + \sum_{i=1}^2 \|q_i\|_{BV((0,T) \times (0,L))} \leq C \left( \|U^0\|_{\mathbf{BV}(0,L)} + \|u_b\|_{BV(0,T)} \right)$$

where the generic constant  $C$  is independent on  $\varepsilon$ .

*Proof.* Setting  $U^\delta(t, x) := (u_1^\delta(t, x), u_2^\delta(t, x), q_1^\delta(t, x), q_2^\delta(t, x), u_0^\delta(t, x))$ , one has from the previous estimates :

$$\|U^\delta\|_{\mathbf{W}_{t,x}^{1,1}((0,T) \times (0,L))} \leq C \left( \|U^{0,\delta}\|_{\mathbf{W}_x^{1,1}(0,L)} + \|u_b^\delta\|_{W^{1,1}(0,T)} \right)$$

Now thanks to Lemma 4.4, one estimates the right hand side with respect to the BV norm of the data :

$$\|U^{0,\delta}\|_{\mathbf{W}_x^{1,1}(0,L)} \leq C \left( 1 + \|U^0\|_{\mathbf{BV}(0,L)} \right), \quad \|u_b^\delta\|_{W^{1,1}(0,T)} \leq C(1 + \|u_b\|_{BV(0,T)})$$

We are in the hypotheses of [32, Theorem 5.2.1. p. 222] : by  $L^1$ -continuity, shown in Theorem 2.7,  $U^\delta$  tends to  $U := (u_1, u_2, q_1, q_2, u_0)$  in  $L^1((0, T) \times (0, L))$  strongly, when  $\delta$  vanishes. Then for any open set  $V \subset (0, T) \times (0, L)$  one has

$$\|U\|_{\mathbf{BV}_{t,x}(V)} \leq \liminf_{\delta \rightarrow 0} \|U^\delta\|_{\mathbf{BV}_{t,x}((0,T) \times (0,L))} = \liminf_{\delta \rightarrow 0} \|U^\delta\|_{\mathbf{W}_{t,x}^{1,1}((0,T) \times (0,L))}$$

and since the  $\mathbf{BV}$  bound of the sequence  $(U^\delta)_\delta$  is uniformly bounded with respect to  $\delta$ , the result extends by [32, Remark 5.2.2. p. 223] to the whole set  $(0, T) \times (0, L)$ .  $\square$

**5. Proof of the convergence result (Theorem 2.5).** It is divided into two steps.

- 1. Convergence :** From Lemma 4.2, Lemma 4.3 and Corollary 1, sequences  $(u_1^\varepsilon)_\varepsilon$  and  $(u_2^\varepsilon)_\varepsilon$  are uniformly bounded in  $L^\infty \cap BV((0, T) \times (0, L))$ . Thanks to the Helly's theorem [3, 18], we deduce that, up to extraction of a subsequence,

$$\begin{aligned}u_1^\varepsilon &\xrightarrow{\varepsilon \rightarrow 0} u_1 \text{ strongly in } L^1([0, T] \times [0, L]), \\ u_2^\varepsilon &\xrightarrow{\varepsilon \rightarrow 0} u_2 \text{ strongly in } L^1([0, T] \times [0, L]),\end{aligned}$$

with limit function  $u_1, u_2 \in L^\infty \cap BV((0, T) \times (0, L))$ .

By equations (7a), one shows by testing with the appropriate  $C^1$  compactly supported functions in  $(0, T) \times (0, L)$  and using the definition of the BV norm (cf for instance, [32, p. 220–221]):

$$\|q_1^\varepsilon - u_1^\varepsilon\|_{L^1((0,T) \times (0,L))} \leq C\varepsilon \|u_1^\varepsilon\|_{BV((0,T) \times (0,L))}$$

which tends to zero as  $\varepsilon$  goes to 0 thanks to the bounds in Corollary 1 and Lemma 4.7. Therefore,  $q_1^\varepsilon \xrightarrow{\varepsilon \rightarrow 0} u_1$  strongly in  $L^1((0, T) \times (0, L))$ . By the same argument, we show that  $q_2^\varepsilon \xrightarrow{\varepsilon \rightarrow 0} u_2$  strongly in  $L^1((0, T) \times (0, L))$ . Moreover, since  $G$  is Lipschitz-continuous, we have, when  $\varepsilon$  goes to zero,

$$\|G(q_2^\varepsilon) - G(u_2)\|_{L^1((0, T) \times (0, L))} \longrightarrow 0.$$

For the convergence of  $u_0^\varepsilon$ , let us first denote  $u_0$  a solution to the equation

$$a_0 \partial_t u_0 = K_1(u_1 - u_0) + K_2(u_2 - u_0) + G(u_2).$$

Then, taking the last equation of system (7), subtracting by this latter equation and multiplying by  $\text{sgn}(u_0^\varepsilon - u_0)$ , we get in a weak sense that

$$\begin{aligned} a_0 \partial_t |u_0^\varepsilon - u_0| &\leq K_1 |q_1^\varepsilon - u_1| + K_2 |q_2^\varepsilon - u_2| + (K_1 + K_2) |u_0 - u_0^\varepsilon| + |G(q_2^\varepsilon) - G(u_2)| \\ &\leq K_1 |q_1^\varepsilon - u_1| + K_2 |q_2^\varepsilon - u_2| + (K_1 + K_2) |u_0 - u_0^\varepsilon| + \|G'\|_\infty |q_2^\varepsilon - u_2|. \end{aligned}$$

Using Grönwall's Lemma, we get, after an integration on  $[0, L]$ ,

$$\begin{aligned} \int_0^L |u_0^\varepsilon - u_0|(t, x) dx &\leq \int_0^L e^{\frac{(K_1 + K_2)}{a_0} t} |u_0 - u_0^\varepsilon|(0, x) dx \\ &\quad + \frac{K_1}{a_0} \int_0^L \int_0^T e^{\frac{(K_1 + K_2)}{a_0} (t-s)} |q_1^\varepsilon - u_1|(s, x) ds dx \\ &\quad + \frac{(\|G'\|_\infty + K_2)}{a_0} \int_0^L \int_0^T e^{\frac{(K_1 + K_2)}{a_0} (t-s)} |q_2^\varepsilon - u_2|(s, x) ds dx. \end{aligned}$$

Thus, one concludes that

$$u_0^\varepsilon \xrightarrow{\varepsilon \rightarrow 0} u_0 \text{ strongly in } L^1((0, T) \times (0, L)).$$

## 2. The limit system :

We pass to the limit in (15), the weak formulation of system (7). Suppose that  $\phi \in \mathcal{S}_5$ . Taking  $\phi_1 = \phi_3$  and  $\phi_2 = \phi_4$  in (15) we may pass to the limit  $\varepsilon \rightarrow 0$  and we obtain

$$\begin{aligned} &\int_0^T \int_0^L u_1((a_1 + a_3) \partial_t \phi_1 + \alpha \partial_x \phi_1) dx dt + \alpha \int_0^T u_b(t) \phi_1(t, 0) dt \\ &+ \int_0^L (a_1 u_1^0(x) + a_3 q_1^0(x)) \phi_1(0, x) dx \\ &+ \int_0^T \int_0^L u_2((a_2 + a_4) \partial_t \phi_2 - \alpha \partial_x \phi_2) dx dt + \int_0^L (a_2 u_2^0(x) + a_4 q_2^0(x)) \phi_2(0, x) dx \\ &+ \int_0^T \int_0^L (a_0 u_0 \partial_t \phi_3 + K_1(u_1 - u_0)(\phi_3 - \phi_1) + K_2(u_2 - u_0)(\phi_3 - \phi_2) \\ &+ G(u_2)(\phi_3 - \phi_2)) dx dt + \int_0^L a_0 u_0(0, x) \phi_3(0, x) dx = 0. \end{aligned}$$

which is exactly (14) with initial data coming from system (7). Finally, since the solution of the limit system is unique, we deduce that the whole sequence converges. This concludes the proof of Theorem 2.5.

**6. Numerical validation.** For the sake of simplicity, we assume, in what follows that  $a_i = 1$ , for  $i \in \{0, 4\}$  when we discretise problems (7) and (8)-(10). We discretize the domain  $(0, L)$  with a uniform mesh in space of size  $N_x$  setting :

$$\Delta x := L/N_x,$$

In order to avoid an  $\varepsilon$ -dependent CFL, we discretize the linear parts of the source term in (7) in an implicit way, while we keep the transport part explicit and use a MUSCL second order discretization in space [8]. This leads to write the scheme, for  $i \in \{1, N_x - 1\}$

$$\left\{ \begin{array}{l} \left(1 + \frac{\Delta t}{\varepsilon}\right) u_{i,1}^{n+1} - \frac{\Delta t}{\varepsilon} q_{i,1}^{n+1} \\ \quad = (1 - \lambda_5) u_{i,1}^n + \lambda_5 u_{i-1,1}^n - \frac{\lambda_5(1 - \lambda_5)}{2} (\phi_{i+1/2,1}^n - \phi_{i-1/2,1}^n) \\ \left(1 + \frac{\Delta t}{\varepsilon}\right) u_{i,2}^{n+1} - \frac{\Delta t}{\varepsilon} q_{i,2}^{n+1} \\ \quad = (1 - \lambda_5) u_{i,2}^n + \lambda_5 u_{i+1,2}^n - \frac{\lambda_5(1 - \lambda_5)}{2} (\phi_{i+1/2,2}^n - \phi_{i-1/2,2}^n) \\ -\frac{\Delta t}{\varepsilon} u_{i,1}^{n+1} + \left(1 + \frac{\Delta t}{\varepsilon} + \Delta t K_1\right) q_{i,1}^{n+1} - \Delta t K_1 u_{i,0}^{n+1} = q_{i,1}^n \\ -\frac{\Delta t}{\varepsilon} u_{i,2}^{n+1} + \left(1 + \frac{\Delta t}{\varepsilon} + \Delta t K_2\right) q_{i,2}^{n+1} - \Delta t K_2 u_{i,0}^{n+1} = q_{i,2}^n - \Delta t G(q_{i,2}^n) \\ -\Delta t K_1 q_{i,1}^{n+1} - \Delta t K_2 q_{i,2}^{n+1} + (1 + \Delta t(K_1 + K_2)) u_{i,0}^{n+1} = u_{i,0}^n + \Delta t G(q_{i,2}^n) \end{array} \right. \quad (44)$$

where the time step is chosen s.t.

$$\lambda_5 := \frac{\Delta t}{\Delta x} \alpha$$

is less than 1, and the flux limiters are defined as

$$\begin{aligned} \phi_{i+1/2,1}^n &:= \text{minmod}(u_{i+1,1}^n - u_{i,1}^n, u_{i,1}^n - u_{i-1,1}^n), \\ \phi_{i+1/2,2}^n &:= \text{minmod}(u_{i+2,2}^n - u_{i+1,2}^n, u_{i+1,2}^n - u_{i,2}^n), \end{aligned}$$

and the minmod function is defined in a Generalised minmod limiter [8] :

$$\text{minmod}(1, r) := \max\left(0, \min\left(2r, \frac{1+r}{2}, 2\right)\right).$$

We complement the scheme with initial and boundary conditions :

$$u_{i,j}^0 := \frac{1}{\Delta x} \int_{i\Delta x}^{(i+1)\Delta x} u_j^0, \quad \forall j \in \{0, 1, 2\}, \quad q_{i,j}^0 := \frac{1}{\Delta x} \int_{i\Delta x}^{(i+1)\Delta x} q_j^0, \quad \forall j \in \{1, 2\}$$

and

$$u_{0,1}^{n+1} := u_n^b = \frac{1}{\Delta t} \int_{n\Delta t}^{(n+1)\Delta t} u_b(t) dt, \quad u_{N_x,2}^{n+1} := u_{N_x-1,1}^{n+1}, \quad \forall n \in \{0, N_t - 1\}. \quad (45)$$

**Proposition 2.** *Under the CFL condition :  $\lambda_5 \in (0, 1]$  which is independent of  $\varepsilon$ , the scheme (44) preserves positivity and the discrete version of the  $L^\infty((0, T); L^\infty \cap L^1((0, L)))$ -norm. Moreover, under the same condition, it is possible to prove that*

$$\Delta t \sum_{n=1}^{N_t-1} \sum_{i=1}^{N_x-1} \left\{ \sum_{j=0}^2 |u_{i,j}^{n+1} - u_{i,j}^n| + \sum_{j=1}^2 |q_{i,j}^{n+1} - q_{i,j}^n| \right\} < C$$

For the first part, the proof uses at the discrete level similar ideas as in Lemmas 4.1 and 4.2. For the second part, one adds the initial layer (19) to the numerical solutions for each time step and proceeds as in Proposition 1 and Lemma 1. Since the ideas are very similar and the properties of the transport part are rather standard once it is written in a convex form [8, Harten's Lemma], the proof is left to the reader for the sake of concision.

For various values of  $\varepsilon$ , we compare the solutions provided by (44) with a similar discretisation of (8)-(10), reading :

$$\left\{ \begin{array}{l} \left(1 + \frac{\Delta t K_1}{2}\right) v_{i,1}^{n+1} - \frac{\Delta t K_1}{2} v_{i,0}^{n+1} \\ \quad = (1 - \lambda_3) v_{i,1}^n + \lambda_3 v_{i-1,1}^n - \frac{\lambda_3(1 - \lambda_3)}{4} (\phi_{i+1/2,1}^n - \phi_{i-1/2,1}^n) \\ \left(1 + \frac{\Delta t K_2}{2}\right) v_{i,2}^{n+1} - \frac{\Delta t K_2}{2} v_{i,0}^{n+1} \\ \quad = (1 - \lambda_3) v_{i,2}^n + \lambda_3 v_{i+1,2}^n - \frac{\lambda_3(1 - \lambda_3)}{4} (\phi_{i+1/2,2}^n - \phi_{i-1/2,2}^n) - \Delta t G(v_{i,1}^n) \\ -\Delta t K_1 v_{i,1}^{n+1} - \Delta t K_2 v_{i,2}^{n+1} + (1 + \Delta t(K_1 + K_2)) v_{i,0}^{n+1} = v_{i,0}^n + \Delta t G(v_{i,1}^n) \end{array} \right.$$

where

$$\lambda_3 := \frac{\Delta t}{2\Delta x} \alpha = \frac{\lambda_5}{2}$$

We initialize the data setting :

$$v_{i,1}^0 = \frac{u_{i,1}^0 + q_{i,1}^0}{2}, \quad v_{i,2}^0 = \frac{u_{i,2}^0 + q_{i,2}^0}{2}, \quad v_{i,0}^0 = u_{i,0}^0.$$

and boundary conditions are similar to (45). We compare  $(u_{i,j}^n)_{n,i,j}$  and  $(v_{i,j}^n)_{n,i,j}$  using the discrete  $L^1$  norm in space and time. Indeed we compute

$$E_{\varepsilon, \Delta x} := \Delta t \Delta x \sum_{n=0}^{N_t} \sum_{i=0}^{N_x} \left\{ \sum_{j=1}^2 |2v_{i,j}^n - (u_{i,j}^n + q_{i,j}^n)| + |v_{i,0}^n - u_{i,0}^n| \right\},$$

the results are displayed in Fig. 2.

**Appendix A. Stationary system.** Although the study of the rate of convergence for the dynamical system is quite complicated, explicit computations may be performed for the stationary system. This system reads, for  $x \in (0, L)$ ,

$$\alpha \partial_x \bar{u}_1 = \frac{1}{\varepsilon} (\bar{q}_1 - \bar{u}_1) \tag{46a}$$

$$- \alpha \partial_x \bar{u}_2 = \frac{1}{\varepsilon} (\bar{q}_2 - \bar{u}_2) \tag{46b}$$

$$0 = \frac{1}{\varepsilon} (\bar{u}_1 - \bar{q}_1) + K_1 (\bar{u}_0 - \bar{q}_1) \tag{46c}$$

$$0 = \frac{1}{\varepsilon} (\bar{u}_2 - \bar{q}_2) + K_2 (\bar{u}_0 - \bar{q}_2) - G(\bar{q}_2) \tag{46d}$$

$$0 = K_1 (\bar{q}_1 - \bar{u}_0) + K_2 (\bar{q}_2 - \bar{u}_0) + G(\bar{q}_2), \tag{46e}$$

completed with the boundary conditions

$$\bar{u}_1(0) = u_b, \quad \bar{u}_2(L) = \bar{u}_1(L).$$






FIGURE 2. Numerical error estimates

This system may be reduced by noticing that, after adding the equations in (46), we obtain

$$\alpha \partial_x (\bar{u}_1 - \bar{u}_2) = 0.$$

From the boundary condition at  $x = L$ , we deduce that  $\bar{u}_1 = \bar{u}_2$ , we denote  $\bar{u} := \bar{u}_1 = \bar{u}_2$ . With (46a)–(46b), we get

$$2\bar{u} = \bar{q}_1 + \bar{q}_2. \quad (47)$$

From (46e), we deduce

$$\bar{u}_0 = \frac{K_1 \bar{q}_1 + K_2 \bar{q}_2}{K_1 + K_2} + \frac{G(\bar{q}_2)}{K_1 + K_2}. \quad (48)$$

Injecting this latter expression and (47) into (46c), we obtain

$$\bar{q}_1 \left( 1 + 2\varepsilon \frac{K_1 K_2}{K_1 + K_2} \right) = \frac{2\varepsilon K_1}{K_1 + K_2} (K_2 \bar{q}_2 + G(\bar{q}_2)) + \bar{q}_2.$$

With (46e), we deduce

$$\bar{u} = \bar{q}_2 + \frac{\varepsilon K_1}{K_1 + K_2 + 2\varepsilon K_1 K_2} G(\bar{q}_2). \quad (49)$$

Using also (46a), we obtain the following Cauchy problem

$$\begin{cases} \alpha \left( K_1 + K_2 + \varepsilon (2K_1 K_2 + K_1 G'(\bar{q}_2)) \right) \partial_x \bar{q}_2 = K_1 G(\bar{q}_2), \\ \bar{q}_2(x=0) = \bar{q}_2^0, \quad \text{with } \bar{q}_2^0 + \frac{\varepsilon K_1}{K_1 + K_2 + 2\varepsilon K_1 K_2} G(\bar{q}_2^0) = u_b. \end{cases} \quad (50)$$

Solving this system, we deduce  $\bar{q}_2$ , then we compute  $\bar{u}$ ,  $\bar{q}_1$ , and  $\bar{u}_0$  thanks to relations (49), (47), and (48).

When,  $\varepsilon = 0$ , system (50) reads

$$\begin{cases} \alpha (K_1 + K_2) \partial_x \bar{q}_{2,0} = K_1 G(\bar{q}_{2,0}), \\ \bar{q}_{2,0}(x=0) = u_b. \end{cases} \quad (51)$$

In order to have semi-explicit expression, we introduce an antiderivative of  $1/G$  denoted  $\mathcal{G}$ , i.e.  $\mathcal{G}' = \frac{1}{G}$ . Then, integrating (50) we get

$$\begin{aligned} & \alpha(K_1 + K_2 + 2\varepsilon K_1 K_2) \mathcal{G}(\bar{q}_2) + \alpha\varepsilon \ln \mathcal{G}(\bar{q}_2) \\ &= K_1 x + \alpha(K_1 + K_2 + 2\varepsilon K_1 K_2) \mathcal{G}(\bar{q}_2^0) + \alpha\varepsilon \ln \mathcal{G}(\bar{q}_2^0). \end{aligned}$$

By the same token on (51), we have

$$\alpha(K_1 + K_2) \mathcal{G}(\bar{q}_{2,0}) = K_1 x + \alpha(K_1 + K_2) \mathcal{G}(u_b).$$

Subtracting the last two equalities, we obtain

$$\begin{aligned} (K_1 + K_2)(\mathcal{G}(\bar{q}_2) - \mathcal{G}(\bar{q}_{2,0})) &= (K_1 + K_2)(\mathcal{G}(\bar{q}_2^0) - \mathcal{G}(u_b)) + 2\varepsilon K_1 K_2 (\mathcal{G}(\bar{q}_2^0) - \mathcal{G}(\bar{q}_2)) \\ &\quad + \varepsilon(\ln \mathcal{G}(\bar{q}_2^0) - \ln \mathcal{G}(\bar{q}_2)). \end{aligned}$$

Thanks to a Taylor expansion, there exist  $\zeta_2$  between  $\bar{q}_2$  and  $\bar{q}_{2,0}$  and  $\zeta_2^0$  between  $\bar{q}_2^0$  and  $u_b$  such that

$$\begin{aligned} \bar{q}_2 - \bar{q}_{2,0} &= \frac{G(\zeta_2)}{G(\zeta_2^0)} (\bar{q}_2^0 - u_b) + 2\varepsilon K_1 K_2 (\mathcal{G}(\bar{q}_2^0) - \mathcal{G}(\bar{q}_2)) + \varepsilon(\ln \mathcal{G}(\bar{q}_2^0) - \ln \mathcal{G}(\bar{q}_2)) \\ &= -\frac{\varepsilon K_1 G(\zeta_2) G(\bar{q}_2^0)}{(K_1 + K_2 + 2\varepsilon K_1 K_2) G(\zeta_2^0)} + 2\varepsilon K_1 K_2 (\mathcal{G}(\bar{q}_2^0) - \mathcal{G}(\bar{q}_2)) \\ &\quad + \varepsilon(\ln \mathcal{G}(\bar{q}_2^0) - \ln \mathcal{G}(\bar{q}_2)), \end{aligned}$$

where we use (50) for the last equality. Hence we conclude that, formally, the convergence of  $\bar{q}_2$  towards  $\bar{q}_2^0$  is of order  $\varepsilon$ .

**Appendix B. Mild solutions of transport equations.** We define the problem :

$$\begin{cases} \partial_t u + \alpha \partial_x u = \omega(w - u) & (t, x) \in (0, T) \times (0, L) \\ \partial_t v - \alpha \partial_x v = \omega(z - v) & (t, x) \in (0, T) \times (0, L) \\ u(t, 0) = u_b(t), \quad v(t, L) = v_b(t) & t \in (0, T) \\ u(0, x) = u_0(x), \quad v(0, x) = v_0(x) & (t, x) \in \{0\} \times (0, L) \end{cases} \quad (52)$$

where the data  $(w, z, u_0, v_0, u_b, v_b)$  satisfy

$$(w, z) \in L^\infty((0, T) \times (0, L))^2, \quad (u_0, v_0) \in L^\infty(0, L)^2, \quad (u_b, v_b) \in L^\infty(0, T)^2,$$

and  $(\alpha, \omega) \in (\mathbb{R}_+)^2$ . We define by Duhamel's formula

$$u(t, x) := \begin{cases} u_b(t - x/\alpha) \exp(-\omega x/\alpha) + \\ \quad + \omega \int_{-x/\alpha}^0 \exp(-\omega(s + x/\alpha)) w(t + s, x + \alpha s) ds & \text{if } t > x/\alpha \\ u_0(x - \alpha t) \exp(-\omega t) + \omega \int_{-t}^0 \exp(-\omega(s + t)) w(t + s, x + \alpha s) ds & \text{otherwise} \end{cases} \quad (53)$$

and  $v$  is similarly defined as a function of  $(v_0, v_b, z)$ . Then one has, as in [20, Theorem 2.1, Lemma 2.1 and Lemma 3.1], that

**Theorem B.1.** *Let  $(u, v)$  be solutions of (52) defined in the sense of characteristics through Duhamel's formula (53), then  $u$  (resp.  $v$ ) satisfies (52) in the weak sense :*

$$\begin{aligned} \int_0^T \int_0^L \Phi(u(t, x)) (-\partial_t - \partial_x + \omega) \varphi \, dx \, dt + \left[ \int_0^L \Phi(u(t, x)) \varphi(t, x) \, dx \right]_{t=0}^{t=T} \\ + \left[ \int_0^T \Phi(u(t, x)) \varphi(t, x) \, dt \right]_{x=0}^{x=L} = \int_0^T \Phi'(u(t, x)) \omega w(t, x) \varphi(t, x) \, dx \, dt \end{aligned}$$

for all  $\varphi \in C^\infty([0, T] \times [0, L])$  and every  $\Phi \in \text{Lip}(\mathbb{R})$ .

**Acknowledgments.** The authors acknowledge the unknown referees for their comments and remarks which help in improving the paper.

## REFERENCES

- [1] J. F. Bertram, R. N. Douglas-Denton, B. Diouf et al., [Human nephron number: Implications for health and disease](#), *Pediatr. Nephrol.*, **26** (2011), 1529–1533.
- [2] J. S. Clemmer, W. A. Pruett, T. G. Coleman, J. E. Hall and R. L. Hester, [Mechanisms of blood pressure salt sensitivity: New insights from mathematical modeling](#), *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, **312** (2016), R451–R466.
- [3] C. M. Dafermos, [Hyperbolic Conservation Laws in Continuum Physics](#), 4th edition, volume **325**, Berlin: Springer, 2016.
- [4] A. Edwards, M. Auberson, S. Ramakrishnan and O. Bonny, A model of uric acid transport in the rat proximal tubule, *Am. J. Physiol. Renal Physiol.*, (2019), F934–F947.
- [5] B. C. Fry, A. Edwards and A. T. Layton, [Impact of nitric-oxide-mediated vasodilation and oxidative stress on renal medullary oxygenation: A modeling study](#), *Am. J. Physiol. Renal Physiol.*, **310** (2016), F237–F247.
- [6] V. Giovangigli, Z.-B. Yang and W.-A. Yong, [Relaxation limit and initial-layers for a class of hyperbolic-parabolic systems](#), *SIAM J. Math. Anal.*, **50** (2018), 4655–4697.
- [7] E. Giusti, [Minimal Surfaces and Functions of Bounded Variation](#), volume **80**. Birkhäuser/Springer, Basel, 1984.
- [8] E. Godlewski and P.-A. Raviart, [Hyperbolic Systems of Conservation Laws](#), volume **3/4**. Paris: Ellipses, 1991.
- [9] K. Hallow and Y. Gebremichael, [A quantitative systems physiology model of renal function and blood pressure regulation: Application in salt-sensitive hypertension](#), *CPT Pharmacometrics Syst. Pharmacol.*, **6** (2017), 393–400.
- [10] K. W. Hargitay B, [The multiplication principle as the basis for concentrating urine in the kidney](#), *Journal of the American Society of Nephrology*, **12** (2001), 1566–1586.
- [11] M. Heida, R. I. A. Patterson and D. R. M. Renger, [Topologies and measures on the space of functions of bounded variation taking values in a Banach or metric space](#), *J. Evol. Equ.*, **19** (2019), 111–152.
- [12] F. James, [Convergence results for some conservation laws with a reflux boundary condition and a relaxation term arising in chemical engineering](#), *SIAM J. Math. Anal.*, **29** (1998), 1200–1223.
- [13] S. Jin and Z. Xin, [The relaxation schemes for systems of conservation laws in arbitrary space dimensions](#), *Commun. Pure Appl. Math.*, **48** (1995), 235–276.
- [14] A. T. Layton, V. Vallon and A. Edwards, [A computational model for simulating solute transport and oxygen consumption along the nephrons](#), *Am. J. Physiol. Renal Physiol.*, **311** (2016), F1378–F1390.
- [15] A. T. Layton, [Mathematical modeling of kidney transport](#), *Wiley Interdiscip. Rev. Syst. Biol. Med.*, **5** (2013), 557–573.
- [16] A. T. Layton and A. Edwards, [Mathematical Modeling in Renal Physiology](#), Springer, 2014.
- [17] H. E. Layton, [Distribution of henle's loops may enhance urine concentrating capability](#), *Biophys. J.*, **49** (1986), 1033–1040.
- [18] P. G. LeFloch, [Hyperbolic Systems of Conservation Laws. The Theory of Classical and Non-classical Shock Waves](#), Basel: Birkhäuser, 2002.

- [19] M. Marulli, A. Edwards, V. Milišić and N. Vauchelet, [On the role of the epithelium in a model of sodium exchange in renal tubules](#), *Math. Biosci.*, **321** (2020), 108308, 12 pp.
- [20] V. Milišić and D. Oelz, [On the asymptotic regime of a model for friction mediated by transient elastic linkages](#), *J. Math. Pures Appl. (9)*, **96** (2011), 484–501.
- [21] R. Moss and S. R. Thomas, [Hormonal regulation of salt and water excretion: A mathematical model of whole kidney function and pressure natriuresis](#), *Am. J. Physiol. Renal Physiol.*, **306** (2014), F224–F248.
- [22] R. Natalini and A. Terracina, [Convergence of a relaxation approximation to a boundary value problem for conservation laws](#), *Comm. Partial Differential Equations*, **26** (2001), 1235–1252.
- [23] A. Nieves-Gonzalez, C. Clausen, A. Layton, H. Layton and L. Moore, Transport efficiency and workload distribution in a mathematical model of the thick ascending limb, *Am. J. Physiol. Renal Physiol.*, **304** (2012), F653–F664.
- [24] B. Perthame, *Transport Equations Biology*, Basel: Birkhäuser, 2007.
- [25] B. Perthame, N. Seguin and M. Tournus, [A simple derivation of BV bounds for inhomogeneous relaxation systems](#), *Commun. Math. Sci.*, **13** (2015), 577–586.
- [26] J. L. Stephenson, Urinary Concentration and Dilution: Models, *American Cancer Society*, (2011), 1349–1408.
- [27] M. Tournus, *Modèles D'échanges Ioniques Dans le Rein: Théorie, Analyse Asymptotique et Applications Numériques*, PhD thesis, Université Pierre et Marie Curie, France, 2013.
- [28] M. Tournus, A. Edwards, N. Seguin and B. Perthame, [Analysis of a simplified model of the urine concentration mechanism](#), *Netw. Heterog. Media*, **7** (2012), 989–1018.
- [29] M. Tournus, N. Seguin, B. Perthame, S. R. Thomas and A. Edwards, [A model of calcium transport along the rat nephron](#), *Am. J. Physiol. Renal Physiol.*, **305** (2013), F979–F994.
- [30] A. M. Weinstein, [A mathematical model of the rat nephron: Glucose transport](#), *Am. J. Physiol. Renal Physiol.*, **308** (2015), F1098–F1118.
- [31] A. M. Weinstein, [A mathematical model of the rat kidney:  \$K^+\$ -induced natriuresis](#), *Am. J. Physiol. Renal Physiol.*, **312** (2017), F925–F950.
- [32] W. P. Ziemer, [Weakly Differentiable Functions. Sobolev Spaces and Functions of Bounded Variation](#), Graduate Texts in Mathematics, **120**. Springer-Verlag, New York, 1989.

Received June 2020; revised April 2021; early access July 2021.

E-mail address: [marta.marulli3@unibo.it](mailto:marta.marulli3@unibo.it)

E-mail address: [milisic@math.univ-paris13.fr](mailto:milisic@math.univ-paris13.fr)

E-mail address: [vauchelet@math.univ-paris13.fr](mailto:vauchelet@math.univ-paris13.fr)