



Research article

A characteristics approach to shock formation in 2D Euler with azimuthal symmetry and entropy

Isaac Neal¹, Steve Shkoller^{1,*} and Vlad Vicol²

¹ Department of Mathematics, University of California Davis, One Shields Avenue, Davis, CA 95616, USA

² Courant Institute of Mathematical Sciences, New York University, 251 Mercer St, New York, NY 10012, USA

* **Correspondence:** Email: shkoller@math.ucdavis.edu; Tel: +1-530-754-0389; Fax: +1-530-752-6635.

Abstract: We provide a detailed analysis of the shock formation process for the non-isentropic 2d Euler equations in azimuthal symmetry. We prove that from an open set of smooth and generic initial data, solutions of the Euler equations form a first singularity or gradient blow-up. This first singularity is termed a Hölder $C^{\frac{1}{3}}$ *pre-shock*, and our analysis provides the first detailed description of this cusp solution. The novelty of this work relative to [1] is that we herein consider a much larger class of initial data, allow for a non-constant initial entropy, allow for a non-trivial sub-dominant Riemann variable, and introduce a host of new identities to avoid apparent derivative loss due to entropy gradients. The method of proof is also new and robust, exploring the transversality of the three different characteristic families to transform space derivatives into time derivatives. Our main result provides a fractional series expansion of the Euler solution about the pre-shock, whose coefficients are computed from the initial data.

Keywords: Euler equations; shock formation; singularity formation; characteristics

Mathematics Subject Classification: 35Q31, 35L67

1. Introduction

Investigating shock formation and development is one of the central problems of hyperbolic PDE. Establishing *shock formation* (gradient blowup) from smooth initial data, in a *constructive* manner, is crucial for analyzing the dynamics of the resulting discontinuous shock waves. A precise description of the solution at the *pre-shock* (the spacetime set where smooth solutions first form cusps) is what allows for a full characterization of singularity propagation, especially in multiple space dimensions (see § 1.2 for details).

This paper establishes shock formation for smooth solutions of the non-isentropic two-dimensional compressible Euler equations in azimuthal symmetry. When compared to [2] this work gives a detailed

description of the solution near the pre-shock as a fractional power series. This paper also goes beyond [1] by establishing shock formation in the *non-isentropic* setting, and with *minimal constraints* imposed on the initial data (see § 1.2 for details).

Beyond the result itself, we develop a new robust proof strategy for establishing shock formation for a complex system of hyperbolic PDEs with multiple wave speeds. Instead of appealing to modulated self-similar analysis (cf. [1, 2]), we use new variables that satisfy pointwise and integral identities and accurately capture the compressible Euler dynamics (see § 1.3 for details).

1.1. The compressible Euler equations

The Euler equations of gas dynamics consist of the three conservation laws for momentum, mass, and energy, given respectively by

$$\partial_t(\rho u) + \operatorname{div}(\rho u \otimes u + pI) = 0, \quad (1.1a)$$

$$\partial_t \rho + \operatorname{div}(\rho u) = 0, \quad (1.1b)$$

$$\partial_t E + \operatorname{div}((p + E)u) = 0. \quad (1.1c)$$

In two space dimensions, the focus of this paper, $u : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}^2$ denotes the velocity vector field, $\rho : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}_+$ denotes the strictly positive density function, $E : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}$ denotes the total energy function, and $p : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}$ denotes the pressure function which is related to (u, ρ, E) by the identity $p = (\gamma - 1)(E - \frac{1}{2}\rho|u|^2)$, where $\gamma > 1$ denotes the adiabatic exponent. For the analysis of the shock formation process, it is convenient to replace conservation of energy (1.1c) with transport of entropy

$$\partial_t S + u \cdot \nabla S = 0. \quad (1.1d)$$

Here, $S : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}$ denotes the specific entropy, and the equation-of-state for pressure is written as

$$p(\rho, S) = \frac{1}{\gamma} \rho^\gamma e^S. \quad (1.2)$$

In preparation for reducing the equations to a more symmetric form, using Riemann-type variables, we introduce the adiabatic exponent

$$\alpha = \frac{\gamma-1}{2}$$

so that the (rescaled) sound speed reads

$$\sigma = \frac{1}{\alpha} \sqrt{\frac{\partial p}{\partial \rho}} = \frac{1}{\alpha} e^{\frac{S}{2}} \rho^\alpha. \quad (1.3)$$

With this notation, the ideal gas equation of state (1.2) becomes

$$p = \frac{\alpha^2}{\gamma} \rho \sigma^2. \quad (1.4)$$

The Euler equations (1.1a), (1.1b), and (1.1d), as a system for (u, σ, S) , are then given by

$$\partial_t u + (u \cdot \nabla)u + \alpha \sigma \nabla \sigma = \frac{\alpha}{2\gamma} \sigma^2 \nabla S, \quad (1.5a)$$

$$\partial_t \sigma + (u \cdot \nabla)\sigma + \alpha \sigma \operatorname{div} u = 0, \quad (1.5b)$$

$$\partial_t S + (u \cdot \nabla)S = 0. \quad (1.5c)$$

We let $\omega = \nabla^\perp \cdot u$ denote the scalar vorticity, and define the *specific vorticity* by $\zeta = \frac{\omega}{\rho}$. A straightforward computation shows that ζ is a solution to

$$\partial_t \zeta + (u \cdot \nabla) \zeta = \frac{\alpha \sigma}{\gamma \rho} \nabla^\perp \sigma \cdot \nabla S. \quad (1.6)$$

The term $\frac{\alpha \sigma}{\gamma \rho} \nabla^\perp \sigma \cdot \nabla S$ on the right side of (1.6) can also be written as $\rho^{-3} \nabla^\perp \rho \cdot \nabla p$ and is referred to as *baroclinic torque*.

The goal of this paper is to give a constructive proof of shock formation for (1.5), from smooth initial data, via a method powerful enough to capture a high-order series expansion of all fields at the preshock, information which is in turn necessary to study the shock development problem. More precisely, we prove:

Theorem 1.1 (Main result, abbreviated). *From smooth, non-isentropic initial data with azimuthal symmetry lying in an open set*, there exist smooth solutions to the 2d Euler equations (1.1) that form a gradient blowup singularity at a computable time T_* [†] and spatial location. More specifically, there exists $\xi_* \in \mathbb{T}$ such that when the 2d Euler equations are expressed in polar coordinates as in (2.1), the azimuthal component of the flow u_θ and the sound speed σ form $C^{0, \frac{1}{3}}$ cusps along the ray $\theta = \xi_*$ at the time of the blowup, and are given by the fractional series expansions*

$$\begin{aligned} u_\theta(r, \theta, T_*) &= r(\mathbf{b}_0 + \mathbf{b}_1(\theta - \xi_*)^{1/3} + \mathbf{b}_2(\theta - \xi_*)^{2/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|)), \\ \sigma(r, \theta, T_*) &= r(\mathbf{c}_0 + \mathbf{b}_1(\theta - \xi_*)^{1/3} + \mathbf{b}_2(\theta - \xi_*)^{2/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|)), \end{aligned}$$

for θ in a neighborhood of radius $\sim \varepsilon^3$,[‡] while the radial component u_r of the flow, the specific entropy S , and the specific vorticity ζ remain $C^{1, \frac{1}{3}}$, with fractional series expansions

$$\begin{aligned} u_r(r, \theta, T_*) &= r(\hat{\mathbf{a}}_0 + \hat{\mathbf{a}}_3(\theta - \xi_*) + \hat{\mathbf{a}}_4(\theta - \xi_*)^{4/3} + \mathcal{O}(\varepsilon^{-1/2}|\theta - \xi_*|^{5/3})), \\ S(r, \theta, T_*) &= \mathbf{k}_0 + \mathbf{k}_3(\theta - \xi_*) + \mathbf{k}_4(\theta - \xi_*)^{4/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{5/3}), \\ \zeta(r, \theta, T_*) &= \mathbf{v}_0 + \mathbf{v}_3(\theta - \xi_*) + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{4/3}). \end{aligned}$$

Here, the constants $\hat{\mathbf{a}}_0, \hat{\mathbf{a}}_3, \hat{\mathbf{a}}_4, \mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2, \mathbf{c}_0, \mathbf{k}_0, \mathbf{k}_3, \mathbf{k}_4$, and \mathbf{v}_0 are $\mathcal{O}(1)$ while \mathbf{v}_3 is $\mathcal{O}(\varepsilon^{-1})$.[§]

1.2. Motivation and prior results

We recall that the classical proofs of finite-time singularity formation for the compressible Euler equations and related hyperbolic systems are not constructive (see e.g. [3–5] for small smooth perturbations near constant states, and [6, 7] for large data). We refer the reader to [8–10] for an extensive bibliographic account of this classical theory.

A constructive proof of blowup, and equally importantly, a detailed description of the solution at the pre-shock, is necessary in order to establish shock development. By definition, shock development refers to the instantaneous development of the discontinuous shock wave from the $C^{0, \frac{1}{3}}$ Hölder cusp at the pre-shock. This is especially true in multiple space dimensions: while the theory of weak solutions

*See § 2.3-2.4 for the details of the pertinent set of initial data.

†We abuse notation here, because the time T_* used here differs from the time T_* referenced in the rest of the paper by a constant dependent on $\gamma > 1$. See § 2.1.

‡Here ε^{-1} is a large parameter quantifying the absolute size of slope of the initial data. See § 2.3 for details.

§See § 2.2 for the details of our use of $\mathcal{O}(\cdot)$ and \sim .

for 1D hyperbolic systems is well-developed (see e.g. [8]), many of the techniques used in the 1D theory either do not apply in multiple space dimensions[¶] or are not precise enough to be useful for the shock development problem, which requires bounds on derivatives of the solution.

Departing from the weak solutions perspective, Lebaud [12] established shock formation and development for the one-dimensional p -system (a variant on 1D isentropic Euler). These results were expanded upon by Chen and Dong [13] and Kong [14]. Studying shock development in the p -system does not per se prove anything about physical solutions of Euler, because physical solutions of Euler that have shocks cannot be isentropic (see § 2.2 of [1] or § 3 of [15] for details). Moreover, non-isentropic solutions of Euler are generically not irrotational due to a misalignment of pressure and entropy gradients (see (1.6) above and § 4 of [15] for the 3D case), so physical solutions which have shocks are also generically not irrotational. Studying shock development for piecewise isentropic or even piecewise irrotational solutions of Euler is called the *restricted shock development* problem. For the restricted shock development problem, Christodoulou established shock formation and development for irrotational flows in his landmark books [16, 17]. Yin [18] wrote the first paper establishing shock formation and development for non-isentropic Euler, but confined to spherical symmetry (see also [19]). Luk and Speck [20] proved shock formation for the 2D isentropic Euler equations in the presence of vorticity by an extension of Christodoulou's geometric framework to allow for vorticity transport. In [21], they later generalized their 2D result to the full 3D non-isentropic setting.

A different perspective was taken by Buckmaster, Shkoller, and Vicol [2,22,23], who used modulated self-similar variables to construct the first gradient singularity (a *point shock*) from generic smooth initial data. In [2] they constructed shocks for 2D isentropic Euler in azimuthal symmetry and characterized the shock profile as an asymptotically self-similar, stable 1D blowup profile. After that, they proved for the first time that the 3D isentropic Euler equations generically form a stable *point shock*, even in the presence of vorticity [22]. The important generalization to the full non-isentropic setting was achieved in [23], where it is also shown that irrotational data instantaneously creates vorticity due to baroclinic torque, and the vorticity remains uniformly bounded up to and including the time of the first gradient singularity.

The analysis of the Euler evolution beyond the time of the first gradient blowup was recently addressed by Shkoller and Vicol [24] by studying the so-called Maximal Globally Hyperbolic Development (MGHD) of smooth and compressive Cauchy data. This can be understood to be the largest (local) spacetime that contains a smooth (and invertible) evolution of the Cauchy data. The future temporal boundary of this spacetime consists of the codimension-2 manifold of pre-shocks (containing the spacetime set of first gradient catastrophes), the singular set (a downstream hypersurface of gradient blowups emanating from the pre-shock manifold), and the Cauchy horizon (an upstream hypersurface emanating from the pre-shock set which the smooth Euler solution can never cross). A partial construction of the MGHD was also obtained by Abbrescia and Speck [25] who were able to evolve the Euler solution up the union of the pre-shock set and the singular set (but the upstream evolution up to the Cauchy horizon was not treated).

Buckmaster, Drivas, Shkoller, and Vicol [1] established for the first time shock development in the presence of vorticity, by working in azimuthal symmetry. By improving upon [2], the solution at the pre-shock is described in [1] by a fractional series, assuming that the flow is initially isentropic ($k_0 \equiv 0$ in (2.5c) below) and that the subdominant Riemann variable vanishes ($z_0 \equiv 0$ in (2.5b) below).

[¶]For example, the BV estimates utilized in the classical theory of shocks for 1D hyperbolic systems fails for $d \geq 2$. See [11].

They then used this detailed description of the solution to establish shock development for 2D Euler within the class of azimuthal solutions. The paper [1] is the first to also confirm the production of both a discontinuous shock wave and two surfaces of cusp singularities emanating from the pre-shock, as predicted by Landau and Lifschitz [26].

1.3. New ideas

This paper breaks with [2] and [1] by forgoing the use of self-similar variables. Instead, we use only the fine structure of the Euler system written in the characteristic coordinates that correspond to the three different wave speeds present in the system. We show that the sound speed remains bounded from below up to the time of the first blowup (see Proposition 4.1), which means that the three wave speeds remain uniformly transverse to one another up to the blowup time. This transversality allows us to prove useful integral bounds (see Lemma 3.1 and § 4) and allows us to exchange space derivatives for time derivatives (see § 5), which can be integrated to obtain identities for the higher-order derivatives of our variables. This exchange of space for time derivatives via transversality is the key new idea of this work.

The implementation of this idea is made possible by using the special *differentiated Riemann variables* introduced in [1]. These new variables, labeled q^w and q^z , evolve along the characteristics of the fastest and slowest wave speeds, respectively, and they do not experience derivative loss (see § 3 of [1] or § 3.2 below). Whereas [1] utilized q^w and q^z for studying shock development, we use q^w and q^z to also establish shock formation in the non-isentropic setting. Using pointwise and integral identities for q^w and q^z , we are able to obtain estimates for our variables and their derivatives up to the blowup time without first establishing the uniqueness or location of the blowup label; we instead derive the uniqueness and location of the blowup label as a result of our estimates (see § 10.3).

We note that because we avoid self-similar analysis, we are able to place far fewer assumptions on our initial data than in [1]. When compared to [1], we also obtain a higher-order fractional series expansion of the solution at the time of blowup (see Theorem 2.1).

2. Azimuthal symmetry

2.1. The Euler equations in polar coordinates and azimuthal symmetry

The 2D Euler equations (1.5) take the following form in polar coordinates for the variables (u_θ, u_r, ρ, S) :

$$\left(\partial_t + u_r \partial_r + \frac{1}{r} u_\theta \partial_\theta\right) u_r - \frac{1}{r} u_\theta^2 + \alpha \sigma \partial_r \sigma = \frac{\alpha}{2\gamma} \sigma^2 \partial_r S, \quad (2.1a)$$

$$\left(\partial_t + u_r \partial_r + \frac{1}{r} u_\theta \partial_\theta\right) u_\theta + \frac{1}{r} u_r u_\theta + \alpha \frac{\sigma}{r} \partial_\theta \sigma = \frac{\alpha}{2\gamma} \frac{\sigma^2}{r} \partial_\theta S, \quad (2.1b)$$

$$\left(\partial_t + u_r \partial_r + \frac{1}{r} u_\theta \partial_\theta\right) \sigma + \alpha \sigma \left(\frac{1}{r} u_r + \partial_r u_r + \frac{1}{r} \partial_\theta u_\theta\right) = 0, \quad (2.1c)$$

$$\left(\partial_t + u_r \partial_r + \frac{1}{r} u_\theta \partial_\theta\right) S = 0. \quad (2.1d)$$

We introduce the new variables[¶]

$$u_\theta(r, \theta, t) = rb(\theta, t), \quad u_r(r, \theta, t) = ra(\theta, t), \quad \sigma(r, \theta, t) = rc(\theta, t), \quad S(r, \theta, t) = k(\theta, t). \quad (2.2)$$

[¶]Note that our symmetry constraints make S discontinuous at the origin unless S is constant. For this reason, a classical solution of the 2D Euler equations (1.5) is recovered from the azimuthal variables (a, b, c, k) via (2.2) on the punctured plane. Alternatively, we may restrict the domain of evolution for 2D Euler to an annular domain pushed forward under the flow of u (see [2, § 2.1]).

The system (2.1) then takes the form

$$(\partial_t + b\partial_\theta) a + a^2 - b^2 + \alpha c^2 = 0 \quad (2.3a)$$

$$(\partial_t + b\partial_\theta) b + \alpha c \partial_\theta c + 2ab = \frac{\alpha}{2\gamma} c^2 \partial_\theta k \quad (2.3b)$$

$$(\partial_t + b\partial_\theta) c + \alpha c \partial_\theta b + \gamma ac = 0 \quad (2.3c)$$

$$(\partial_t + b\partial_\theta) k = 0. \quad (2.3d)$$

For simplicity of the presentation, we will set $\gamma = 2$ from here on; note however that all statements in this paper apply *mutatis mutandis* to the case of a general $\gamma > 1$. The Riemann functions w and z are defined by

$$w = b + c, \quad z = b - c, \quad (2.4a)$$

$$b = \frac{1}{2}(w + z), \quad c = \frac{1}{2}(w - z). \quad (2.4b)$$

It is convenient to rescale time, letting $\partial_t \mapsto \frac{3}{4}\partial_{\tilde{t}}$, and for notational simplicity, we continue to write t for \tilde{t} . With this temporal rescaling employed, the system (2.3c) can be equivalently written as

$$\partial_t w + \lambda_3 \partial_\theta w = -\frac{8}{3}aw + \frac{1}{24}(w - z)^2 \partial_\theta k, \quad (2.5a)$$

$$\partial_t z + \lambda_1 \partial_\theta z = -\frac{8}{3}az + \frac{1}{24}(w - z)^2 \partial_\theta k, \quad (2.5b)$$

$$\partial_t k + \lambda_2 \partial_\theta k = 0, \quad (2.5c)$$

$$\partial_t a + \lambda_2 \partial_\theta a = -\frac{4}{3}a^2 + \frac{1}{3}(w + z)^2 - \frac{1}{6}(w - z)^2. \quad (2.5d)$$

where the three wave speeds are given by

$$\lambda_1 = \frac{1}{3}w + z < \lambda_2 = \frac{2}{3}w + \frac{2}{3}z < \lambda_3 = w + \frac{1}{3}z. \quad (2.6)$$

We note that (2.3c) takes the form

$$\partial_t c + \lambda_2 \partial_\theta c + \frac{1}{2}c \partial_\theta \lambda_2 = -\frac{8}{3}ac. \quad (2.7)$$

Finally, we denote the specific vorticity (1.6) in azimuthal symmetry by

$$\varpi = 4(w + z - \partial_\theta a) c^{-2} e^k, \quad (2.8)$$

which satisfies the evolution equation

$$\partial_t \varpi + \lambda_2 \partial_\theta \varpi = \frac{8}{3}a\varpi + \frac{4}{3}e^k \partial_\theta k. \quad (2.9)$$

2.2. Notation

In most of what follows, there will be an important parameter $\varepsilon > 0$, and $a \lesssim b$ will be used to signify that $a \leq Cb$ for some constant C independent of ε and any variables x, θ , or t . However, the constant can depend on the implicit constants in the assumptions on the initial data in § 2.3 and can depend on our choice of $\gamma > 1$ for the pressure law**. We will use the notation $a \sim b$ to express $a \lesssim b \lesssim a$. We will also write

$$f = \mathcal{O}(g)$$

**We have already chosen to fix $\gamma = 2$ for the entirety of this paper, but our result will hold for arbitrary $\gamma > 1$, and the value of γ will effect the constants.

to express that $|f| \lesssim g$ everywhere in the relevant domain. We will express bounds of the type

$$f(x, t) = \begin{cases} \mathcal{O}(b_1) & |x| \leq \varepsilon^2 \\ \mathcal{O}(b_2) & |x| \geq \varepsilon^2 \end{cases} \quad \text{simply as} \quad f = \mathcal{B}(b_1; b_2).$$

Often below we will have functions f defined on $\mathbb{T} \times [0, T_*)$ and maps $\Psi : \mathbb{T} \times [0, T_*) \rightarrow \mathbb{T}$, and we will use the notation

$$f \circ \Psi(x, t) := f(\Psi(x, t), t).$$

When such an inverse exists, we will write Ψ^{-1} to denote the function such that $\Psi^{-1} \circ \Psi(x, t) = \Psi \circ \Psi^{-1}(x, t) = x$ for all t .

While the spatial variable θ for (2.5) lies in \mathbb{T} , we will often identify \mathbb{T} with the interval $(-\pi, \pi]$.

2.3. Assumptions on the Initial Data

Our initial data will be $w_0, z_0, k_0, a_0 \in H^6(\mathbb{T})$, where z_0, k_0 , and a_0 all satisfy

$$\|\partial_x^j k_0\|_{L^\infty} \lesssim \varepsilon^{\gamma_j}, \quad \|\partial_x^j a_0\|_{L^\infty} \lesssim \varepsilon^{\alpha_j}, \quad \|\partial_x^j z_0\|_{L^\infty} \lesssim \varepsilon^{\beta_j}, \quad (2.10)$$

for $j = 0, 1, 2, 3, 4, 5$, where $\alpha_j, \beta_j, \gamma_j$ are fixed constants satisfying the relations

- $\alpha_0, \beta_0, \gamma_0 \geq 0$,
- $\gamma_1 \geq \mu, \alpha_1 \geq 0$,
- $\gamma_j \geq \mu - j$ for $j = 2, 3, 4, 5$,
- $\alpha_j \geq \mu + 1 - j$ for $j = 2, 3, 4, 5$,
- $\beta_j \geq \mu - j$ for $j = 1, 2, 3, 4, 5$.

Here $\mu > 0$ is a fixed positive constant that is a lower bound on the ℓ^∞ distance of our vector of parameters $(\alpha_2, \dots, \alpha_5, \gamma_1, \dots, \gamma_5, \beta_1, \dots, \beta_5)$ from the boundary of the open set defined by the constraints $\beta_1 > -1$, $\gamma_1 > 0$, etc. Additionally, we assume that w_0 satisfies

1. $w_0 \sim 1$,
2. $w'_0(0) := -\frac{1}{\varepsilon}$ and $|w'_0(x)| < \varepsilon^{-1}$ for all $x \neq 0$,
3. $w'_0(x) \geq -\frac{1}{\varepsilon} + C\varepsilon^{\frac{\mu}{2}-1}$ for all $|x| \geq \varepsilon^{3/2}$, and some constant $C > 0$.
4. $w''_0(x) \sim \varepsilon^{-4}$ for all $|x| \leq \varepsilon^{3/2}$,
5. $|\partial_x^4 w_0(x)| \lesssim \varepsilon^{\mu-5}$ for all $|x| \leq \varepsilon^2$,
6. $\|\partial_x^5 w_0\|_{L^\infty} \lesssim \varepsilon^{-7}$,

and that z_0 satisfies

$$\max z_0 < \min w_0. \quad (2.11)$$

Note that an immediate consequence of our assumptions is that w_0 must also satisfy

- $w''_0(0) = 0$,

- $|w_0''(x)| \lesssim \varepsilon^{-2}$ for $|x| \leq \varepsilon^2$,
- $\|w_0''\|_{L^\infty} \lesssim \varepsilon^{-\frac{5}{2}}$,
- $\|w_0'''\|_{L^\infty} \lesssim \varepsilon^{-4}$,
- $\|\partial_x^4 w_0\|_{L^\infty} \lesssim \varepsilon^{-\frac{11}{2}}$.

The following additional constraints are not at all necessary for proving our theorem, but they do make the formulas of the proof below cleaner

$$\alpha_0 = \beta_0 = \gamma_0 = \alpha_1 = 0, \quad \beta_1 \leq 0, \quad \text{and} \quad \alpha_j, \beta_j, \gamma_j \leq 1 \quad \forall j = 0, 1, 2, 3, 4, 5. \tag{2.12}$$

Note that the constraints made here on the first five derivatives of (w_0, z_0, k_0, a_0) are much less stringent than those imposed in [1]. In [1], the authors assume that k_0 is constant, z_0 is identically 0, and that w_0' and a_0 have support with diameter $\mathcal{O}(\varepsilon^{1/2})$, among other constraints. Here we do away with such unnecessary hypotheses. Additionally, the result of this paper applies to a wide range of parameters $(\alpha_j, \beta_j, \gamma_j)$, whereas in [1] the authors only work with $(\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4) = (1, 0, 0, 0)$, which is only one point in our admissible range for these parameters.

In what follows, we will parametrize time so that the initial time is always $t = -\varepsilon$. The local well-posedness theory of (1.5) implies that for any $(w_0, z_0, k_0, a_0) \in H^6(\mathbb{T})$ there exists a time $T_* \in (-\varepsilon, +\infty]$ such that there exists a unique C^1 solution (w, z, k, a) of (2.5) satisfying $(w, z, k, a)|_{t=-\varepsilon} = (w_0, z_0, k_0, a_0)$. Furthermore, (w, z, k, a) is guaranteed to be in $C^0([-\varepsilon, T_*]; H^6(\mathbb{T})) \cap C^1([-\varepsilon, T_*]; H^5(\mathbb{T}))$. Additionally, it follows from the standard theory of (1.5) that if $T_* < \infty$ then

$$\int_{-\varepsilon}^{T_*} \|\partial_\theta w(t)\|_{L^\infty} + \|\partial_\theta z(t)\|_{L^\infty} + \|\partial_\theta k(t)\|_{L^\infty} + \|\partial_\theta a(t)\|_{L^\infty} dt = \infty. \tag{2.13}$$

The inequalities above can be made into open constraints by making them strict inequalities. While the two pointwise constraints that require w_0' to attain its unique global minimum at $x = 0$ and $w_0'(0) = -\frac{1}{\varepsilon}$ are not open constraints, for any suitably small perturbation of initial data (w_0, z_0, k_0, a_0) satisfying all of the above constraints, one can recover the two pointwise constraints by translating in space and rescaling the solution in time. Since the spatial translation and time rescaling can be made sufficiently small, there exists an open set of initial data around the functions (w_0, z_0, k_0, a_0) described above for which the results of Theorem 2.1 below still hold. Thus, the shock formation we describe is stable.

2.4. Statement of the main theorem

Theorem 2.1 (Main theorem). *For $\mu > 0$, $\varepsilon > 0$ sufficiently small, and initial data $(w, z, k, a)|_{t=-\varepsilon} = (w_0, z_0, k_0, a_0)$ in the open set described in § 2.3, there exists a blowup time T_* with $|T_*| \lesssim \varepsilon^{1+\mu}$, a unique blowup location $\xi_* \in \mathbb{T}$, and unique C^1 solutions (w, z, k, a) to (2.5) on $\mathbb{T} \times [-\varepsilon, T_*)$ such that $|x_*| \lesssim \varepsilon^{2+\mu}$,*

$$w(\cdot, T_*) \in C^{0, \frac{1}{3}}(\mathbb{T}), \quad z(\cdot, T_*), k(\cdot, T_*), a(\cdot, T_*), \varpi(\cdot, T_*) \in C^{1, \frac{1}{3}}(\mathbb{T}),$$

where ϖ is the specific vorticity (see (2.8)). Furthermore, there exists a unique blowup label $x_* \in (-\pi, \pi]$ such that

$$\lim_{t \rightarrow T_*} \eta(x_*, t) = \xi_*$$

where η is the 3-characteristic defined in § 3.1. In a neighborhood $\theta \in \eta([- \varepsilon^2, \varepsilon^2], T_*)$ of radius $\sim \varepsilon^3$ the functions $w(\cdot, T_*)$, $z(\cdot, T_*)$, $k(\cdot, T_*)$, and $a(\cdot, T_*)$ have the following fractional series expansions:

There exist constants $\mathring{a}_0^w, \mathring{a}_1^w, \mathring{a}_2^w$ with

$$|\mathring{a}_0^w| \lesssim 1, \quad |\mathring{a}_1^w| \lesssim 1, \quad |\mathring{a}_2^w| \lesssim 1,$$

such that

$$\begin{aligned} w(\theta, T_*) &= \mathring{a}_0^w + \mathring{a}_1^w(\theta - \xi_*)^{1/3} + \mathring{a}_2^w(\theta - \xi_*)^{2/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|), \\ \partial_\theta w(\theta, T_*) &= \frac{1}{3}\mathring{a}_1^w(\theta - \xi_*)^{-2/3} + \frac{2}{3}\mathring{a}_2^w(\theta - \xi_*)^{-1/3} + \mathcal{O}(\varepsilon^{-1}), \\ \partial_\theta^2 w(\theta, T_*) &= -\frac{2}{9}\mathring{a}_1^w(\theta - \xi_*)^{-5/3} - \frac{2}{9}\mathring{a}_2^w(\theta - \xi_*)^{-4/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{-1}), \\ \partial_\theta^3 w(\theta, T_*) &= \frac{10}{27}\mathring{a}_1^w(\theta - \xi_*)^{-8/3} + \frac{8}{27}\mathring{a}_2^w(\theta - \xi_*)^{-7/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{-2}). \end{aligned} \quad (2.14)$$

There exist constants $\mathring{a}_0^z, \mathring{a}_3^z, \mathring{a}_4^z$ with

$$|\mathring{a}_0^z| \lesssim 1, \quad |\mathring{a}_3^z| \lesssim \varepsilon^{\mu-1}, \quad |\mathring{a}_4^z| \lesssim \varepsilon^{\mu-1},$$

such that

$$\begin{aligned} z(\theta, T_*) &= \mathring{a}_0^z + \mathring{a}_3^z(\theta - \xi_*) + \mathring{a}_4^z(\theta - \xi_*)^{4/3} + \mathcal{O}(\varepsilon^{\mu-2}|\theta - \xi_*|^{5/3}), \\ \partial_\theta z(\theta, T_*) &= \mathring{a}_3^z + \frac{4}{3}\mathring{a}_4^z(\theta - \xi_*)^{1/3} + \mathcal{O}(\varepsilon^{\mu-2}|\theta - \xi_*|^{2/3}), \\ \partial_\theta^2 z(\theta, T_*) &= \frac{4}{9}\mathring{a}_4^z(\theta - \xi_*)^{-2/3} + \mathcal{O}(\varepsilon^{\mu-2}|\theta - \xi_*|^{-1/3}), \\ \partial_\theta^3 z(\theta, T_*) &= -\frac{8}{27}\mathring{a}_4^z(\theta - \xi_*)^{-5/3} + \mathcal{O}(\varepsilon^{\mu-2}|\theta - \xi_*|^{4/3}). \end{aligned} \quad (2.15)$$

There exist constants $\mathring{a}_0^k, \mathring{a}_3^k, \mathring{a}_4^k$ with

$$|\mathring{a}_0^k| \lesssim 1, \quad |\mathring{a}_3^k| \lesssim \varepsilon^\mu, \quad |\mathring{a}_4^k| \lesssim \varepsilon^{\gamma_2+1\wedge\mu},$$

such that

$$\begin{aligned} k(\theta, T_*) &= \mathring{a}_0^k + \mathring{a}_3^k(\theta - \xi_*) + \mathring{a}_4^k(\theta - \xi_*)^{4/3} + \mathcal{O}(\varepsilon^{\gamma_2\wedge\mu-1}|\theta - \xi_*|^{5/3}), \\ \partial_\theta k(\theta, T_*) &= \mathring{a}_3^k + \frac{4}{3}\mathring{a}_4^k(\theta - \xi_*)^{1/3} + \mathcal{O}(\varepsilon^{\gamma_2\wedge\mu-1}|\theta - \xi_*|^{2/3}), \\ \partial_\theta^2 k(\theta, T_*) &= \frac{4}{9}\mathring{a}_4^k(\theta - \xi_*)^{-2/3} + \mathcal{O}(\varepsilon^{\gamma_2\wedge\mu-1}|\theta - \xi_*|^{-1/3}), \\ \partial_\theta^3 k(\theta, T_*) &= -\frac{8}{27}\mathring{a}_4^k(\theta - \xi_*)^{-5/3} + \mathcal{O}(\varepsilon^{\gamma_2\wedge\mu-1}|\theta - \xi_*|^{4/3}). \end{aligned} \quad (2.16)$$

There exist constants $\mathring{a}_0^a, \mathring{a}_3^a, \mathring{a}_4^a$ with

$$|\mathring{a}_0^a| \lesssim 1, \quad |\mathring{a}_3^a| \lesssim 1, \quad |\mathring{a}_4^a| \lesssim 1,$$

such that

$$\begin{aligned} a(\theta, T_*) &= \mathring{a}_0^a + \mathring{a}_3^a(\theta - \xi_*) + \mathring{a}_4^a(\theta - \xi_*)^{4/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{5/3}), \\ \partial_\theta a(\theta, T_*) &= \mathring{a}_3^a + \frac{4}{3}\mathring{a}_4^a(\theta - \xi_*)^{1/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{2/3}), \\ \partial_\theta^2 a(\theta, T_*) &= \frac{4}{9}\mathring{a}_4^a(\theta - \xi_*)^{-2/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{-1/3}), \\ \partial_\theta^3 a(\theta, T_*) &= -\frac{8}{27}\mathring{a}_4^a(\theta - \xi_*)^{-5/3} + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{4/3}). \end{aligned} \quad (2.17)$$

There exist constants $\mathring{a}_0^\varpi, \mathring{a}_3^\varpi$ with

$$|\mathring{a}_0^\varpi| \lesssim 1, \quad |\mathring{a}_3^\varpi| \lesssim \varepsilon^{-1},$$

such that

$$\begin{aligned} \varpi(\theta, T_*) &= \mathring{a}_0^\varpi + \mathring{a}_3^\varpi(\theta - \xi_*) + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{4/3}), \\ \partial_\theta \varpi(\theta, T_*) &= \mathring{a}_3^\varpi + \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{1/3}), \\ \partial_\theta^2 \varpi(\theta, T_*) &= \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{-2/3}), \\ \partial_\theta^3 \varpi(\theta, T_*) &= \mathcal{O}(\varepsilon^{-1}|\theta - \xi_*|^{-5/3}). \end{aligned} \tag{2.18}$$

Moreover, the C^5 regularity away from the pre-shock is characterized by

$$\begin{aligned} \max_{n \leq 5} |\partial_\theta^n w(\eta(x, t), t)| + |\partial_\theta^n z(\eta(x, t), t)| + |\partial_\theta^n k(\eta(x, t), t)| + |\partial_\theta^n a(\eta(x, t), t)| \\ \lesssim \mathcal{B}(\varepsilon^{-7}[\frac{1}{2\varepsilon}(T_* - t) + c(\varepsilon + t)\varepsilon^{-4}(x - x_*)^2]^{-1}; \varepsilon^{-16}). \end{aligned} \tag{2.19}$$

Theorem 1.1 clearly follows from Theorem 2.1 as an immediate corollary.

2.5. Outline of the proof of Theorem 2.1

In this paper, we will show that the classical solution (w, z, k, a) of (2.5) with the initial data specified in § 2.3 breaks down in finite time, and that this occurs when the flow η of the fastest wave speed λ_3 ceases to be a diffeomorphism. More specifically, the blowup time T_* will be characterized as the first time t when $\min_x \eta_x(x, t) = 0$. We will also establish that there is a unique Lagrangian label x_* for which $\eta_x(x_*, T_*) = 0$, which will imply that η_{xx} vanishes at (x_*, T_*) as well. While $w, z, k, a, \partial_\theta z, \partial_\theta k$, and $\partial_\theta a$ will be shown to remain bounded on $\mathbb{T} \times [-\varepsilon, T_*]$, $\partial_\theta w$ will be shown to go to $-\infty$ at the point $(\xi_*, T_*) := (\eta(x_*, T_*), T_*)$ and remain smooth elsewhere. The key ingredient for implementing the above-described strategy is to show that the functions $w \circ \eta, z \circ \eta, k \circ \eta$, and $a \circ \eta$ remain as smooth as their initial data, *uniformly* up to T_* . The authors of [1] proved such uniform estimates using self-similar analysis, but only in a special case.^{††} In this paper, we prove uniform C^5 estimates for $(w, z, k, a) \circ \eta$ on $\mathbb{T} \times [-\varepsilon, T_*]$, even in the most general setting, not by relying on self-similar variables, but by instead using the transversality of various families of characteristics. This allows us to also consider a much broader class of initial data than previously considered in [1]. Once we have shown that all the variables stay smooth along the η characteristic, we obtain our functional description of the solution near (ξ_*, T_*) by inverting the map $x \mapsto \eta(x, t)$ for (x, t) near the point (x_*, T_*) . In light of the constraints $\eta_x(x_*, T_*) = \eta_{xx}(x_*, T_*) = 0$, this amounts to the inversion of what is to leading order a cubic polynomial, resulting in fractional series expansions of w, z, k , and a near (ξ_*, T_*) in terms of powers of $(\theta - \xi_*)^{1/3}$.

This paper is organized as follows:

1. In § 4 we bound $|T_*|$ and prove that $\partial_\theta w$ must become infinite at time T_* . We use a simple bootstrap argument to get estimates for w, z, k, a and their first derivatives up to time $\varepsilon \wedge T_*$. Using these estimates, we show that η_x must have a zero before time $t = \varepsilon$, and conclude that $|T_*| \lesssim \varepsilon^{1+\mu}$. This implies that $\varepsilon \wedge T_* = T_*$ and therefore all of our estimates and identities hold up to time T_* . The fact that $\partial_\theta w$ must blow up then follows immediately from the fact that $\partial_\theta z, \partial_\theta k$, and $\partial_\theta a$ remain bounded up to time T_* (see (2.13) above).

^{††}The authors of [1] work in the case where z and k are identically zero and many more constraints are placed on w_0 and a_0 . See § 2.3 above for a discussion.

2. Next we show that $w \circ \eta, z \circ \eta, k \circ \eta$, and $a \circ \eta$ remain smooth up to time T_* . To do this, we first establish crucial identities in § 5, which result from the fact that the wave speeds are uniformly transverse to one another. Then in § 6 - 9 we prove pointwise bounds on z, k, a and their derivatives in terms of w and its derivatives by analyzing how our new variables evolve along the multiple wave speeds. This allows us to conclude in § 10 that w, z, k , and a all remain smooth along η .
3. Last we establish that the singularity occurs at a unique point $(\xi_*, T_*) \in \mathbb{T} \times [-\varepsilon, T_*]$ and we invert η near this point to obtain fractional series expansions for w, z, k , and a . We do this by establishing in § 10 that there is a unique point $(x_*, T_*) \in \mathbb{T} \times [-\varepsilon, T_*]$ where η_x vanishes and that $\eta_{xx}(x_*, T_*) = 0$ as well. Since $\eta(x, T_*) = \xi_* + \eta_{xxx}(x_*, T_*)(x - x_*)^3 + \mathcal{O}(|x - x_*|^4)$ near (x_*, T_*) , it follows (see § 11) that $(x - x_*) \sim (\theta - \xi_*)^{1/3}$ for small $|x - x_*|$ at time T_* , and the Taylor series expansions of the smooth functions $w \circ \eta(\cdot, T_*), z \circ \eta(\cdot, T_*), k \circ \eta(\cdot, T_*)$, and $a \circ \eta(\cdot, T_*)$ near x_* become fractional series expansions of $w(\cdot, T_*), z(\cdot, T_*), k(\cdot, T_*)$, and $a(\cdot, T_*)$ near ξ_* .

3. Preliminaries

3.1. The characteristics

Let $\varphi > 0$, and let Ψ be the flow of $\lambda := (1 - \varphi)w + (\frac{1}{3} + \varphi)z$.

$$\begin{aligned} \Psi_x &= e^{\int_{-\varepsilon}^t \partial_\theta \lambda \circ \Psi}. \\ \partial_t c + \lambda \partial_\theta c &= -(\varphi \partial_\theta w + (\frac{2}{3} - \varphi) \partial_\theta z + \frac{8}{3} a) c. \end{aligned} \tag{3.1}$$

If $c > 0$ everywhere, this tells us that

$$\begin{aligned} -\frac{1}{\varphi} \partial_t (\log c \circ \Psi) &= (\partial_\theta w + (\frac{2}{3} \frac{1}{\varphi} - 1) \partial_\theta z + \frac{8}{3} \frac{1}{\varphi} a) \circ \Psi. \\ \implies \partial_\theta \lambda \circ \Psi &= -\frac{1-\varphi}{\varphi} \partial_t (\log c \circ \Psi) + ((2 - \frac{2}{3} \frac{1}{\varphi})) \partial_\theta z - \frac{8}{3} \frac{1-\varphi}{\varphi} a \circ \Psi. \\ \implies \Psi_x &= (\frac{c_0}{c \circ \Psi})^{\frac{1-\varphi}{\varphi}} e^{\int_{-\varepsilon}^t (2 - \frac{2}{3} \frac{1}{\varphi}) \partial_\theta z - \frac{8}{3} \frac{1-\varphi}{\varphi} a \circ \Psi}. \end{aligned} \tag{3.2}$$

If $c \sim 1$ and $\partial_\theta z, a$ are bounded, then this lets us conclude that $\Psi_x \sim 1$. We will prove in the next section that $c \sim 1$ and that $\partial_\theta z, a$ are indeed bounded on $\mathbb{T} \times [-\varepsilon, T_*]$, so everything that follows is relevant.

In the case where $\varphi = \frac{2}{3}$, we have $\lambda = \lambda_1$, the first wave speed. Let ψ denote the corresponding flow, the so-called **1-characteristic**. Its first derivative satisfies

$$\psi_x = \left(\frac{c_0}{c \circ \psi} \right)^{\frac{1}{2}} e^{\int_{-\varepsilon}^t (\partial_\theta z - \frac{4}{3} a) \circ \psi}, \tag{3.3}$$

while its second derivative obeys

$$\begin{aligned} \psi_{xx} &= \psi_x \left(\frac{1}{2} \frac{c'_0}{c_0} + \int_{-\varepsilon}^t \psi_x (\partial_\theta^2 z - \frac{4}{3} \partial_\theta a) \circ \psi \right) - \frac{1}{2} \psi_x^2 \frac{\partial_\theta c \circ \psi}{c \circ \psi} \\ &=: \psi_x \Psi - \frac{1}{2} \psi_x^2 \frac{\partial_\theta c \circ \psi}{c \circ \psi}. \end{aligned} \tag{3.4}$$

$$=: \psi_x^2 (Q_1 - \frac{1}{2} c^{-1} \partial_\theta c) \circ \psi. \tag{3.5}$$

When $\varphi = \frac{1}{3}$, we have $\lambda = \lambda_2$ and the corresponding flow is the **2-characteristic**, ϕ . The first derivative of ϕ satisfies

$$\phi_x = \left(\frac{c_0}{c \circ \phi} \right)^2 e^{-\frac{16}{3} \int_{-\varepsilon}^t a \circ \phi} \quad (3.6)$$

while its second derivative obeys

$$\begin{aligned} \phi_{xx} &= \phi_x \left(2 \frac{c'_0}{c_0} - \frac{16}{3} \int_{-\varepsilon}^t \partial_x(a \circ \phi) \right) - 2\phi_x^2 \frac{\partial_\theta c \circ \phi}{c \circ \phi} \\ &=: \phi_x \Phi - 2\phi_x^2 \frac{\partial_\theta c \circ \phi}{c \circ \phi} \end{aligned} \quad (3.7)$$

$$=: \phi_x^2 (Q_2 - 2c^{-1} \partial_\theta c) \circ \phi. \quad (3.8)$$

When $\varphi = 0$, we have $\lambda = \lambda_3$ and the corresponding flow is the **3-characteristics**, η . Note that our analysis for $\varphi > 0$ breaks down for η , but also that w is essentially transported along η .

3.2. q^w and q^z

Our system (2.5) can be written as

$$\partial_t \vec{x} + A \partial_\theta \vec{x} = \vec{b} \quad (3.9)$$

where

$$\vec{x} := \begin{bmatrix} w \\ z \\ k \\ a \end{bmatrix}, \quad A := \begin{pmatrix} \lambda_3 & 0 & -\frac{1}{6}c^2 & 0 \\ 0 & \lambda_1 & -\frac{1}{6}c^2 & 0 \\ 0 & 0 & \lambda_2 & 0 \\ 0 & 0 & 0 & \lambda_2 \end{pmatrix}, \quad \vec{b} := \begin{bmatrix} -\frac{8}{3}aw \\ -\frac{8}{3}az \\ 0 \\ -\frac{4}{3}a^2 + \frac{1}{3}(w+z)^2 - \frac{1}{6}(w-z)^2 \end{bmatrix}.$$

Taking ∂_θ of (3.9) and diagonalizing A gives us

$$\partial_t \vec{y} + D \vec{y} = Q(\vec{x}, \vec{y})$$

where $D = \text{diag}(\lambda_3, \lambda_1, \lambda_2, \lambda_2)$, $\vec{y} := (\partial_\theta w - \frac{1}{4}c \partial_\theta k, \partial_\theta z + \frac{1}{4}c \partial_\theta k, \partial_\theta k, \partial_\theta a)$, and $Q : \mathbb{R}^8 \rightarrow \mathbb{R}^4$ is a third-order polynomial. This motivates the introduction of the following variables:

$$q^w := \partial_\theta w - \frac{1}{4}c \partial_\theta k \quad \text{and} \quad q^z := \partial_\theta z + \frac{1}{4}c \partial_\theta k. \quad (3.10)$$

One can check using the identities in § A.1 that

$$\partial_t(q^w \circ \eta \eta_x) = \left(-\frac{8}{3}a + \frac{1}{12}c \partial_\theta k\right) \circ \eta (q^w \circ \eta \eta_x) + \frac{1}{12}(c \partial_\theta k) \circ \eta (q^z \circ \eta \eta_x) - \frac{8}{3} \partial_x(a \circ \eta) w \circ \eta. \quad (3.11)$$

$$\partial_t(q^z \circ \psi \psi_x) = \left(-\frac{8}{3}a - \frac{1}{12}c \partial_\theta k\right) \circ \psi (q^z \circ \psi \psi_x) - \frac{1}{12}(c \partial_\theta k) \circ \psi (q^w \circ \psi \psi_x) - \frac{8}{3} \partial_x(a \circ \psi) z \circ \psi. \quad (3.12)$$

If we define

$$I_t(x) := e^{\frac{1}{8}k \circ \eta - \frac{8}{3} \int_{-\varepsilon}^t a \circ \eta} \quad (3.13)$$

then our equation for $\partial_t(q^w \circ \eta_{\lambda_x})$ gives us the Duhamel formula

$$\eta_x q^w \circ \eta = I_t \left[(w'_0 - \frac{1}{4} c_0 k'_0) e^{-\frac{1}{8} k_0} + \frac{1}{12} \int_{-\varepsilon}^t I_\tau^{-1} \eta_x (c \partial_\theta k q^z) \circ \eta \, d\tau - \frac{8}{3} \int_{-\varepsilon}^t I_\tau^{-1} w \circ \eta \partial_x (a \circ \eta) \, d\tau \right]. \quad (3.14)$$

It follows immediately from the definitions of λ_3 and q^w that

$$\eta_x = 1 + \int_{-\varepsilon}^t \eta_x \partial_\theta \lambda_3 \circ \eta \, d\tau = 1 + \int_{-\varepsilon}^t \eta_x q^w \circ \eta \, d\tau + \frac{1}{4} \int_{-\varepsilon}^t \partial_x (k \circ \eta) (c \circ \eta) \, d\tau + \frac{1}{3} \int_{-\varepsilon}^t \partial_x (z \circ \eta) \, d\tau. \quad (3.15)$$

Identity (3.16) will be used in § 4.3, and (3.14) and (3.15) will be used in § 3.3, 4.3, 4.4, and 10. Similarly, $q^z \circ \psi \psi_x$ satisfies the Duhamel formula

$$\begin{aligned} q^z \circ \psi \psi_x &= (z'_0 + \frac{1}{4} c_0 k'_0) e^{-\int_{-\varepsilon}^t (\frac{8}{3} a + \frac{1}{12} c \partial_\theta k) \circ \psi} - \frac{1}{12} \int_{-\varepsilon}^t e^{-\int_{-\varepsilon}^t (\frac{8}{3} a + \frac{1}{12} c \partial_\theta k) \circ \psi} \psi_x (c \partial_\theta k q^w) \circ \psi \\ &\quad - \frac{8}{3} \int_{-\varepsilon}^t e^{-\int_{-\varepsilon}^t (\frac{8}{3} a + \frac{1}{12} c \partial_\theta k) \circ \psi} \psi_x (\partial_\theta a z) \circ \psi \, d\tau. \end{aligned} \quad (3.16)$$

3.3. Integral bounds

Let $\varphi > 0$, and let Ψ be the flow of $\lambda := (1 - \varphi)w + (\frac{1}{3} + \varphi)z$.

Lemma 3.1. *Suppose that $T \in [-\varepsilon, \varepsilon \wedge T^*]$ and that for all $(\theta, t) \in \mathbb{T} \times [-\varepsilon, T]$ we have*

$$w \sim 1, \quad c \sim 1, \quad |k|, |a| \lesssim 1, \quad |\partial_\theta z| \lesssim \varepsilon^{\beta_1}, \quad |\partial_\theta k| \lesssim \varepsilon^{\gamma_1}, \quad |\partial_\theta a| \lesssim 1.$$

Then, for all $\varphi > 0$, we have

$$\int_{-\varepsilon}^t |\partial_\theta w \circ \Psi(x, \tau)| \, d\tau \lesssim \frac{1}{\varphi} \left(\frac{\varepsilon + t}{\varepsilon} \right) \quad (3.17)$$

with a constant uniform in $\varphi > 0$, $(x, t) \in \mathbb{T} \times [-\varepsilon, T]$.

Proof of Lemma 3.1. Fix $\varphi > 0$ and define

$$g(x, t) := \eta^{-1}(\Psi(x, t), t).$$

We compute that

$$\begin{aligned} \partial_t g(x, t) &= \partial_t \eta^{-1}(\Psi(x, t), t) + \partial_t \eta^{-1}(\Psi(x, t), t) \Psi_t(x, t) \\ &= \frac{-\eta_t(g(x, t), t) + \Psi_t(x, t)}{\eta_x(g(x, t), t)} \\ &= \frac{-\lambda_3 \circ \eta((g(x, t), t) + \lambda \circ \Psi(x, t))}{\eta_x(g(x, t), t)} \\ &= \left(\frac{-\lambda_3 \circ \eta + \lambda \circ \eta}{\eta_x} \right) (g(x, t), t) \\ &= -2\varphi \frac{c \circ \eta}{\eta_x} (g(x, t), t). \end{aligned} \quad (3.18)$$

Note that $\partial_t g(x, t) < 0$ everywhere. We also know that

$$\begin{aligned}\Psi(x, t) &= x + \int_{-\varepsilon}^t \lambda \circ \Psi(x, \tau) \, d\tau, \quad \text{and} \\ \Psi(x, t) &= \eta(g(x, t), t) = g(x, t) + \int_{-\varepsilon}^t \lambda_3 \circ \eta(g(x, t), \tau) \, d\tau, \\ \text{so } x - g(x, t) &= \int_{-\varepsilon}^t \lambda_3 \circ \eta(g(x, t), \tau) - \lambda \circ \Psi(x, \tau) \, d\tau.\end{aligned}$$

Our hypotheses allow us to conclude (see (4.2) below) that

$$I_t(x)e^{-\frac{1}{8}k_0(x)} = 1 + \mathcal{O}(\varepsilon + t)$$

for times $t \in [-\varepsilon, T]$. Using our hypotheses, along with this equation and (3.14), we conclude that for all $(x, t) \in \mathbb{T} \times [-\varepsilon, T]$, we have

$$\sup_{[-\varepsilon, t]} |\eta_x q^w \circ \eta| \leq \varepsilon^{-1}(1 + \mathcal{O}(\varepsilon^{\gamma_1})) + \mathcal{O}(\varepsilon^{0 \wedge \beta_1 + \gamma_1})(\varepsilon + t) \sup_{[-\varepsilon, t]} \eta_x.$$

Plugging this into (3.15) and using the fact that $T \leq \varepsilon$ gives us

$$\begin{aligned}\sup_{[-\varepsilon, t]} \eta_x &\leq 1 + (\varepsilon + t)\varepsilon^{-1}(1 + \mathcal{O}(\varepsilon^{\gamma_1})) + \mathcal{O}(\varepsilon^{\beta_1})(\varepsilon + t) \sup_{[-\varepsilon, t]} \eta_x \\ &\leq 1 + 2(1 + \mathcal{O}(\varepsilon^{\gamma_1})) + \mathcal{O}(\varepsilon^{\beta_1 + 1}) \sup_{[-\varepsilon, t]} \eta_x. \\ \implies \sup_{[-\varepsilon, t]} \eta_x &\leq \frac{1 + 2(1 + \mathcal{O}(\varepsilon^\mu))}{1 - \mathcal{O}(\varepsilon^\mu)} \leq 4.\end{aligned}$$

The last inequality is true for $\varepsilon > 0$ taken to be small enough, since $\mu > 0$. Plugging this into (3.14) and letting ε be sufficiently small gives us

$$\begin{aligned}\eta_x |q^w \circ \eta| &\leq \varepsilon^{-1}(1 + \mathcal{O}(\varepsilon^\mu)) \\ \implies q^w \circ \eta &= \frac{w'_0 I_t e^{-\frac{1}{8}k_0} + \mathcal{O}(\varepsilon^{\gamma_1})}{\eta_x}.\end{aligned}\tag{3.19}$$

It follows that

$$q^w \circ \Psi(x, t) = -\frac{1}{\varphi} \partial_t g(x, t) \frac{w'_0(g(x, t)) + \mathcal{O}(1)}{2c \circ \Psi(x, t)}.$$

Since $\partial_t g < 0$, it follows that

$$|q^w \circ \Psi(x, t)| \lesssim -\frac{1}{\varphi} \partial_t g(x, t) \frac{1}{\varepsilon}.$$

So

$$\int_{-\varepsilon}^t |q^w \circ \Psi(x, \tau)| \, d\tau \lesssim \frac{x - g(x, t)}{\varphi \varepsilon} \lesssim \frac{1}{\varphi} \left(\frac{\varepsilon + t}{\varepsilon} \right).$$

Our result follows immediately from this inequality and our hypotheses. \square

4. Initial Estimates

4.1. Zeroth Order Estimates

Proposition 4.1. For ε small enough, the following estimates hold for all $t \in [-\varepsilon, \varepsilon \wedge T_*]$:

$$w \sim 1, \quad c \sim 1, \quad \phi_x \sim 1, \\ \|\partial_\theta k\|_{L^\infty} \lesssim \|k'_0\|_{L^\infty}, \quad \|a\|_{L^\infty} \leq \|a_0\|_{L^\infty} + \mathcal{O}(\varepsilon), \quad \|z\|_{L^\infty} \leq \|z_0\|_{L^\infty} + \mathcal{O}(\varepsilon).$$

Proof of Proposition 4.1. This follows from an easy bootstrap argument. Let $t \leq \varepsilon \wedge T_*$. If we assume all of the listed bounds hold up to time t for some constants, then it follows that (3.6) holds up to time t . k satisfies $k \circ \phi = k_0$ and

$$\phi_x \partial_\theta k \circ \phi = k'_0. \quad (4.1)$$

Additionally, (2.5) gives us Duhamel formulas for $w \circ \eta$, $z \circ \psi$, and $a \circ \phi$. Using these Duhamel formulas along with (2.11), (3.6), (4.1), and the fact that $t \leq \varepsilon$ it is straightforward to improve our bounds for all times before t , provided the constants we assumed in our bootstrap hypothesis are appropriate and ε is small enough. \square

Using these estimates, it is easy to show that for $\varepsilon > 0$ sufficiently small we obtain that

$$|I_t e^{-\frac{1}{8}k_0} - 1| \leq \mathcal{O}(\varepsilon + t) \quad \forall x \in \mathbb{T}, -\varepsilon \leq t \leq \varepsilon \wedge T_*. \quad (4.2)$$

4.2. $\partial_\theta a$ bounds

Using (3.6) and (4.1) we have

$$\begin{aligned} \partial_t(\varpi \circ \phi) &= \frac{8}{3}(a\varpi) \circ \phi + \frac{4}{3}e^{k_0} \partial_\theta k \circ \phi \\ &= \frac{8}{3}(a\varpi) \circ \phi + \frac{4}{3}e^{k_0} k'_0 \phi_x^{-1} \\ &= \frac{8}{3}(a\varpi) \circ \phi + \frac{4}{3} \frac{k'_0}{c_0^2} e^{k_0} \mathcal{I}_t^2(c \circ \phi)^2, \end{aligned} \quad (4.3)$$

where

$$\mathcal{I}_t := e^{\frac{8}{3} \int_{-\varepsilon}^t a \circ \phi}. \quad (4.4)$$

Therefore,

$$\varpi \circ \phi = \varpi_0 \mathcal{I}_t + \frac{4}{3} c_0^{-2} k'_0 e^{k_0} \mathcal{I}_t \int_{-\varepsilon}^t \mathcal{I}_\tau (c^2 \circ \phi) d\tau. \quad (4.5)$$

Note that

$$\phi_x = c_0^2 \mathcal{I}_t^{-2} c^{-2} \circ \phi. \quad (4.6)$$

This relation will be useful for estimating the higher derivatives of a .

Since

$$\varpi_0 = 4c_0^{-2}(w_0 + z_0 - a'_0)e^{k_0}, \quad (4.7)$$

our assumptions on our initial data let us conclude that $|\varpi_0| \lesssim 1$, and therefore for all $(\theta, t) \in \mathbb{T} \times [-\varepsilon, T]$ we have

$$|\varpi| \lesssim 1. \quad (4.8)$$

Since

$$\partial_{\theta} a = w + z - \frac{1}{4} c^2 \varpi e^{-k}, \quad (4.9)$$

it follows that

$$|\partial_{\theta} a| \lesssim 1 \quad (4.10)$$

for all times $t \in [-\varepsilon, \varepsilon \wedge T_*]$.

Using (4.10) and the bounds on the initial data, we conclude that

$$|\Phi| \lesssim \varepsilon^{-1}, \quad (4.11)$$

for all times $t \in [-\varepsilon, \varepsilon \wedge T_*]$.

4.3. $\partial_{\theta} z$ bounds

Proposition 4.2. *For all $(x, t) \in \mathbb{T} \times [-\varepsilon, \varepsilon \wedge T_*]$ we have*

$$\eta_x |q^w \circ \eta| \leq 2\varepsilon^{-1}, \quad \eta_x \leq 4, \quad |q^z \circ \psi| \lesssim \|z'_0\|_{L^\infty}, \quad \psi_x \sim 1.$$

Proof of 4.2. We will use a bootstrap argument. Let $T \in [-\varepsilon, \varepsilon \wedge T_*)$ and let our bootstrap assumption be that

$$|q^z| \leq C \|z'_0\|_{L^\infty}$$

for all $(\theta, t) \in \mathbb{T} \times [-\varepsilon, T]$ and a constant C to be determined. Since we are assuming $|\partial_{\theta} z| \lesssim \varepsilon^{\beta_1}$ for times $t \in [-\varepsilon, T]$, it follows from (3.3) and our estimates from § 4.1 that $\psi_x \sim 1$ with constants independent of C for times $t \in [-\varepsilon, T]$, provided that ε is small enough relative to C .

Using our bootstrap assumption, along with the estimates from § 4.1 and 4.2, we can conclude (see Lemma 3.1 and its proof) that for all $(x, t) \in \mathbb{T} \times [-\varepsilon, T]$, we have

$$\begin{aligned} \eta_x &\leq 4, \\ \eta_x |q^w \circ \eta| &\leq 2\varepsilon^{-1}, \\ \int_{-\varepsilon}^t |q^w \circ \psi(x, \tau)| d\tau &\lesssim 1. \end{aligned}$$

Using this last estimate along with the estimates from § 4.1, 4.2, it follows from (3.16) and the fact that $\psi_x \sim 1$ that

$$|q^z \circ \psi \psi_x - z'_0| \lesssim \varepsilon^{\gamma_1}$$

for times $t \in [-\varepsilon, T]$. It follows that

$$|q^z \circ \psi| \leq \|\psi_x^{-1}\|_{L^\infty(\mathbb{T} \times [-\varepsilon, T])} (\|z'_0\|_{L^\infty} + \mathcal{O}(\varepsilon^{\gamma_1}))$$

for $t \in [-\varepsilon, T]$. Since $\gamma_1 > 0 \geq \beta_1$, it follows that if we let ε become small enough, we obtain that

$$|q^z| \leq 2 \|\psi_x^{-1}\|_{L^\infty(\mathbb{T} \times [-\varepsilon, T])} \|z'_0\|_{L^\infty}$$

for all $t \in [-\varepsilon, T]$. If C is chosen large enough and ε is chosen small enough, this improves upon our second bootstrap assumption. \square

It follows as an immediate corollary of this proposition that

$$|\partial_{\theta} z| \lesssim \varepsilon^{\beta_1}, \quad (4.12)$$

$$\eta_x \lesssim 1, \quad (4.13)$$

$$\psi_x \sim 1, \quad (4.14)$$

for all times $t \in [-\varepsilon, \varepsilon \wedge T_*]$.

4.4. Bounding $|T_*|$

Now our estimates will let us conclude that η_x behaves roughly the same as it would if w were the solution of Burger's equation with initial data w_0 and η were the flow of w . Using Proposition 4.1, (4.2), (4.10), (4.12), and (4.13) in equation (3.14) gives us

$$\eta_x q^w \circ \eta = \left(-\frac{1}{\varepsilon} + (w'_0 + \frac{1}{\varepsilon})\right) I_t e^{-\frac{1}{\varepsilon} k_0} + \mathcal{O}(\varepsilon^{\gamma_1})$$

for all times $t \in [-\varepsilon, \varepsilon \wedge T_*]$. Plugging this into (3.15) and using the same bounds produces

$$\eta_x = 1 + \left(-\frac{1}{\varepsilon} + (w'_0 + \frac{1}{\varepsilon})\right) \int_{-\varepsilon}^t I_{\tau} e^{-\frac{1}{\varepsilon} k_0} d\tau + \mathcal{O}(\varepsilon^{\beta_1+1}) \quad (4.15)$$

for $t \in [-\varepsilon, \varepsilon \wedge T_*]$. Evaluating (4.15) at $x = 0$ and using (4.2) gives

$$\begin{aligned} \eta_x(0, t) &= 1 - (\varepsilon + t)\varepsilon^{-1} + \mathcal{O}(\varepsilon^{\mu}) \\ &= -\frac{t}{\varepsilon} + \mathcal{O}(\varepsilon^{\mu}). \end{aligned}$$

Since this is true for all $t \in [-\varepsilon, \varepsilon \wedge T_*]$, it follows that we must have $T_* \lesssim \varepsilon^{1+\mu}$ if ε is chosen small enough. Therefore, $T_* = \varepsilon \wedge T_*$, and everything we have proven for $t \in [-\varepsilon, \varepsilon \wedge T_*]$ is true for $t \in [-\varepsilon, T_*]$.

We can also prove a lower bound on T_* . Since $w'_0(x) + \frac{1}{\varepsilon} \geq 0$ for all x , it follows from (4.15) and (4.2) that

$$\eta_x \geq -\frac{t}{\varepsilon} + \mathcal{O}(\varepsilon^{\mu})$$

everywhere. Therefore, $|T_*| \lesssim \varepsilon^{1+\mu}$, else $\partial_{\theta} w, \partial_{\theta} z, \partial_{\theta} k$ and $\partial_{\theta} a$ would all stay bounded up to T_* .

We can also obtain a lower bound for η_x away from 0. Indeed, since $w'_0(x) + \frac{1}{\varepsilon} \geq C\varepsilon^{\frac{\mu}{2}-1}$ for $|x| \geq \varepsilon^{3/2}$, we have

$$\begin{aligned} \eta_x &\geq -\frac{t}{\varepsilon} + C\varepsilon^{\frac{\mu}{2}-1}(\varepsilon + t) + \mathcal{O}(\varepsilon^{\mu}) \\ &= (T_* - t)\left[\frac{t}{\varepsilon} - C\varepsilon^{\frac{\mu}{2}-1}\right] + C\varepsilon^{\frac{\mu}{2}} + \mathcal{O}(\varepsilon^{\mu}) \\ &\geq C\varepsilon^{\frac{\mu}{2}} + \mathcal{O}(\varepsilon^{\mu}) \\ &\gtrsim \varepsilon^{\frac{\mu}{2}}. \end{aligned} \quad (4.16)$$

Using Lemma 3.1 and the estimates proven in this section, we can now conclude that the bound (3.17) holds for all $\varphi > 0, (x, t) \in \mathbb{T} \times [-\varepsilon, T_*]$. This fact will be used so frequently in the rest of the paper that we will not bother to cite it.

5. Transversality

Let $\varphi > 0$ and let Ψ be the flow of $\lambda := (1 - \varphi)w + (\frac{1}{3} + \varphi)z$.

$$\begin{aligned} -\frac{1}{\varphi}\Psi_x\partial_t(\partial_\theta c \circ \Psi) &= \frac{1}{\varphi}\Psi_{xt}\partial_\theta c \circ \Psi - \frac{1}{\varphi}\partial_{tx}(c \circ \Psi) \\ &= \frac{1}{\varphi}\Psi_{xt}\partial_\theta c \circ \Psi + \Psi_x(\partial_\theta w + (\frac{2}{3}\frac{1}{\varphi} - 1)\partial_\theta z + \frac{8}{3}\frac{1}{\varphi}a) \circ \Psi(\partial_\theta c \circ \Psi) \\ &\quad + \Psi_x(c\partial_\theta^2 w + (\frac{2}{3}\frac{1}{\varphi} - 1)c\partial_\theta^2 z + \frac{8}{3}\frac{1}{\varphi}\partial_\theta ac) \circ \Psi. \\ \implies -\frac{1}{\varphi}\partial_t(\Psi_x(c^{-1}\partial_\theta c) \circ \Psi) &= -\frac{1}{\varphi}\Psi_{tx}(c^{-1}\partial_\theta c) \circ \Psi - \Psi_x(c^{-2}\partial_\theta c) \circ \Psi(-\frac{1}{\varphi}\partial_t(c \circ \Psi)) \\ &\quad + (c^{-1} \circ \Psi)(-\frac{1}{\varphi}\Psi_x\partial_t(\partial_\theta c \circ \Psi)) \\ &= \Psi_x(\partial_\theta^2 w + (\frac{2}{3}\frac{1}{\varphi} - 1)\partial_\theta^2 z + \frac{8}{3}\frac{1}{\varphi}\partial_\theta a) \circ \Psi. \end{aligned}$$

Therefore, if $h : \mathbb{T} \times [-\varepsilon, T_*) \rightarrow \mathbb{R}$ is any differentiable function, we have

$$\begin{aligned} \Psi_x(h\partial_\theta^2 w) \circ \Psi &= -\frac{1}{\varphi}\partial_t(\Psi_x(c^{-1}\partial_\theta ch) \circ \Psi) - \Psi_x((\frac{2}{3}\frac{1}{\varphi} - 1)h\partial_\theta^2 z + \frac{8}{3}\frac{1}{\varphi}\partial_\theta ah) \circ \Psi \\ &\quad - \Psi_x(c^{-1}\partial_\theta c) \circ \Psi(-\frac{1}{\varphi}\partial_t(h \circ \Psi)). \end{aligned}$$

This gives us the following equation:

$$\begin{aligned} \partial_x((h\partial_\theta w) \circ \Psi) &= -\frac{1}{\varphi}\partial_t(\Psi_x(c^{-1}\partial_\theta ch) \circ \Psi) \\ &\quad - \Psi_x((\frac{2}{3}\frac{1}{\varphi} - 1)h\partial_\theta^2 z + \frac{8}{3}\frac{1}{\varphi}\partial_\theta ah) \circ \Psi \\ &\quad + \Psi_x[(\partial_\theta h\partial_\theta w) \circ \Psi - (c^{-1}\partial_\theta c) \circ \Psi(-\frac{1}{\varphi}\partial_t(h \circ \Psi))]. \end{aligned} \quad (5.1)$$

The last term in this expression motivates the following definition:

Definition 5.1 (Transversality). *A differentiable function $h : \mathbb{T} \times [-\varepsilon, T_*) \rightarrow \mathbb{R}$ is **transversal** (or **1-transversal**) if it is bounded and there exist both a constant $\varphi > 0$ and bounded functions A, B , and C such that*

$$\begin{cases} \partial_\theta h = Ac^{-1}\partial_\theta c + B \\ \partial_t h + \lambda\partial_\theta h = -\varphi A\partial_\theta w - \varphi C \end{cases}$$

Here $\lambda = (1 - \varphi)w + (\frac{1}{3} + \varphi)z$, as in the above discussion. If in addition A, B , and C are themselves transversal functions, we say that h is **2-transversal**. We recursively define h to be **n -transversal** if A, B , and C are $(n - 1)$ -transversal.

A few remarks about transversal functions:

- If h satisfies the transversality condition for one $\varphi_0 > 0$, then it satisfies the transversality condition for all $\varphi > 0$. If indeed, if we have

$$\begin{cases} \partial_\theta h = Ac^{-1}\partial_\theta c + B \\ \partial_t h + \lambda_0\partial_\theta h = -\varphi_0 A\partial_\theta w - \varphi_0 C \end{cases}$$

for some $\varphi_0 > 0$ then for any other $\varphi > 0$ we have

$$\begin{cases} \partial_\theta h = Ac^{-1}\partial_\theta c + B \\ \partial_t h + \lambda\partial_\theta h = -\varphi A\partial_\theta w + (\varphi - \varphi_0)A\partial_\theta z + 2(\varphi_0 - \varphi)cB - \varphi_0 C \end{cases}.$$

Since $\partial_{\theta}z$ is bounded, h still satisfies the transversality condition for φ , albeit with a different choice of bounded function C . So the notion of a transversal function is independent of our choice of $\varphi > 0$.

- Note that while being transversal does not depend on the choice of φ (as the previous bullet illustrated), and A and B are independent of φ , the function C changes based on φ .
- If h is a bounded function with bounded derivatives, then h is trivially transversal, with $A = 0$, $B = \partial_{\theta}h$ and $C = -\frac{1}{\varphi}(\partial_t h + \lambda \partial_{\theta} h)$.
- If functions h_1, h_2 are n -transversal, then $h_1 + h_2$ is n -transversal. Indeed, we have

$$\begin{cases} \partial_{\theta}(h_1 + h_2) = (A_1 + A_2)c^{-1}\partial_{\theta}c + B_1 + B_2 \\ (\partial_t + \lambda\partial_{\theta})(h_1 + h_2) = -\varphi(A_1 + A_2)\partial_{\theta}w - \varphi(C_1 + C_2) \end{cases} .$$

- If functions h_1, h_2 are n -transversal, then their product is n -transversal. Indeed, we have

$$\begin{cases} \partial_{\theta}(h_1 h_2) = (A_1 h_2 + A_2 h_1)c^{-1}\partial_{\theta}c + B_1 h_2 + B_2 h_1 \\ (\partial_t + \lambda\partial_{\theta})(h_1 h_2) = -\varphi(A_1 h_2 + A_2 h_1)\partial_{\theta}w - \varphi(C_1 h_2 + C_2 h_1) \end{cases} .$$

- If h is n -transversal and $h \sim 1$ then h^{-1} is also n -transversal. Indeed,

$$\begin{cases} \partial_{\theta}(h^{-1}) = -h^{-2}Ac^{-1}\partial_{\theta}c - h^{-2}B \\ \partial_t(h^{-1}) + \lambda\partial_{\theta}(h^{-1}) = -\varphi(-h^{-2}A)\partial_{\theta}w - \varphi(-h^{-2}C) \end{cases} .$$

- If $F : \mathbb{R} \rightarrow \mathbb{R}$ is smooth and h is n -transversal, then $F \circ h$ is n -transversal. Indeed, we have

$$\begin{cases} \partial_{\theta}(F \circ h) = (AF' \circ h)c^{-1}\partial_{\theta}c + BF' \circ h \\ (\partial_t + \lambda\partial_{\theta})(F \circ h) = -\varphi(AF' \circ h)\partial_{\theta}w - \varphi CF' \circ h \end{cases} .$$

This rule will be especially useful for $F(x) = e^x$.

- c is transversal with $A = c$, $B = 0$, and $C = 4ac$ when $\varphi = \frac{2}{3}$. It follows inductively that if a is n -transversal, then c is $(n + 1)$ -transversal. At this point, we already know that a is at least 1-transversal because it is uniformly C^1 , so c is currently proven to be at least 2-transversal. $c \sim 1$, so c^{-1} is also 2-transversal. The fact that both c and c^{-1} are transversal was the main ingredient used in the computation of (5.1).

The following lemma will be used in § 7.3, § 8.3, and § 9.3.

Lemma 5.2 (Identities for transversal functions along 1-characteristics). *If $h : \mathbb{T} \times [-\varepsilon, T_*) \rightarrow \mathbb{R}$ is transversal with*

$$\begin{cases} \partial_{\theta}h = Ac^{-1}\partial_{\theta}c + B \\ -\frac{3}{2}\partial_t(h \circ \psi) = (A\partial_{\theta}w + C) \circ \psi \end{cases}$$

then we have

$$\partial_x((h\partial_{\theta}w) \circ \psi) = -\frac{3}{2}\partial_t(\psi_x(c^{-1}\partial_{\theta}ch) \circ \psi) + \psi_x([B - \frac{1}{2}c^{-1}C]\partial_{\theta}w) \circ \psi + \psi_x(\frac{1}{2}c^{-1}C\partial_{\theta}z - 4\partial_{\theta}ah) \circ \psi. \quad (5.2)$$

and

$$\begin{aligned} \partial_x(\psi_x(h\partial_\theta w)\circ\psi) &= -\frac{3}{2}\partial_t(\psi_x^2(c^{-1}\partial_\theta ch)\circ\psi) + \psi_x^2([Qh + B - \frac{1}{2}c^{-1}C + \frac{3}{4}c^{-1}h\partial_\theta z]\partial_\theta w)\circ\psi \\ &\quad + \psi_x^2(\frac{1}{2}c^{-1}C\partial_\theta z - 4\partial_\theta ah - \frac{3}{4}c^{-1}h\partial_\theta z^2)\circ\psi. \end{aligned} \quad (5.3)$$

From these two equations. we obtain the bounds

$$|\partial_x((h\partial_\theta w)\circ\psi) + \frac{3}{2}\partial_t(\psi_x(c^{-1}\partial_\theta ch)\circ\psi)| \lesssim \|h\|_{L^\infty} + \varepsilon^{\beta_1}\|C\|_{L^\infty} + (\|B\|_{L^\infty} + \|C\|_{L^\infty})|\partial_\theta w\circ\psi|. \quad (5.4)$$

and

$$\begin{aligned} |\partial_x(\psi_x(h\partial_\theta w)\circ\psi) + \frac{3}{2}\partial_t(\psi_x^2(c^{-1}\partial_\theta ch)\circ\psi)| &\lesssim \varepsilon^{2\beta_1}\|h\|_{L^\infty} + \varepsilon^{\beta_1}\|C\|_{L^\infty} \\ &\quad + (\varepsilon^{-1}\|h\|_{L^\infty} + \|B\|_{L^\infty} + \|C\|_{L^\infty})|\partial_\theta w\circ\psi|. \end{aligned} \quad (5.5)$$

Proof of Lemma 5.2. (5.2) follows immediately from (5.1). To prove (5.3),

$$\begin{aligned} \partial_x(\psi(h\partial_\theta w)\circ\psi) &= \psi_{xx}(h\partial_\theta w)\circ\psi + \psi_x\partial_x((h\partial_\theta w)\circ\psi) \\ &= \psi_x^2([Q - \frac{1}{2}c^{-1}\partial_\theta c]h\partial_\theta w)\circ\psi - \partial_t(\psi_x^2(c^{-1}\partial_\theta ch)\circ\psi) + \frac{3}{2}\psi_{xt}\psi_x(c^{-1}\partial_\theta ch)\circ\psi \\ &\quad + \psi_x([B - \frac{1}{2}c^{-1}C]\partial_\theta w)\circ\psi + \psi_x(\frac{1}{2}c^{-1}C\partial_\theta z - 4\partial_\theta ah)\circ\psi \\ &= -\frac{3}{2}\partial_t(\psi_x^2(c^{-1}\partial_\theta ch)\circ\psi) + \psi_x^2([Qh + B - \frac{1}{2}c^{-1}C + \frac{3}{4}c^{-1}h\partial_\theta z]\partial_\theta w)\circ\psi \\ &\quad + \psi_x^2(\frac{1}{2}c^{-1}C\partial_\theta z - 4\partial_\theta ah - \frac{3}{4}c^{-1}h\partial_\theta z^2)\circ\psi. \end{aligned}$$

The inequalities follow immediately from the equations and the first-order estimates. \square

The following lemma will be used in § 7.2, § 8.2, and § 9.2.

Lemma 5.3 (Identities for transversal functions along 2-characteristics). *If $h : \mathbb{T} \times [-\varepsilon, T_*) \rightarrow \mathbb{R}$ is a differentiable function satisfying the transversality condition*

$$\begin{cases} \partial_\theta h = Ac^{-1}\partial_\theta c + B \\ -3\partial_t(h\circ\phi) = (A\partial_\theta w + C)\circ\phi \end{cases}$$

then we have

$$\partial_x((h\partial_\theta w)\circ\phi) = -3\partial_t(\phi_x(c^{-1}\partial_\theta ch)\circ\phi) + \phi_x(B\partial_\theta w - Cc^{-1}\partial_\theta c - h\partial_\theta^2 z - 8\partial_\theta ah)\circ\phi. \quad (5.6)$$

and

$$\begin{aligned} \partial_x(\phi_x^2(h\partial_\theta w)\circ\phi) &= -3\partial_t(\phi_x^3(c^{-1}\partial_\theta c\partial_\theta w)\circ\phi) \\ &\quad + \phi_x^3(B\partial_\theta w - Cc^{-1}\partial_\theta c + 4c^{-1}\partial_\theta ch\partial_\theta z - h\partial_\theta^2 z - 8\partial_\theta ah)\circ\phi \\ &\quad + 2\phi_x^2\Phi(h\partial_\theta w)\circ\phi. \end{aligned} \quad (5.7)$$

Proof of Lemma 5.3. (5.6) follows immediately from (5.1). The proof of (5.7) is an easy computation using (5.1) and (3.7). \square

The following lemma will first be used in § 8.3, so there is no circularity in its proof. See § 6.1 for the definition of \mathcal{E} and § 6.3 for the definition of f .

Lemma 5.4 (Identities for 2-transversal functions along 1-characteristics). *If $h : \mathbb{T} \times [-\varepsilon, T_*) \rightarrow \mathbb{R}$ is 2-transversal with*

$$\begin{aligned} \partial_\theta h &= Ac^{-1}\partial_\theta c + B \\ -\frac{3}{2}\partial_t(h \circ \psi) &= (A\partial_\theta w + C) \circ \psi \\ \partial_\theta A &= A_A c^{-1}\partial_\theta c + B_A \\ -\frac{3}{2}\partial_t(A \circ \psi) &= (A_A \partial_\theta w + C_A) \circ \psi \\ \partial_\theta B &= A_B c^{-1}\partial_\theta c + B_B \\ -\frac{3}{2}\partial_t(B \circ \psi) &= (A_B \partial_\theta w + C_B) \circ \psi \\ \partial_\theta C &= A_C c^{-1}\partial_\theta c + B_C \\ -\frac{3}{2}\partial_t(C \circ \psi) &= (A_C \partial_\theta w + C_C) \circ \psi \end{aligned}$$

then we have

$$\begin{aligned} \partial_x^2((h\partial_\theta w) \circ \psi) &= -\frac{3}{2}\partial_t(\psi_x^2(c^{-1}\partial_\theta^2 ch + [-\frac{3}{2}h + A]c^{-2}\partial_\theta c^2 + [Q_1 + 2B - \frac{1}{2}c^{-1}C]c^{-1}\partial_\theta c) \circ \psi) \\ &+ \psi_x^2([BQ_1 + B_B - \frac{1}{2}c^{-1}B_C]\partial_\theta w) \circ \psi \\ &- \psi_x^2([Q_1C + C_B - \frac{1}{2}c^{-1}C_C + 2ac^{-1}C + \frac{1}{2}c^{-1}\partial_\theta zC - 2\partial_\theta ah + 4\partial_\theta aA]c^{-1}\partial_\theta c) \circ \psi \\ &- \psi_x^2([\frac{1}{4}c^{-1}\partial_\theta zC - \frac{1}{2}A_C c^{-1}\partial_\theta z + \frac{1}{4}C\partial_\theta k]c^{-1}\partial_\theta c) \circ \psi \\ &+ \psi_x^2(\frac{1}{2}f - 4\partial_\theta ahQ_1 - 4\partial_\theta^2 ah - 4\partial_\theta aB - \frac{1}{8}c\mathcal{E} + \frac{1}{2}B_C c^{-1}\partial_\theta z) \circ \psi \end{aligned} \quad (5.8)$$

and

$$\begin{aligned} \partial_x^2(h \circ \psi) &= -\frac{3}{4}\partial_t(\psi_x^2(Ac^{-2}\partial_\theta c) \circ \psi) \\ &+ \psi_x^2([\frac{1}{2}Ac^{-1}Q_1 + \frac{1}{2}B_A c^{-1} - \frac{1}{4}C_A c^{-2} + Aac^{-2} + \frac{3}{8}Ac^{-2}\partial_\theta z]\partial_\theta w) \circ \psi \\ &+ \psi_x^2([\frac{1}{4}Ac^{-1}\partial_\theta z - \frac{1}{2}B + A_B - \frac{1}{2}A_A c^{-1}\partial_\theta + \frac{1}{4}Ac^{-1}\partial_\theta z + \frac{1}{4}A\partial_\theta k]c^{-1}\partial_\theta c) \circ \psi \\ &+ \psi_x^2(\frac{1}{4}C_A c^{-2}\partial_\theta z - Aac^{-2}\partial_\theta z - 2\partial_\theta aAc^{-1} - \frac{3}{8}Ac^{-2}\partial_\theta z^2) \circ \psi \\ &+ \psi_x^2(BQ_1 - \frac{1}{2}Ac^{-1}Q_1\partial_\theta z + B_B - \frac{1}{2}B_A c^{-1}\partial_\theta z - \frac{1}{2}Ac^{-1}f + \frac{1}{8}A\mathcal{E}) \circ \psi. \end{aligned} \quad (5.9)$$

Proof of Lemma 5.4. Taking ∂_x of (5.2) gives us

$$\begin{aligned} \partial_x^2((h\partial_\theta w) \circ \psi) &= -\frac{3}{2}\partial_{tx}^2(\psi_x(c^{-1}\partial_\theta c) \circ \psi) + \partial_x(\psi_x([B - \frac{1}{2}c^{-1}C]\partial_\theta w) \circ \psi) \\ &+ \partial_x(\psi_x(\frac{1}{2}c^{-1}C\partial_\theta z - 4\partial_\theta ah) \circ \psi). \end{aligned}$$

If we define $\tilde{h} := B - \frac{1}{2}c^{-1}C$, then the rules for transversal functions tell us that

$$\begin{aligned} B_{\tilde{h}} &= B_B - \frac{1}{2}c^{-1}B_C \\ C_{\tilde{h}} &= C_B - \frac{1}{2}c^{-1}C_C + 2ac^{-1}C. \end{aligned}$$

Applying (5.3) to \tilde{h} and simplifying gives us (5.8).

For the next identity, we see that

$$\partial_x^2(h \circ \psi) = \partial_x(\psi_x(\frac{1}{2}Ac^{-1}\partial_\theta w) \circ \psi) + \partial_x(\psi_x(B - \frac{1}{2}Ac^{-1}\partial_\theta z) \circ \psi).$$

Applying (5.3) to the function $\frac{1}{2}Ac^{-1}$ and simplifying gives us (5.9). \square

The following lemma will be used in § 8 and § 9.

Lemma 5.5 (Classes of transversal functions). *Let $\varphi > 0$, and let Ψ be the flow of $\lambda = (1 - \varphi)w + (\frac{1}{3} + \varphi)z$. Then*

1. *If h is a transversal function and H is defined by*

$$H \circ \Psi(x, t) := \int_{-\varepsilon}^t h \circ \Psi(x, \tau) d\tau$$

then H is transversal.

2. $\Psi_x \circ \Psi^{-1}$ *is a transversal function.*

3. *If h is a transversal function and K is defined by*

$$K \circ \Psi(x, t) := \int_{-\varepsilon}^t (h \partial_{\theta} w) \circ \Psi(x, \tau) d\tau$$

then K is transversal.

4. *If h is a 2-transversal function and H is defined by*

$$H \circ \Psi(x, t) := \int_{-\varepsilon}^t h \circ \Psi(x, \tau) d\tau$$

then H is 2-transversal.

5. $\Psi_x \circ \Psi^{-1}$ *is a 2-transversal function.*

6. *If h is a 2-transversal function and K is defined by*

$$K \circ \Psi(x, t) := \int_{-\varepsilon}^t (h \partial_{\theta} w) \circ \Psi(x, \tau) d\tau$$

then K is 2-transversal.

Proof of Lemma 5.5. In this proof, h satisfies

$$\begin{cases} \partial_{\theta} h = Ac^{-1} \partial_{\theta} c + B \\ -\frac{1}{\varphi} \partial_t (h \circ \Psi) = (A \partial_{\theta} w + C) \circ \Psi \end{cases} .$$

(i) Since

$$\begin{cases} \partial_{\theta} H = \left(\Psi_x^{-1} \int_{-\varepsilon}^t (Ac^{-1} \partial_{\theta} c + B) \circ \Psi d\tau \right) \circ \Psi^{-1} \\ -\frac{1}{\varphi} \partial_t (H \circ \Psi) = -\frac{1}{\varphi} h \circ \Psi \end{cases} \quad (5.10)$$

it follows from (3.17) and the fact that $\Psi_x \sim 1$ that H is transversal.

(ii) We know that a is transversal, and it will be proven in § 6.3 that $\partial_{\theta} z$ is transversal. Therefore, part (i) applies to the function

$$H \circ \Psi = \int_{-\varepsilon}^t \left(2 - \frac{2}{3} \frac{1}{\varphi} \right) \partial_{\theta} z - \frac{8}{3} \frac{1 - \varphi}{\varphi} a \circ \Psi.$$

Since $F(x) = e^x$ is smooth, it follows that

$$e^{\int_{-\varepsilon}^t (2 - \frac{2}{3}\frac{1}{\varphi}) \partial_{\theta z} - \frac{8}{3}\frac{1-\varphi}{\varphi} a) \circ \Psi \circ \Psi^{-1}}$$

is transversal. We already know that c and c^{-1} are both 2-transversal, so it now follows from (3.2) that $\Psi_x \circ \Psi^{-1}$ is transversal.

(iii) Using (5.1) tells us that

$$\begin{aligned} \partial_{\theta} K \circ \psi &= -\frac{1}{\varphi} (hc^{-1} \partial_{\theta} c) \circ \Psi + \frac{1}{\varphi} \Psi_x^{-1} c_0^{-1} c_0' h_0 \\ &\quad - \Psi_x^{-1} \int_{-\varepsilon}^t \Psi_x \left(\left(\frac{2}{3} \frac{1}{\varphi} - 1 \right) h \partial_{\theta}^2 z + \frac{8}{3} \frac{1}{\varphi} \partial_{\theta} a h \right) \circ \Psi \, d\tau + \Psi_x^{-1} \int_{-\varepsilon}^t \Psi_x (B \partial_{\theta} w - c^{-1} \partial_{\theta} c C) \circ \Psi \, d\tau, \\ -\frac{1}{\varphi} \partial_t (K \circ \Psi) &= -\frac{1}{\varphi} (h \partial_{\theta} w) \circ \Psi. \end{aligned} \quad (5.11)$$

It now follows from (6.18) that K is transversal.

(iv) This follows immediately from applying (ii) and (iii) to (5.10).

(v) It will be proven in § 6.2 that $\partial_{\theta} a$ is transversal, from which it will follow that a is 2-transversal, and it will be proven in § 7.3 that $\partial_{\theta} z$ is 2-transversal. Since $F(x) = e^x$ is smooth, it follows from (iv) that

$$e^{\int_{-\varepsilon}^t (2 - \frac{2}{3}\frac{1}{\varphi}) \partial_{\theta z} - \frac{8}{3}\frac{1-\varphi}{\varphi} a) \circ \Psi \circ \Psi^{-1}}$$

is 2-transversal. Since c^{-1} is 2-transversal, it now follows from (3.2) that $\Psi_x \circ \Psi^{-1}$ is 2-transversal.

(vi) We prove in § 6.2, 6.3, 7.1, and 7.3 that $\partial_{\theta} a$, $\partial_{\theta} z$, \mathcal{E} , and f are all transversal, so our result follows from applying (i), (ii), and (iii) to (5.11). □

6. Second Derivative Estimates

6.1. $\partial_{\theta}^2 k$ bounds

Differentiating (4.1) and plugging in (3.7) gives us

$$\begin{aligned} \phi_x^2 \partial_{\theta}^2 k \circ \phi &= k_0'' - \phi_{xx} \partial_{\theta} k \circ \phi \\ &= k_0'' - \phi_x \Phi \partial_{\theta} k \circ \phi + 2\phi_x^2 \frac{\partial_{\theta} c \circ \phi \partial_{\theta} k \circ \phi}{c \circ \phi} \\ &= k_0'' - \Phi k_0' + 2\phi_x^2 \frac{\partial_{\theta} c \circ \phi \partial_{\theta} k \circ \phi}{c \circ \phi}. \end{aligned} \quad (6.1)$$

If we define

$$\mathcal{E} := \partial_{\theta}^2 k - 2c^{-1} \partial_{\theta} c \partial_{\theta} k, \quad (6.2)$$

then it follows that $\mathcal{E} = [\phi_x^{-2} (k_0'' - \Phi k_0')] \circ \phi^{-1}$ and from (4.11) we conclude that

$$|\mathcal{E}| \lesssim \varepsilon^{\gamma_2 \wedge \gamma_1 - 1}. \quad (6.3)$$

$$\implies |\partial_{\theta}^2 k| \lesssim \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} + \varepsilon^{\gamma_1} |\partial_{\theta} w|. \quad (6.4)$$

With this notation, we can write

$$\begin{aligned}\partial_\theta^2 k &= (2\partial_\theta k)c^{-1}\partial_\theta c + \mathcal{E} \\ -\frac{3}{2}\partial_t(\partial_\theta k \circ \psi) &= (2\partial_\theta k)\partial_\theta w + c\mathcal{E},\end{aligned}\tag{6.5}$$

so $\partial_\theta k$ is transversal. The fact that $\partial_\theta k$ is transversal will be used through § 7 and § 6.

6.2. $\partial_\theta^2 a$ bounds

Using (3.6) and (A.1a) we have

$$\begin{aligned}\mathcal{I}_t \partial_x(c^2 \circ \phi) &= 2\phi_x \mathcal{I}_t \left[-\frac{3}{2}\partial_t(c \circ \phi) - (4ac + c\partial_\theta z) \circ \phi \right] \\ &= c_0^2 \mathcal{I}_t^{-1} [3\partial_t(c^{-1} \circ \phi) - (8c^{-1}a + 2c^{-1}\partial_\theta z) \circ \phi] \\ &= c_0^2 [3\partial_t(\mathcal{I}_t^{-1} c^{-1} \circ \phi) - 2\mathcal{I}_t^{-1}(c^{-1}\partial_\theta z) \circ \phi].\end{aligned}\tag{6.6}$$

Therefore, differentiating (4.5) gives us

$$\begin{aligned}\partial_x(\varpi \circ \phi) &= \partial_x(\varpi_0 \mathcal{I}_t) - 4\frac{k'_0 e^{k_0}}{c_0} \mathcal{I}_t + 3\frac{k'_0 e^{k_0}}{c \circ \phi} - \frac{8}{3}k'_0 e^{k_0} \mathcal{I}_t \int_{-\varepsilon}^t \mathcal{I}_\tau^{-1}(c^{-1}\partial_\theta z) \circ \phi \, d\tau \\ &\quad + \frac{4}{3}\partial_x\left(\frac{k'_0 e^{k_0}}{c_0^2} \mathcal{I}_t\right) \int_{-\varepsilon}^t \mathcal{I}_\tau(c^2 \circ \phi) \, d\tau + \frac{4}{3}\frac{k'_0 e^{k_0}}{c_0^2} \mathcal{I}_t \int_{-\varepsilon}^t \partial_x(\mathcal{I}_\tau)(c^2 \circ \phi) \, d\tau.\end{aligned}\tag{6.7}$$

It is easy to check that

$$|\mathcal{I}'_t| \lesssim \varepsilon.\tag{6.8}$$

It follows from this bound and (6.7) that

$$|\partial_\theta \varpi \circ \phi| \lesssim |\varpi'_0| + \varepsilon^{2+1\wedge\gamma_1}.$$

By differentiating the equation (4.7) and using our assumptions on the initial data, we conclude that $|\varpi'_0| \lesssim \varepsilon^{-1}$. Therefore,

$$|\partial_\theta \varpi| \lesssim \varepsilon^{-1}.\tag{6.9}$$

Differentiating (4.9) in space and using our first derivative estimates along with (6.9) gives us

$$\begin{aligned}\partial_\theta^2 a &= 2[\partial_\theta a - c - 2z]c^{-1}\partial_\theta c + 2\partial_\theta z - \frac{1}{4}c^2\partial_\theta \varpi e^{-k} + \frac{1}{4}c^2\partial_\theta k \varpi e^{-k} \\ -\frac{3}{2}\partial_t(\partial_\theta a \circ \psi) &= (2[\partial_\theta a - c - 2z]\partial_\theta w + 4a\partial_\theta a - \frac{1}{4}c^3\partial_\theta \varpi e^{-k} + \frac{1}{4}c^3\partial_\theta k \varpi e^{-k}) \circ \psi.\end{aligned}\tag{6.10}$$

So (6.9) lets us conclude that $\partial_\theta a$ is transversal, which will be used in § 7 and § 8. This equation for $\partial_\theta^2 a$ and our estimate (6.9) also lets us conclude that

$$|\partial_\theta^2 a| \lesssim \varepsilon^{-1} + |\partial_\theta w|.\tag{6.11}$$

It now follows from (6.11) that

$$|\partial_x \Phi| \lesssim \varepsilon^{-2} + |w'_0|.\tag{6.12}$$

6.3. $\partial_\theta^2 z$ bounds

Let us introduce the new variable

$$f := \partial_\theta^2 z - \frac{1}{2}c^{-1}\partial_\theta c \partial_\theta z + \frac{1}{4}c\partial_\theta^2 k. \quad (6.13)$$

Using the identities from § A.1 along with (6.2) gives us

$$\begin{aligned} \partial_t(f \circ \psi) &:= (\partial_t(\log c \circ \psi) + \frac{1}{2}\partial_t(k \circ \psi) - 3\partial_\theta z \circ \psi)(f \circ \psi) \\ &+ (\frac{1}{12}c^2\partial_\theta k \partial_\theta^2 k - \frac{1}{2}c^{-1}\partial_\theta c \partial_\theta z^2) \circ \psi - \frac{1}{8}\mathcal{E}(\frac{1}{3}c\partial_\theta w - c\partial_\theta z) \circ \psi \\ &- \frac{8}{3}(\partial_\theta^2 a z + \frac{3}{2}\partial_\theta a \partial_\theta z - \frac{1}{2}\partial_\theta a c^{-1}\partial_\theta c z) \circ \psi. \end{aligned} \quad (6.14)$$

If we define

$$J_t := e^{\frac{1}{2}k \circ \psi - 3 \int_{-\varepsilon}^t \partial_\theta z \circ \psi}, \quad (6.15)$$

then (6.14) gives us the Duhamel formula

$$\begin{aligned} f \circ \psi &= c_0^{-1}(z_0'' - \frac{1}{2}c_0^{-1}c_0'z_0' + \frac{1}{2}c_0k_0'')e^{-\frac{1}{2}k_0}J_t c \circ \psi + \frac{1}{12}c \circ \psi J_t \int_{-\varepsilon}^t J_\tau^{-1}(c\partial_\theta k \partial_\theta^2 k) \circ \psi \, d\tau \\ &- \frac{1}{2}c \circ \psi J_t \int_{-\varepsilon}^t J_\tau^{-1}(c^{-2}\partial_\theta c \partial_\theta z^2) \circ \psi \, d\tau - \frac{1}{8}c \circ \psi J_t \int_{-\varepsilon}^t J_\tau^{-1}\mathcal{E}(\frac{1}{3}\partial_\theta w - \partial_\theta z) \circ \psi \, d\tau \\ &- \frac{8}{3}c \circ \psi J_t \int_{-\varepsilon}^t J_\tau^{-1}(\partial_\theta^2 a c^{-1}z + \frac{3}{2}\partial_\theta a c^{-1}\partial_\theta z - \frac{1}{2}\partial_\theta a c^{-2}\partial_\theta c z) \circ \psi \, d\tau. \end{aligned} \quad (6.16)$$

It now follows from (6.4), (6.3), and (6.11) that

$$|f| \lesssim \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1}. \quad (6.17)$$

It follows immediately from this bound and (6.4) that

$$|\partial_\theta^2 z| \lesssim \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1} + \varepsilon^{\beta_1} |\partial_\theta w|. \quad (6.18)$$

This bound tells us that

$$|\Psi| \lesssim \varepsilon^{-1}. \quad (6.19)$$

We can also conclude that $\partial_\theta z$ is transversal. Indeed

$$\begin{aligned} \partial_\theta^2 z &= [\frac{1}{2}\partial_\theta z - \frac{1}{2}c\partial_\theta k]c^{-1}\partial_\theta c + f - \frac{1}{4}c\mathcal{E} \\ -\frac{3}{2}\partial_t(\partial_\theta z \circ \psi) &= ([\frac{1}{2}\partial_\theta z - \frac{1}{2}c\partial_\theta k]\partial_\theta w + \frac{1}{2}c\partial_\theta k \partial_\theta z - \frac{1}{4}c^2\mathcal{E} + 4\partial_\theta a z + a\partial_\theta z) \circ \psi. \end{aligned} \quad (6.20)$$

The fact that $\partial_\theta z$ is transversal will be used in § 7 and § 8.

7. Third derivative estimates

7.1. $\partial_\theta^3 k$ bounds

Using the fact that $\mathcal{E} = [\phi_x^{-2}(k_0'' - \Phi k_0')] \circ \phi^{-1}$, we can compute that

$$\partial_\theta \mathcal{E} = [\phi_x^{-3}(k_0''' - 3\Phi k_0'' + (2\Phi^2 - \partial_x \Phi)k_0')] \circ \phi^{-1} + 4\mathcal{E}c^{-1}\partial_\theta c \quad (7.1)$$

Define

$$\tilde{\mathcal{E}} := [\phi_x^{-3}(k_0''' - 3\Phi k_0'' + (2\Phi^2 - \partial_x \Phi)k_0')] \circ \phi^{-1}. \quad (7.2)$$

We know from (4.11) and (6.12) that

$$|\tilde{\mathcal{E}}| \lesssim \varepsilon^{\gamma_1} |w_0'' \circ \phi^{-1}| + \varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \gamma_1 - 2}. \quad (7.3)$$

Since

$$\begin{aligned} \partial_\theta \mathcal{E} &= 4\mathcal{E}c^{-1} \partial_\theta c + \tilde{\mathcal{E}} \\ -\frac{3}{2} \partial_t (\mathcal{E} \circ \psi) &= (4\mathcal{E} \partial_\theta w - 8\partial_\theta a \partial_\theta k + \tilde{\mathcal{E}}c) \circ \psi. \end{aligned} \quad (7.4)$$

it follows that \mathcal{E} is transversal and therefore $\partial_\theta k$ is 2-transversal.

Taking ∂_θ of (6.2) gives us

$$\partial_\theta^3 k = \tilde{\mathcal{E}} + 6\mathcal{E}c^{-1} \partial_\theta c + 2c^{-2} \partial_\theta c^2 \partial_\theta k + 2c^{-1} \partial_\theta^2 c \partial_\theta k. \quad (7.5)$$

It follows that

$$\begin{aligned} |\partial_\theta^3 k| &\lesssim \varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \gamma_1 - 2} + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} |\partial_\theta w| \\ &\quad + \varepsilon^{\gamma_1} |w_0'' \circ \phi^{-1}| + \varepsilon^{\gamma_1} |\partial_\theta w|^2 + \varepsilon^{\gamma_1} |\partial_\theta^2 w|. \end{aligned} \quad (7.6)$$

7.2. $\partial_\theta^3 a$ bounds

Taking ∂_x of (6.7) gives us

$$\begin{aligned} \partial_x^2 (\varpi \circ \phi) &= \partial_x^2 (\varpi_0 I_t) - 4\partial_x \left(\frac{k_0' e^{k_0}}{c_0} I_t \right) + 4 \frac{\partial_x (k_0' e^{k_0})}{c \circ \phi} - 4k_0' e^{k_0} \phi_x \frac{\partial_\theta c \circ \phi}{(c \circ \phi)^2} \\ &\quad - \frac{8}{3} \partial_x (k_0' e^{k_0} I_t) \int_{-\varepsilon}^t I_\tau^{-1} (c^{-1} \partial_\theta z) \circ \phi \, d\tau - \frac{8}{3} k_0' e^{k_0} I_t \int_{-\varepsilon}^t \partial_x (I_\tau^{-1} (c^{-1} \partial_\theta z) \circ \phi) \, d\tau \\ &\quad + \frac{4}{3} \partial_x^2 \left(\frac{k_0' e^{k_0}}{c_0^2} I_t \right) \int_{-\varepsilon}^t I_\tau (c^2 \circ \phi) \, d\tau + \frac{8}{3} \partial_x \left(\frac{k_0' e^{k_0}}{c_0^2} I_t \right) \int_{-\varepsilon}^t \partial_x (I_\tau) (c^2 \circ \phi) \, d\tau \\ &\quad + \frac{4}{3} \partial_x \left(\frac{k_0' e^{k_0}}{c_0^2} I_t \right) \int_{-\varepsilon}^t I_\tau \partial_x (c^2 \circ \phi) \, d\tau + \frac{4}{3} \frac{k_0' e^{k_0}}{c_0^2} I_t \int_{-\varepsilon}^t \partial_x (I_\tau) \partial_x (c^2 \circ \phi) \, d\tau \\ &\quad + \frac{4}{3} \frac{k_0' e^{k_0}}{c_0^2} I_t \int_{-\varepsilon}^t \partial_x^2 (I_\tau) (c^2 \circ \phi) \, d\tau. \end{aligned}$$

It is easy to use (3.7) and (6.11) to obtain that $|I_t'|, |\partial_x^2 I_{t,\tau}| \lesssim 1$. Using (6.6), we have that

$$\begin{aligned} \partial_x (I_t) \partial_x (c^2 \circ \phi) &= \frac{8}{3} \left(\int_{-\varepsilon}^t \partial_x (a \circ \phi) \right) I_t \partial_x (c^2 \circ \phi) \\ &= c_0^2 \left[8\partial_t \left(\int_{-\varepsilon}^t \partial_x (a \circ \phi) \right) I_t^{-1} c^{-1} \circ \phi - 8\partial_x (a \circ \phi) I_t^{-1} c^{-1} \circ \phi + 2\partial_x (I_t^{-1}) (c^{-1} \partial_\theta z) \circ \phi \right]. \end{aligned} \quad (7.7)$$

Using (6.6) and (7.7), we have that

$$\partial_x \left(\frac{k_0' e^{k_0}}{c_0^2} I_t \right) \int_{-\varepsilon}^t I_\tau \partial_x (c^2 \circ \phi) \, d\tau = -3 \frac{\partial_x (k_0' e^{k_0} I_t)}{c_0} + 6 \frac{c_0' k_0' e^{k_0}}{c_0^2} I_t$$

$$\begin{aligned}
& + \left[3\partial_x(k'_0 e^{k_0}) + 8k'_0 e^{k_0} \int_{-\varepsilon}^t \partial_x(a \circ \phi) d\tau - 6\frac{c'_0 k'_0 e^{k_0}}{c_0} \right] c^{-1} \circ \phi \\
& - 2\partial_x(k'_0 e^{k_0} I_t) \int_{-\varepsilon}^t I_\tau^{-1}(c^{-1} \partial_{\theta z}) \circ \phi d\tau \\
& + 4\frac{c'_0 k'_0 e^{k_0}}{c_0} I_t \int_{-\varepsilon}^t I_\tau^{-1}(c^{-1} \partial_{\theta z}) \circ \phi d\tau. \\
\frac{k'_0 e^{k_0}}{c_0^2} I_t \int_{-\varepsilon}^t \partial_x(I_\tau) \partial_x(c^2 \circ \phi) d\tau & = 8k'_0 e^{k_0} \left(\int_{-\varepsilon}^t \partial_x(a \circ \phi) \right) c^{-1} \circ \phi + 2k'_0 e^{k_0} I_t \int_{-\varepsilon}^t \partial_x(I_\tau^{-1})(c^{-1} \partial_{\theta z}) \circ \phi d\tau \\
& - 8c_0^2 k'_0 e^{k_0} I_t \int_{-\varepsilon}^t I_\tau^{-3} (\partial_{\theta} a c^{-3}) \circ \phi d\tau.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
\partial_x^2(\varpi \circ \phi) & = \partial_x^2(\varpi_0 I_t) - 8\frac{\partial_x(k'_0 e^{k_0} I_t)}{c_0} + 12\frac{c'_0 k'_0 e^{k_0}}{c_0^2} I_t \\
& + \left[8\partial_x(k'_0 e^{k_0}) + \frac{64}{3}k'_0 e^{k_0} \int_{-\varepsilon}^t \partial_x(a \circ \phi) d\tau - 8\frac{c'_0 k'_0 e^{k_0}}{c_0} \right] c^{-1} \circ \phi - 4k'_0 e^{k_0} \phi_x(c^{-2} \partial_{\theta} c) \circ \phi \\
& - \frac{16}{3}\partial_x(k'_0 e^{k_0} I_t) \int_{-\varepsilon}^t I_\tau^{-1}(c^{-1} \partial_{\theta z}) \circ \phi d\tau + \frac{16}{3}\frac{c'_0 k'_0 e^{k_0}}{c_0} I_t \int_{-\varepsilon}^t I_\tau^{-1}(c^{-1} \partial_{\theta z}) \circ \phi d\tau \\
& + \frac{4}{3}\partial_x^2\left(\frac{k'_0 e^{k_0}}{c_0^2} I_t\right) \int_{-\varepsilon}^t I_\tau(c^2 \circ \phi) d\tau + \frac{8}{3}\partial_x\left(\frac{k'_0 e^{k_0}}{c_0^2} I_t\right) \int_{-\varepsilon}^t \partial_x(I_\tau)(c^2 \circ \phi) d\tau \\
& + \frac{4}{3}\frac{k'_0 e^{k_0}}{c_0^2} I_t \int_{-\varepsilon}^t \partial_x^2(I_\tau)(c^2 \circ \phi) d\tau \\
& + \frac{2}{3}c_0^2 k'_0 e^{k_0} I_t \int_{-\varepsilon}^t I_\tau^{-3} ([c^{-4} \partial_{\theta z} + c^{-3} \partial_{\theta} k] \partial_{\theta} w) \circ \phi d\tau \\
& - \frac{2}{3}c_0^2 k'_0 e^{k_0} I_t \int_{-\varepsilon}^t I_\tau^{-3} (c^{-4} \partial_{\theta} z^2 + c^{-3} \partial_{\theta} k \partial_{\theta} z) \circ \phi d\tau \\
& + c_0^2 k'_0 e^{k_0} I_t \int_{-\varepsilon}^t I_\tau^{-3} \left(\frac{2}{3}c^{-2} \mathcal{E} - \frac{8}{3}c^{-3} f - \frac{32}{3} \partial_{\theta} a c^{-3} \right) \circ \phi d\tau. \tag{7.8}
\end{aligned}$$

Rearranging this and using $k \circ \phi = k_0$, (4.1), (4.6), and (3.8), we have

$$\partial_{\theta}^2 \varpi = A c^{-1} \partial_{\theta} c + B \tag{7.9}$$

where

$$\begin{aligned}
A & := 2\partial_{\theta} \varpi - 4c^{-1} e^k \partial_{\theta} k, \tag{7.10} \\
B \circ \phi & := -(Q_2 c) \circ \phi \\
& + [c_0^{-4} I_t^4 \partial_x^2(\varpi_0 I_t) - 8c_0^{-5} I_t^4 \partial_x(k'_0 e^{k_0} I_t) + 12c_0^{-6} c'_0 k'_0 e^{k_0} I_t^5] c^4 \circ \phi \\
& + \left[8\partial_x(k'_0 e^{k_0}) + \frac{64}{3}k'_0 e^{k_0} \int_{-\varepsilon}^t \partial_x(a \circ \phi) d\tau - 8\frac{c'_0 k'_0 e^{k_0}}{c_0} \right] c_0^{-4} I_t^4 c^3 \circ \phi \\
& - \frac{16}{3}c_0^{-4} I_t^4 \partial_x(k'_0 e^{k_0} I_t) \int_{-\varepsilon}^t I_\tau^{-1}(c^{-1} \partial_{\theta z}) \circ \phi d\tau c^4 \circ \phi + \frac{16}{3}c_0^{-5} c'_0 k'_0 e^{k_0} I_t^5 \int_{-\varepsilon}^t I_\tau^{-1}(c^{-1} \partial_{\theta z}) \circ \phi d\tau c^4 \circ \phi
\end{aligned}$$

$$\begin{aligned}
 &+ \frac{4}{3}c_0^{-4}I_t^4\partial_x^2\left(\frac{k_0'e^{k_0}}{c_0^2}I_t\right)\int_{-\varepsilon}^t I_\tau(c^2\circ\phi)\,d\tau c^4\circ\phi + \frac{8}{3}c_0^{-4}I_t^4\partial_x\left(\frac{k_0'e^{k_0}}{c_0^2}I_t\right)\int_{-\varepsilon}^t \partial_x(I_\tau)(c^2\circ\phi)\,d\tau c^4\circ\phi \\
 &+ \frac{4}{3}c_0^{-6}k_0'e^{k_0}I_t^5\int_{-\varepsilon}^t \partial_x^2(I_\tau)(c^2\circ\phi)\,d\tau c^4\circ\phi \\
 &+ \frac{2}{3}c_0^{-2}k_0'e^{k_0}I_t^5\left[\int_{-\varepsilon}^t I_\tau^{-3}([c^{-4}\partial_\theta z + c^{-3}\partial_\theta k]\partial_\theta w)\circ\phi\,d\tau \right. \\
 &\quad \left. - \int_{-\varepsilon}^t I_\tau^{-3}(c^{-4}\partial_\theta z^2 + c^{-3}\partial_\theta k\partial_\theta z)\circ\phi\,d\tau\right]c^4\circ\phi \\
 &+ c_0^{-2}k_0'e^{k_0}I_t^5\int_{-\varepsilon}^t I_\tau^{-3}\left(\frac{2}{3}c^{-2}\mathcal{E} - \frac{8}{3}c^{-3}f - \frac{32}{3}\partial_\theta a c^{-3}\right)\circ\phi\,d\tau c^4\circ\phi.
 \end{aligned} \tag{7.11}$$

Taking two derivatives of (4.7), we find that

$$|\varpi_0''| \lesssim \varepsilon^{-2} + |w_0''|. \tag{7.12}$$

It follows that

$$|B| \lesssim \varepsilon^{-2} + |w_0''\circ\phi^{-1}|. \tag{7.13}$$

We therefore conclude that

$$|\partial_\theta^2\varpi| \lesssim \varepsilon^{-2} + |w_0''\circ\phi^{-1}| + \varepsilon^{-1}|\partial_\theta w|. \tag{7.14}$$

Differentiating (4.3) and using (4.1) allows us to compute that

$$-3\partial_t(\partial_\theta\varpi\circ\phi) = (A\partial_\theta w + C)\circ\phi \tag{7.15}$$

where

$$\begin{aligned}
 C &:= 4e^k c^{-1}\partial_\theta k\partial_\theta z + 2\partial_\theta\varpi\partial_\theta z \\
 &\quad - 8a\partial_\theta\varpi + 8\partial_\theta a\varpi + 4e^k\mathcal{E} + 4e^k\partial_\theta k^2.
 \end{aligned} \tag{7.16}$$

This, along with (7.14), implies that $\partial_\theta\varpi$ is transversal. Therefore (see (6.10)), $\partial_\theta a$ is 2-transversal.

Now taking ∂_θ or (6.10) and using (6.4), (6.18), and (7.14) gives us

$$\partial_\theta^3 a = 2[\partial_\theta a - c - 2z]c^{-1}\partial_\theta^2 c + \mathcal{O}(\varepsilon^{-2} + |w_0''\circ\phi^{-1}| + \varepsilon^{-1}|\partial_\theta w| + |\partial_\theta w|^2). \tag{7.17}$$

Now, one can compute that

$$\begin{aligned}
 \partial_x^2(a\circ\phi) &= \phi_x^2(-[1 + 2c^{-1}z]\partial_\theta w + [3 + 2c^{-1}z]\partial_\theta z)\circ\phi \\
 &\quad + \phi_x^2\left(\frac{1}{4}c^2\partial_\theta k\varpi e^{-k} - \frac{1}{4}c^2\partial_\theta\varpi e^{-k} + \partial_\theta a Q_2\right)\circ\phi.
 \end{aligned} \tag{7.18}$$

It now follows from (4.11), (5.7), (6.4), (6.18), and (7.14) that

$$\int_{-\varepsilon}^t \partial_x^3(a\circ\phi)\,d\tau = \phi_x^3([3 + 6c^{-1}z]c^{-1}\partial_\theta c)\circ\phi + \mathcal{O}(\varepsilon^{-1} + \varepsilon|w_0''|). \tag{7.19}$$

Therefore,

$$\Phi_{xx} = 2\frac{c_0^2 c_0''' - 3c_0 c_0' c_0'' + 2(c_0')^3}{c_0^3} - \phi_x^3([16 + 32c^{-1}z]c^{-1}\partial_\theta c)\circ\phi + \mathcal{O}(\varepsilon^{-1} + \varepsilon|w_0''|). \tag{7.20}$$

These equations will be used in § 8.1, § 8.2, § 9.1, and § 9.2.‡‡

‡‡For the fifth-order estimates, one actually has to write out the full formula for (7.19) and (7.20) and work with it. We will omit such straightforward but space-consuming details.

7.3. $\partial_\theta^3 z$ bounds

We know that

$$\psi_x(\partial_\theta f \circ \psi) = \left(\frac{1}{2} \psi_x \partial_\theta k \circ \psi + \psi_x(c^{-1} \partial_\theta c) \circ \psi - 3 \int_{-\varepsilon}^t \psi_x \partial_\theta^2 z \circ \psi \, d\tau \right) f \circ \psi + J_t c \circ \psi \partial_x \left(J_t^{-1}(c^{-1} f) \circ \psi \right).$$

Recall from our Duhamel formula for f that

$$\begin{aligned} & J_t^{-1}(c^{-1} f) \circ \psi \\ &= c_0^{-1}(z_0'' - \frac{1}{2} c_0^{-1} c_0' z_0' + \frac{1}{2} c_0 k_0'') e^{-\frac{1}{2} k_0} \\ &+ \int_{-\varepsilon}^t J_t^{-1} \left(\left[\frac{1}{12} \partial_\theta k^2 - \frac{1}{4} c^{-2} \partial_\theta z^2 - \frac{1}{24} \mathcal{E} - 2 \partial_\theta a c^{-2} z + \frac{8}{3} c^{-1} z + \frac{16}{3} c^{-2} z^2 \right] \partial_\theta w \right) \circ \psi \, d\tau \\ &+ \int_{-\varepsilon}^t J_t^{-1} \left(\frac{1}{12} c \partial_\theta k \mathcal{E} - \frac{1}{12} \partial_\theta k^2 \partial_\theta z + \frac{1}{4} c^{-2} \partial_\theta z^3 + \frac{1}{8} \mathcal{E} \partial_\theta z \right) \circ \psi \, d\tau \\ &+ \int_{-\varepsilon}^t J_t^{-1} \left(2 \partial_\theta a c^{-2} \partial_\theta z z - 8 c^{-1} \partial_\theta z z - \frac{16}{3} c^{-2} \partial_\theta z z^2 - 4 \partial_\theta a c^{-1} \partial_\theta z + \frac{2}{3} c \partial_\theta \varpi e^{-k} z - \frac{2}{3} c \partial_\theta k \varpi e^{-k} z \right) \circ \psi. \end{aligned} \quad (7.21)$$

Taking ∂_x of (7.21) and using (5.4) and (7.3), we have that

$$\begin{aligned} J_t c \circ \psi \partial_x \left(J_t^{-1}(c^{-1} f) \circ \psi \right) &= \psi_x \left(-\frac{1}{8} \partial_\theta c \partial_\theta k^2 + \frac{3}{8} c^{-2} \partial_\theta c \partial_\theta z^2 + \frac{1}{16} \mathcal{E} \partial_\theta c \right) \circ \psi \\ &+ \psi_x \left(3 \partial_\theta a c^{-2} \partial_\theta c z - 4 c^{-1} \partial_\theta c z - 8 c^{-2} \partial_\theta c z^2 \right) \circ \psi \\ &+ \mathcal{O}(\varepsilon^{\beta_3 \wedge \gamma_3 \wedge \beta_2 - 1 \wedge \gamma_2 - 1 \wedge \beta_1 - 2} + \varepsilon^{\beta_1} |w_0''| + \varepsilon^{\beta_1 + \gamma_1 + 1} \|w_0''\|_{L^\infty}). \end{aligned} \quad (7.22)$$

Therefore

$$\begin{aligned} \partial_\theta f &= c^{-1} \partial_\theta c f - \frac{1}{8} \partial_\theta c \partial_\theta k^2 + \frac{3}{8} c^{-2} \partial_\theta c \partial_\theta z^2 + \frac{1}{16} \mathcal{E} \partial_\theta c \\ &+ 3 \partial_\theta a c^{-2} \partial_\theta c z - 4 c^{-1} \partial_\theta c z - 8 c^{-2} \partial_\theta c z^2 \\ &+ \mathcal{O}(\varepsilon^{\beta_3 \wedge \gamma_3 \wedge \beta_2 - 1 \wedge \gamma_2 - 1 \wedge \beta_1 - 2} + \varepsilon^{\beta_1} |w_0'' \circ \psi^{-1}| + \varepsilon^{\beta_1 + \gamma_1 + 1} \|w_0''\|_{L^\infty}). \end{aligned} \quad (7.23)$$

Taking ∂_θ of (6.20) and using these bounds, we conclude

$$\begin{aligned} \partial_\theta^3 z &= \left[\frac{1}{2} \partial_\theta z - \frac{1}{2} c \partial_\theta k \right] c^{-1} \partial_\theta^2 c \\ &+ \mathcal{O}(\varepsilon^{\beta_3 \wedge \gamma_3 \wedge \beta_2 - 1 \wedge \gamma_2 - 1 \wedge \beta_1 - 2} + \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1} |\partial_\theta w| + \varepsilon^{\beta_1} |w_0'' \circ \psi^{-1}| + \varepsilon^{\beta_1} |\partial_\theta w|^2). \end{aligned} \quad (7.24)$$

Since (6.14) can be rewritten as

$$\begin{aligned} -\frac{3}{2} \partial_t(f \circ \psi) &= \left(\left[f - \frac{1}{8} c \partial_\theta k^2 + \frac{3}{8} c^{-1} \partial_\theta z^2 + \frac{1}{16} c \mathcal{E} + 3 \partial_\theta a c^{-1} z - 4 z - 8 c^{-1} z^2 \right] \partial_\theta w \right) \circ \psi \\ &+ \left(\left(4a + \frac{1}{2} c \partial_\theta k + \frac{9}{2} \partial_\theta z \right) f + \frac{1}{8} c \partial_\theta k^2 \partial_\theta z - \frac{1}{8} c^2 \partial_\theta k \mathcal{E} - \frac{3}{8} c^{-1} \partial_\theta z^3 \right) \circ \psi \\ &+ \left(-\frac{3}{16} c \partial_\theta z \mathcal{E} - (3 \partial_\theta a c^{-1} z - 4 z - 8 c^{-1} z^2) \partial_\theta z + 6 \partial_\theta a \partial_\theta z \right) \circ \psi \\ &+ (8 z \partial_\theta z + c^2 \partial_\theta \varpi e^{-k} z + c^2 \partial_\theta \varpi e^{-k} z) \circ \psi. \end{aligned}$$

we can also conclude that f is transversal, and therefore $\partial_\theta z$ is 2-transversal.

8. Fourth derivative estimates

8.1. $\partial_\theta^4 k$ bounds

Taking ∂_x of $\tilde{\mathcal{E}} \circ \phi$ and using (4.1) and (7.20) gives us

$$\begin{aligned} \partial_x(\tilde{\mathcal{E}} \circ \phi) &= \phi_x^{-3}(\partial_x^4 k_0 - 6\Phi k_0''' - (4\partial_x \Phi - 11\Phi^2)k_0'' - (\partial_x^2 \Phi - 7\partial_x \Phi \Phi + 6\Phi^3)k_0') + 6\phi_x(\tilde{\mathcal{E}}c^{-1}\partial_\theta c) \circ \phi \\ &= \phi_x^{-3}(\partial_x^4 k_0 - 6\Phi k_0''' - (4\partial_x \Phi - 11\Phi^2)k_0'' - (2\frac{c_0^2 c_0'' - 3c_0 c_0' c_0'' + 2(c_0')^3}{c_0^3} - 7\partial_x \Phi \Phi + 6\Phi^3)k_0') \\ &\quad + \phi_x([\partial_\theta k[16 + 32c^{-1}z] + 6\tilde{\mathcal{E}}]c^{-1}\partial_\theta c) \circ \phi + \mathcal{O}(\varepsilon^{\gamma_1-1} + \varepsilon^{\gamma_1+1}|w_0''|). \\ \Rightarrow \partial_\theta \tilde{\mathcal{E}} &= [\phi_x^{-4}(\partial_x^4 k_0 - 6\Phi k_0''' - (4\partial_x \Phi - 11\Phi^2)k_0'' - (2\frac{c_0^2 c_0'' - 3c_0 c_0' c_0'' + 2(c_0')^3}{c_0^3} - 7\partial_x \Phi \Phi + 6\Phi^3)k_0') \circ \phi^{-1} \\ &\quad + [\partial_\theta k[16 + 32c^{-1}z] + 6\tilde{\mathcal{E}}]c^{-1}\partial_\theta c + \mathcal{O}(\varepsilon^{\gamma_1-1} + \varepsilon^{\gamma_1+1}|w_0''|)]. \end{aligned}$$

Define

$$\hat{\mathcal{E}} = \partial_\theta \tilde{\mathcal{E}} - [\partial_\theta k[16 + 32c^{-1}z] + 6\tilde{\mathcal{E}}]c^{-1}\partial_\theta c. \quad (8.1)$$

Then (4.11) and (6.12) tell us that

$$|\hat{\mathcal{E}}| \lesssim \varepsilon^{\gamma_4 \wedge \gamma_3 - 1 \wedge \gamma_2 - 2 \wedge \gamma_1 - 3} + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} |w_0'' \circ \phi^{-1}| + \varepsilon^{\gamma_1} |w_0''' \circ \phi^{-1}|.$$

Using (7.18), we compute that

$$\begin{aligned} -3\partial_t(\tilde{\mathcal{E}} \circ \phi) &= (6\tilde{\mathcal{E}}\partial_\theta w + 6\tilde{\mathcal{E}}\partial_\theta z) \circ \phi - 16\phi_x^{-3}k_0''\partial_x(a \circ \phi) \\ &\quad + 16\phi_x^{-3}k_0'(4\Phi\partial_x(a \circ \phi) - \partial_x^2(a \circ \phi)) \\ &= ([\partial_\theta k[16 + 32c^{-1}z] + 6\tilde{\mathcal{E}}]\partial_\theta w + 6\tilde{\mathcal{E}}\partial_\theta z) \circ \phi + \mathcal{O}(\varepsilon^{\gamma_2 \wedge \gamma_1 - 1}). \end{aligned} \quad (8.2)$$

so $\tilde{\mathcal{E}}$ is transversal, and therefore $\partial_\theta k$ is 3-transversal. This will be used in § 9.

Taking ∂_θ of (7.5) gives us

$$\begin{aligned} \partial_\theta^4 k &= \hat{\mathcal{E}} + [16 + 32c^{-1}z]c^{-1}\partial_\theta c\partial_\theta k + 12\tilde{\mathcal{E}}c^{-1}\partial_\theta c \\ &\quad + 20\mathcal{E}c^{-2}\partial_\theta c^2 + 8\mathcal{E}c^{-1}\partial_\theta^2 c + 6c^{-2}\partial_\theta c\partial_\theta^2 c\partial_\theta k + 2c^{-1}\partial_\theta^3 c\partial_\theta k. \end{aligned} \quad (8.3)$$

Note that the terms of order $|\partial_\theta w|^3$ happen to cancel when this computation is done.

It follows from (6.3), (6.18), (7.3), and (7.24) that

$$\begin{aligned} |\partial_\theta^4 k| &\lesssim \varepsilon^{\gamma_4 \wedge \gamma_3 - 1 \wedge \gamma_2 - 1 \wedge \gamma_1 - 3} + \varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \gamma_1 - 2} |\partial_\theta w| \\ &\quad + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} (|w_0'' \circ \phi^{-1}| + |\partial_\theta w|^2 + |\partial_\theta^2 w|) \\ &\quad + \varepsilon^{\gamma_1} (|w_0''' \circ \phi^{-1}| + |w_0'' \circ \phi^{-1}| |\partial_\theta w| + |\partial_\theta w| |\partial_\theta^2 w| + |\partial_\theta^3 w|). \end{aligned} \quad (8.4)$$

8.2. $\partial_\theta^4 a$ bounds

At this point, we can apply Lemma 5.5 to conclude that the variable Q_2 defined in (3.8) is transversal. In fact, Lemma 5.5 allows us to conclude that Q_2 is 2-transversal, but we will not need to use that until § 9.

Recall from § 7.2 that

$$\begin{cases} \partial_\theta^2 \varpi = Ac^{-1} \partial_\theta c + B \\ -3\partial_t(\partial_\theta \varpi \circ \phi) = (A\partial_\theta w + C) \circ \phi \end{cases}$$

where A, B, C are defined by (7.10), (7.11), and (7.16). Since $c, \partial_\theta k, \partial_\theta a, \partial_\theta z, \partial_\theta \varpi, \mathcal{E}$ are all transversal, it is immediate that A and C are transversal. We now also know that f and Q_2 are transversal, so Lemma 5.5 lets us conclude that B is transversal. So $\partial_\theta \varpi$ is 2-transversal. This fact will be utilized in § 8.3.

It is immediately clear from (7.10) that

$$\partial_\theta^3 \varpi = \partial_\theta Ac^{-1} \partial_\theta c - Ac^{-2} \partial_\theta c^2 + Ac^{-1} \partial_\theta^2 c + \partial_\theta B.$$

Since

$$\partial_\theta A = [2A - 4c^{-1} e^k \partial_\theta k] c^{-1} \partial_\theta c + 2B - 4c^{-1} e^k \partial_\theta k^2 - 4c^{-1} e^k \mathcal{E},$$

we conclude that

$$\partial_\theta^3 \varpi = Ac^{-1} \partial_\theta^2 c + [A - 4c^{-1} e^k \partial_\theta k] c^{-2} \partial_\theta c^2 + [2B - 4c^{-1} e^k \partial_\theta k^2 - 4c^{-1} e^k \mathcal{E}] c^{-1} \partial_\theta c + \partial_\theta B \quad (8.5)$$

So, to estimate $\partial_\theta^3 \varpi$, all that remains is to bound $\partial_\theta B$.

We know from (7.19) that

$$\partial_x^3 \mathcal{I}_t = \phi_x^3 ([3 + 6c^{-1} z] c^{-1} \partial_\theta c) \circ \phi + \mathcal{O}(\varepsilon^{-1} + \varepsilon |w_0''|). \quad (8.6)$$

We know from (5.6) that

$$\partial_x \left(\int_{-\varepsilon}^t \mathcal{I}_\tau^{-3} ([c^{-4} \partial_\theta z + c^{-3} \partial_\theta k] \partial_\theta w) \circ \phi \, d\tau \right) = -3\phi_x ([c^{-4} \partial_\theta z + c^{-3} \partial_\theta k] c^{-1} \partial_\theta c) \circ \phi + \mathcal{O}(\varepsilon^{\beta_2+1 \wedge \gamma_2+1 \wedge \beta_1}).$$

It is straightforward to compute that

$$\partial_\theta Q_2 = 2Q_2 c^{-1} \partial_\theta c + \mathcal{O}(\varepsilon^{-2} + |w_0'' \circ \phi^{-1}|).$$

Taking ∂_x^3 of (4.7) produces

$$|\varpi_0'''| \lesssim \varepsilon^{-3} + \varepsilon^{-1} |w_0''| + |w_0'''|. \quad (8.7)$$

Therefore, taking ∂_x of (7.11) and using (7.3) and (7.23), we conclude that

$$|\partial_\theta B| \lesssim \varepsilon^{-3} + \varepsilon^{-1} |w_0'' \circ \phi^{-1}| + |w_0''' \circ \phi^{-1}| + (\varepsilon^{-2} + |w_0'' \circ \phi^{-1}|) |\partial_\theta w|. \quad (8.8)$$

Therefore,

$$\begin{aligned} |\partial_\theta^3 \varpi| &\lesssim \varepsilon^{-3} + \varepsilon^{-1} |w_0'' \circ \phi^{-1}| + |w_0''' \circ \phi^{-1}| \\ &\quad + (\varepsilon^{-2} + |w_0'' \circ \phi^{-1}|) |\partial_\theta w| + \varepsilon^{-1} |\partial_\theta w|^2 + \varepsilon^{-1} |\partial_\theta^2 w|. \end{aligned} \quad (8.9)$$

Now, taking ∂_θ^3 of (4.9) and using (6.4), (6.9), (7.6), (7.14), (7.24), and (8.9) shows that

$$\begin{aligned} |\partial_\theta^4 a| &\lesssim \varepsilon^{-3} + \varepsilon^{-1} |w_0'' \circ \phi^{-1}| + |w_0''' \circ \phi^{-1}| \\ &\quad + (\varepsilon^{-2} + |w_0'' \circ \phi^{-1}|) |\partial_\theta w| + \varepsilon^{-1} |\partial_\theta w|^2 + \varepsilon^{-1} |\partial_\theta^2 w| \\ &\quad + |\partial_\theta w| |\partial_\theta^2 w| + |\partial_\theta^3 w|. \end{aligned} \quad (8.10)$$

8.3. $\partial_\theta^4 z$ bounds

Abusing notation, introduce a function J defined so that

$$J \circ \psi(x, t) = J_t(x).$$

Lemma 5.5 implies that J is 2-transversal. Using the new function J , we can rewrite (7.21) as

$$(J^{-1}c^{-1}f) \circ \psi = c_0^{-1}(z_0'' - \frac{1}{2}c_0^{-1}c_0'z_0' + \frac{1}{2}c_0k_0'')e^{-\frac{1}{2}k_0} + \int_{-\varepsilon}^t (Jh_1 \partial_\theta w) \circ \psi \, d\tau + \int_{-\varepsilon}^t (Jh_2) \circ \psi \, d\tau$$

Given everything that has been proven up to this point, h_1 and h_2 are both 2-transversal. It follows from Lemma 5.5 that $J^{-1}c^{-1}f$ is 2-transversal, and since c and J are both 2-transversal, it thus follows that f is 2-transversal. This will be utilized in § 9.

Using Lemma 5.4 on the functions h_1 and h_2 , along with estimates from the previous sections, we conclude that

$$\left| \partial_x^2 \left(\int_{-\varepsilon}^t (Jh_1 \partial_\theta w) \circ \psi \, d\tau + \int_{-\varepsilon}^t (Jh_2) \circ \psi \, d\tau \right) \right| \lesssim \varepsilon^{\mu-4} + \varepsilon^{\mu-3} |\partial_\theta w \circ \psi| + \varepsilon^{\mu-2} |\partial_\theta w \circ \psi|^2 + \varepsilon^{\mu-2} |\partial_\theta^2 w \circ \psi|.$$

It therefore follows that

$$\left| \partial_x^2 \left((J^{-1}c^{-1}f) \circ \psi \right) \right| \lesssim \varepsilon^{\mu-5} + \varepsilon^{\mu-3} |\partial_\theta w \circ \psi| + \varepsilon^{\mu-2} |\partial_\theta w \circ \psi|^2 + \varepsilon^{\mu-2} |\partial_\theta^2 w \circ \psi|.$$

We conclude that

$$|\partial_\theta^2 f| \lesssim \varepsilon^{\mu-5} + \varepsilon^{\mu-\frac{7}{2}} |\partial_\theta w| + \varepsilon^{\mu-2} |\partial_\theta w|^2 + \varepsilon^{\mu-2} |\partial_\theta^2 w|.$$

It now follows that

$$\begin{aligned} |\partial_\theta^4 z| &\lesssim \varepsilon^{\mu-5} + \varepsilon^{\mu-\frac{7}{2}} |\partial_\theta w| + \varepsilon^{\mu-2} |\partial_\theta w|^2 + \varepsilon^{\mu-2} |\partial_\theta^2 w| \\ &\quad + \varepsilon^{\beta_1} (|\partial_\theta w|^3 + |\partial_\theta w| |\partial_\theta^2 w| + |\partial_\theta^3 w|). \end{aligned} \quad (8.11)$$

9. Fifth Order Estimates

9.1. $\partial_\theta^5 k$ bounds

We already know (see § 8.1) that $\partial_\theta k$ is 3-transversal, and we will not need to show that $\partial_\theta k$ is 4-transversal. $\partial_\theta k$ is 4-transversal, but it doesn't matter for our purposes. One can easily establish the bound

$$\begin{aligned} |\partial_\theta \hat{\mathcal{E}}| &\lesssim \varepsilon^{\gamma_5 \wedge \gamma_4 - 1 \wedge \gamma_3 - 2 \wedge \gamma_2 - 3 \wedge \gamma_1 - 4} + \varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \gamma_1 - 2} |w_0'' \circ \phi^{-1}| + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} |w_0''' \circ \phi^{-1}| \\ &\quad + \varepsilon^{\gamma_1} (|w_0'' \circ \phi^{-1}|^2 + |\partial_x^4 w_0 \circ \phi^{-1}|) \\ &\quad + (\varepsilon^{\gamma_4 \wedge \gamma_3 - 1 \wedge \gamma_2 - 2 \wedge \gamma_1 - 3} + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} |w_0'' \circ \phi^{-1}| + \varepsilon^{\gamma_1} |w_0''' \circ \phi^{-1}|) |\partial_\theta w| \end{aligned}$$

It now follows from taking ∂_θ of (8.3) that

$$\begin{aligned} \partial_\theta^5 k &= 2c^{-1} \partial_\theta^4 c \partial_\theta k + \mathcal{O}(\varepsilon^{\gamma_5 \wedge \gamma_4 - 1 \wedge \gamma_3 - \frac{5}{2} \wedge \gamma_2 - 4 \wedge \gamma_1 - \frac{11}{2}} + \varepsilon^{\gamma_4 \wedge \gamma_3 - 1 \wedge \gamma_2 - \frac{5}{2} \wedge \gamma_1 - 4} |\partial_\theta w| \\ &\quad + \varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \gamma_1 - \frac{5}{2}} (|\partial_\theta w|^2 + |\partial_\theta^2 w|) \\ &\quad + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} (|\partial_\theta w|^3 + |\partial_\theta w| |\partial_\theta^2 w| + |\partial_\theta^3 w|) \\ &\quad + \varepsilon^{\gamma_1} (|\partial_\theta w|^2 |\partial_\theta^2 w| + |\partial_\theta^2 w|^2 + |\partial_\theta w| |\partial_\theta^3 w|)). \end{aligned} \quad (9.1)$$

9.2. $\partial_\theta^5 a$ bounds

Since $c, \partial_\theta k, \partial_\theta a, \partial_\theta z, \partial_\theta \varpi$, and \mathcal{E} are all 2-transversal, it follows immediately that A and C are 2-transversal. f and Q_2 are also 2-transversal, so it follows from Lemma 5.5 that B is 2-transversal. Therefore, $\partial_\theta \varpi$ is 3-transversal. This will be used in § 9.3.

Taking ∂_θ of (8.5) and using the bounds we already have on $A, B, C, \partial_\theta A$, and $\partial_\theta B$ gives us

$$|\partial_\theta^4 \varpi| \lesssim |\partial_\theta^2 B| + \varepsilon^{-4} |\partial_\theta w| + \varepsilon^{-5/2} (|\partial_\theta w|^2 + |\partial_\theta^2 w|) + \varepsilon^{-1} (|\partial_\theta w|^3 + |\partial_\theta w| |\partial_\theta^2 w| + |\partial_\theta^3 w|).$$

$\partial_\theta^2 B$ can be bounded in a manner similar to the way $\partial_\theta B$ was bounded. One simply needs to use a lemma similar to Lemma 5.4 but for the 2-characteristics, which is very straightforward to prove at this point. Then, since $\partial_\theta^2 Q$ and $\partial_x^4 I_t$ can be explicitly computed and bounded ^{§§}, one can bound $\partial_\theta^2 B$ and conclude that

$$|\partial_\theta^4 \varpi| \lesssim \varepsilon^{-11/2} + \varepsilon^{-4} |\partial_\theta w| + \varepsilon^{-5/2} (|\partial_\theta w|^2 + |\partial_\theta^2 w|) + \varepsilon^{-1} (|\partial_\theta w|^3 + |\partial_\theta w| |\partial_\theta^2 w| + |\partial_\theta^3 w|).$$

From here, taking ∂_θ^3 of (6.10) gives

$$\begin{aligned} \partial_\theta^5 a &= 2[\partial_\theta a - c - 2z]c^{-1} \partial_\theta^4 c + \mathcal{O}(\varepsilon^{-11/2} + \varepsilon^{-4} |\partial_\theta w| + \varepsilon^{-5/2} (|\partial_\theta w|^2 + |\partial_\theta^2 w|) \\ &\quad + \varepsilon^{-1} (|\partial_\theta w|^3 + |\partial_\theta w| |\partial_\theta^2 w| + |\partial_\theta^3 w|) \\ &\quad + |\partial_\theta w| |\partial_\theta^3 w| + |\partial_\theta w|^2 |\partial_\theta^2 w| + |\partial_\theta^2 w|^2). \end{aligned} \quad (9.2)$$

9.3. $\partial_\theta^5 z$ bounds

One can use Lemma 5.2, Lemma 5.4, and Lemma 5.5 to derive a lemma for 3-transversal functions analogous to Lemma 5.4. Bounding $\partial_\theta^5 z$ now follows in a manner completely analogous to § 8.3. One obtains

$$\begin{aligned} \partial_\theta^5 z &= [\tfrac{1}{2} \partial_\theta z - \tfrac{1}{2} c \partial_\theta k] c^{-1} \partial_\theta^4 c + \mathcal{O}(\varepsilon^{\mu-13/2} + \varepsilon^{\mu-5} |\partial_\theta w| + \varepsilon^{\mu-7/2} (|\partial_\theta w|^2 + |\partial_\theta^2 w|) \\ &\quad + \varepsilon^{\mu-2} (|\partial_\theta w|^3 + |\partial_\theta w| |\partial_\theta^2 w| + |\partial_\theta^3 w|) \\ &\quad \varepsilon^{\beta_1} (|\partial_\theta w|^4 + |\partial_\theta w| |\partial_\theta^3 w| + |\partial_\theta w|^2 |\partial_\theta^2 w| + |\partial_\theta^2 w|^2)). \end{aligned} \quad (9.3)$$

10. Estimates along η

10.1. Second derivative estimates η

It follows from the first derivative estimates that

$$|\partial_x I_t| \lesssim \varepsilon^{\gamma_1}$$

where I_t is the integrating factor in (3.13). It follows from the second derivative estimates that

$$\begin{aligned} \eta_x |\partial_\theta^2 k \circ \eta| &\lesssim \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} \\ \eta_x |\partial_\theta^2 a \circ \eta| &\lesssim \varepsilon^{-1} \\ \eta_x |\partial_\theta^2 z \circ \eta| &\lesssim \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1} \end{aligned}$$

^{§§}One must write out the full equation for (7.19) in order to do this, which is arduous but straightforward.

Taking ∂_x of (3.14) and using these bounds, we find that

$$|\partial_x(\eta_x q^w \circ \eta)| \leq |w_0''|(1 + \mathcal{O}(\varepsilon)) + \mathcal{O}(\varepsilon^{\gamma_2 \wedge \gamma_1 - 1}) + \mathcal{O}(\varepsilon^{0 \wedge \beta_1 + \gamma_1})(\varepsilon + t) \sup_{[-\varepsilon, t]} \eta_{xx}$$

Taking ∂_x of (3.15) and plugging in this estimate gives us

$$\begin{aligned} \sup_{[-\varepsilon, t]} |\eta_{xx}| &\leq (\varepsilon + t)|w_0''|(1 + \mathcal{O}(\varepsilon)) + \mathcal{O}(\varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1})(\varepsilon + t) + \mathcal{O}(\varepsilon^{\beta_1})(\varepsilon + t) \sup_{[-\varepsilon, t]} |\eta_{xx}|. \\ \implies \sup_{[-\varepsilon, t]} |\eta_{xx}| &\leq \frac{(\varepsilon + t)[|w_0''|(1 + \mathcal{O}(\varepsilon)) + \mathcal{O}(\varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1})]}{1 - \mathcal{O}(\varepsilon^\mu)} \\ &\lesssim (\varepsilon + t)(|w_0''| + \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1}). \end{aligned} \quad (10.1)$$

It follows that

$$|\eta_{xx}(x, t)| \leq \mathcal{B}(\varepsilon^{-2}(\varepsilon + t); \varepsilon^{-5/2}(\varepsilon + t)). \quad (10.2)$$

Plugging (10.1) into our bound for $|\partial_x(\eta_x q^w \circ \eta)|$ gives us

$$|\partial_x(\eta_x q^w \circ \eta)| \leq |w_0''|(1 + \mathcal{O}(\varepsilon)) + \mathcal{O}(\varepsilon^{\gamma_2 \wedge \gamma_1 - 1}) \quad (10.3)$$

Using the η_{xx} bound along with our second derivative bounds, we obtain that

$$|\partial_x^2(k \circ \eta)| \lesssim \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} + \varepsilon^{\gamma_1}(\varepsilon + t)|w_0''| \quad (10.4)$$

$$|\partial_x^2(a \circ \eta)| \lesssim \varepsilon^{-1} + (\varepsilon + t)|w_0''| \quad (10.5)$$

$$|\partial_x^2(z \circ \eta)| \lesssim \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1} + \varepsilon^{\beta_1}(\varepsilon + t)|w_0''| \quad (10.6)$$

$$|\partial_x^2(\varpi \circ \eta)| \lesssim \varepsilon^{-\frac{5}{2}} + \varepsilon^{-1}(\varepsilon + t)|w_0''|. \quad (10.7)$$

Since

$$\partial_x^2 I_t = \left(\frac{1}{8} \partial_x^2(k \circ \eta) - \frac{8}{3} \int_{-\varepsilon}^t \partial_x^2(a \circ \eta) d\tau \right) I_t + \left(\frac{1}{8} \partial_x(k \circ \eta) - \frac{8}{3} \int_{-\varepsilon}^t \partial_x(a \circ \eta) d\tau \right)^2 I_t,$$

it now follows that

$$|\partial_x^2 I_t| \lesssim \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} + \varepsilon^{\gamma_1}(\varepsilon + t)|w_0''|.$$

Last since

$$\eta_x q^w \circ \eta = \partial_x(w \circ \eta) - \frac{1}{4} c \circ \eta \partial_x(k \circ \eta),$$

we know that

$$\partial_x^2(w \circ \eta) = \partial_x(\eta_x q^w \circ \eta) + \frac{1}{4} \partial_x(c \circ \eta) \partial_x(k \circ \eta) + \frac{1}{4} c \circ \eta \partial_x^2(k \circ \eta),$$

and therefore (10.1) and (10.4) imply that

$$|\partial_x^2(w \circ \eta)| \lesssim |w_0''| + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1}. \quad (10.8)$$

Since

$$\eta_x^2 \partial_\theta^2 w \circ \eta = \partial_x^2(w \circ \eta) - \eta_{xx} \partial_\theta w \circ \eta,$$

it follows that

$$\eta_x^2 |\partial_\theta^2 w \circ \eta| \lesssim |w_0''| + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} + \varepsilon^{-1} \frac{|\eta_{xx}|}{\eta_x}. \quad (10.9)$$

$$\implies \eta_x^3 |\partial_\theta^2 w \circ \eta| \lesssim |w_0''| + \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1}. \quad (10.10)$$

10.2. Third derivative estimates along η

Using the third derivative estimates and (10.9) gives us

$$\begin{aligned}\eta_x^2 |\partial_\theta^3 k \circ \eta| &\lesssim \varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \gamma_1 - 2} + \varepsilon^{\gamma_1} |w_0''| + \varepsilon^{\gamma_1} |w_0'' \circ \phi^{-1} \circ \eta| + \varepsilon^{\gamma_1 - 1} \frac{|\eta_{xxx}|}{\eta_x} \\ \eta_x^2 |\partial_\theta^3 a \circ \eta| &\lesssim \varepsilon^{-2} + |w_0''| + |w_0'' \circ \phi^{-1} \circ \eta| + \varepsilon^{-1} \frac{|\eta_{xxx}|}{\eta_x}. \\ \eta_x^2 |\partial_\theta^3 z \circ \eta| &\lesssim \varepsilon^{\beta_3 \wedge \gamma_3 \wedge \beta_2 - 1 \wedge \gamma_2 - 1 \wedge \beta_1 - 2} + \varepsilon^{\beta_1} |w_0''| + \varepsilon^{\beta_1} |w_0'' \circ \psi^{-1} \circ \eta| + \varepsilon^{\beta_1 - 1} \frac{|\eta_{xxx}|}{\eta_x} \\ \eta_x^2 |\partial_\theta^3 \varpi \circ \eta| &\lesssim \varepsilon^{-3} + \varepsilon^{-1} |w_0''| + \varepsilon^{-1} |w_0'' \circ \phi^{-1} \circ \eta| + |w_0''' \circ \phi^{-1} \circ \eta| + \varepsilon^{-2} \frac{|\eta_{xxx}|}{\eta_x}.\end{aligned}$$

These estimates will be useful in § 10.4 and § 10.5.

Multiplying the above bounds by η_x gives us

$$\begin{aligned}\eta_x^3 |\partial_\theta^3 k \circ \eta| &\lesssim \varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \gamma_1 - 2} + \varepsilon^{\gamma_1} |w_0''| + \varepsilon^{\gamma_1} |w_0'' \circ \phi^{-1} \circ \eta| \\ \eta_x^3 |\partial_\theta^3 a \circ \eta| &\lesssim \varepsilon^{-2} + |w_0''| + |w_0'' \circ \phi^{-1} \circ \eta| \\ \eta_x^3 |\partial_\theta^3 z \circ \eta| &\lesssim \varepsilon^{\beta_3 \wedge \gamma_3 \wedge \beta_2 - 1 \wedge \gamma_2 - 1 \wedge \beta_1 - 2} + \varepsilon^{\beta_1} |w_0''| + \varepsilon^{\beta_1} |w_0'' \circ \psi^{-1} \circ \eta| \\ \eta_x^3 |\partial_\theta^3 \varpi \circ \eta| &\lesssim \varepsilon^{-3} + \varepsilon^{-1} |w_0''| + \varepsilon^{-1} |w_0'' \circ \phi^{-1} \circ \eta| + |w_0''' \circ \phi^{-1} \circ \eta|.\end{aligned}$$

Using this, we compute that

$$\begin{aligned}\left| \partial_x^2 (\eta_x q^w \circ \eta) - w_0''' e^{-\frac{1}{8}k_0} I_t \right| &\lesssim \varepsilon^{\gamma_1} |w_0''| + \varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \mu - 2} + \varepsilon^{1 \wedge \beta_1 + \gamma_1 + 1} (|w_0'' \circ \psi^{-1} \circ \eta| + |w_0'' \circ \phi^{-1} \circ \eta|) \\ &\quad + \sup_{[-\varepsilon, t]} |\eta_{xxx}| \varepsilon^{0 \wedge \beta_2 + \gamma_1 + 1 \wedge \gamma_2 + \gamma_1 + 1 \wedge \beta_1 + \gamma_1} + \varepsilon^{0 \wedge \beta_1 + \gamma_1} (\varepsilon + t) \sup_{[-\varepsilon, t]} |\eta_{xxx}| \\ &\lesssim \varepsilon^{\gamma_1} |w_0''| + \varepsilon^{-2 \wedge \gamma_3 \wedge \beta_2 + \gamma_1 - 1 \wedge \gamma_2 - 1 \wedge \beta_1 + \gamma_1 - 2} + \varepsilon^{0 \wedge \beta_1 + \gamma_1} (\varepsilon + t) \sup_{[-\varepsilon, t]} |\eta_{xxx}|.\end{aligned}$$

This is true for all $x \in \mathbb{T}, t \in [-\varepsilon, T_*]$.

Taking ∂_x^2 of (3.15) and using this bound tells us that

$$\begin{aligned}\sup_{[-\varepsilon, t]} |\eta_{xxx}| &\lesssim (\varepsilon + t) (|w_0'''| + \varepsilon^{\mu-4} + \varepsilon^{\beta_1} \sup_{[-\varepsilon, t]} |\eta_{xxx}|) \\ \implies |\eta_{xxx}| &\lesssim (\varepsilon + t) (|w_0'''| + \varepsilon^{\mu-4})\end{aligned}$$

everywhere. Using this bound, we conclude that

$$\left| \eta_{xxx} - w_0''' \int_{-\varepsilon}^t e^{-\frac{1}{8}k_0} I_\tau d\tau \right| \lesssim (\varepsilon + t)^2 \varepsilon^{\beta_1} |w_0'''| + (\varepsilon + t) \varepsilon^{\mu-4}.$$

Since $w_0''' \sim \varepsilon^{-4}$ for $|x| \leq \varepsilon^{3/2}$ and $\|w_0'''\|_{L^\infty} \lesssim \varepsilon^{-4}$, this bound lets us conclude that

$$\eta_{xxx} \sim (\varepsilon + t) \varepsilon^{-4} \quad \forall |x| \leq \varepsilon^{3/2},$$

and

$$|\eta_{xxx}| \lesssim \varepsilon^{-3} \quad \forall (x, t) \in \mathbb{T} \times [-\varepsilon, T_*].$$

We now conclude that

$$\left| \partial_x^2 (\eta_x q^w \circ \eta) \right| \lesssim |w_0'''| + \varepsilon^\mu |w_0''| + \varepsilon^{-2 \wedge \gamma_3 \wedge \beta_2 + \gamma_1 - 1 \wedge \gamma_2 - 1 \wedge \beta_1 + \gamma_1 - 2}$$

We know that for all $(x, t) \in \mathbb{T} \times [-\varepsilon, T_*)$ we have

$$\begin{aligned} |\partial_x^3(k \circ \eta)| &\lesssim (\varepsilon^{\gamma_3 \wedge \gamma_2 - 1 \wedge \gamma_1 - 2} + \varepsilon^{\gamma_1} |w_0''| + \varepsilon^{\gamma_1} |w_0'' \circ \phi^{-1} \circ \eta|) \eta_x + \varepsilon^{\gamma_2 \wedge \gamma_1 - 1} |\eta_{xx}| + \varepsilon^{\gamma_1} |\eta_{xxx}|. \\ |\partial_x^3(a \circ \eta)| &\lesssim (\varepsilon^{-2} + |w_0''| + |w_0'' \circ \phi^{-1} \circ \eta|) \eta_x + \varepsilon^{-1} |\eta_{xx}| + |\eta_{xxx}|. \\ |\partial_x^3(z \circ \eta)| &\lesssim (\varepsilon^{\beta_3 \wedge \gamma_3 \wedge \beta_2 - 1 \wedge \gamma_2 - 1 \wedge \beta_1 - 2} + \varepsilon^{\beta_1} |w_0''| + \varepsilon^{\beta_1} |w_0'' \circ \psi^{-1} \circ \eta|) \eta_x + \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1} |\eta_{xx}| + \varepsilon^{\beta_1} |\eta_{xxx}| \\ |\partial_x^3(\varpi \circ \eta)| &\lesssim (\varepsilon^{-3} + \varepsilon^{-1} |w_0''| + \varepsilon^{-1} |w_0'' \circ \phi^{-1} \circ \eta| + |w_0'' \circ \phi^{-1} \circ \eta|) \eta_x + \varepsilon^{-2} |\eta_{xx}| + \varepsilon^{-1} |\eta_{xxx}|. \end{aligned}$$

Therefore, we have the bounds

$$\begin{aligned} |\partial_x^3(k \circ \eta)| &\lesssim \varepsilon^{\mu-3} \\ |\partial_x^3(a \circ \eta)| &\lesssim \varepsilon^{-3} \\ |\partial_x^3(z \circ \eta)| &\lesssim \varepsilon^{\mu-4} \\ |\partial_x^3(\varpi \circ \eta)| &\lesssim \varepsilon^{-4}. \end{aligned}$$

Since

$$\partial_x^2(\eta_x q^w \circ \eta) = \partial_x^3(w \circ \eta) - \frac{1}{4} \partial_x^2(c \circ \eta) \partial_x(k \circ \eta) - \frac{1}{2} \partial_x(c \circ \eta) \partial_x^2(k \circ \eta) - \frac{1}{4} c \circ \eta \partial_x^3(k \circ \eta).$$

It follows that

$$|\partial_x^3(w \circ \eta)| \lesssim \varepsilon^{-4}. \quad (10.11)$$

Last it is easy to use the bounds on $\partial_x^3(k \circ \eta)$ and $\partial_x^3(a \circ \eta)$ to conclude that

$$|\partial_x^3 I_t| \lesssim \varepsilon^{\mu-3}.$$

10.3. Blowup time, location, and sharp bounds for η_x and η_{xx}

Lemma 10.1 (Existence and uniqueness of blowup label). *There exists a unique label $x_* \in \mathbb{T}$ such that $\eta_x(x_*, T_*) = 0$. Furthermore, we have $|x_*| \lesssim \varepsilon^{\mu+2}$ and*

$$\eta_x(x_*, T_*) = \eta_{xx}(x_*, T_*) = 0.$$

Proof of Lemma 10.1. Due to (4.16), we know that η_x is bound below outside of $(x, t) \in [-\varepsilon^{3/2}, \varepsilon^{3/2}] \times [-\varepsilon, T_*]$. We know that $\eta_{xxx} > 0$ in $[-\varepsilon^{3/2}, \varepsilon^{3/2}] \times (-\varepsilon, T_*]$, so for all $t \in (-\varepsilon, T_*]$ there is at most one zero of $\eta_{xx}(\cdot, t)$ in $(-\varepsilon^{3/2}, \varepsilon^{3/2}]$.

We know from § 10.1 that for all $(x, t) \in \mathbb{T} \times [-\varepsilon, T_*]$ we have

$$|\eta_{xx}(x, t) - w_0''(x) \int_{-\varepsilon}^t e^{-\frac{1}{8}k_0(x)} I_\tau(x) d\tau| \lesssim (\varepsilon + t)(\varepsilon^{\beta_1+1} |w_0''(x)| + \varepsilon^{\beta_2 \wedge \gamma_2 \wedge \beta_1 - 1}).$$

Recall that $|w_0''(x)| \lesssim \varepsilon^{-2}$ for $|x| \leq \varepsilon^2$. It follows that for $|x| \leq \varepsilon^2$ we have

$$|\eta_{xx} - w_0'' \int_{-\varepsilon}^t e^{-\frac{1}{8}k_0} I_\tau d\tau| \lesssim (\varepsilon + t) \varepsilon^{\mu-2}.$$

Since $w_0''(0) = 0$ and $w_0''' \sim \varepsilon^{-4}$ for $|x| \leq \varepsilon^{3/2}$, it follows that

$$|w_0''(x)| \gtrsim \varepsilon^{-4}|x| \text{ and } \operatorname{sgn}(w_0''(x)) = \operatorname{sgn}(x)$$

for $|x| \leq \varepsilon^{3/2}$. Therefore, we have

$$|\eta_{xx}| \gtrsim (\varepsilon + t)\varepsilon^{-4}[|x| - \mathcal{O}(\varepsilon^{\mu+2})] \quad \forall |x| \leq \varepsilon^2.$$

It follows that there exists a constant C such that for $C\varepsilon^{2+\mu} < x < \varepsilon^2$ we have $\eta_{xx}(x, t) > 0$ and for $-\varepsilon^2 < x < -C\varepsilon^{2+\mu}$ we have $\eta_{xx}(x, t) < 0$. So for all $t \in (-\varepsilon, T_*]$, there exists a unique zero of $\eta_{xx}(\cdot, t)$ in $(-\varepsilon^{3/2}, \varepsilon^{3/2})$.

Therefore, we conclude that there exists a C^2 curve $x_* : (-\varepsilon, T_*] \rightarrow \mathbb{R}$ such that

$$\{(x, t) : |x| \leq \varepsilon^{3/2}, \eta_{xx}(x, t) = 0, -\varepsilon < t \leq T_*\} = \{(x_*(t), t) : -\varepsilon < t \leq T_*\}.$$

Furthermore, we know that $|x_*(t)| \leq C\varepsilon^{2+\mu}$ for all t . From here it is easy to conclude that $\eta_{xx}(x, t) < 0$ for $-\varepsilon^{3/2} \leq x < x_*(t)$ and $\eta_{xx}(x, t) > 0$ for $x_*(t) < x \leq \varepsilon^{3/2}$, so that $x_*(t)$ must be the minimizer of $\eta_x(\cdot, t)$ over $[-\varepsilon^{3/2}, \varepsilon^{3/2}]$.

Define $x_* := x_*(T_*)$. We know that $\min_{\mathbb{T}} \eta_x(\cdot, t) \rightarrow 0$ as $t \rightarrow T_*$ and η_x is bound below for $|x| \geq \varepsilon^{3/2}$, so $\eta_x(x_*(t), t) \rightarrow 0$ as $t \rightarrow T_*$. Our result now follows. \square

We can now improve upon our lower bounds for η_x . Let $x_*(t)$ be the curve from the proof of Lemma 10.1. If $t > -\varepsilon$ and $x \in (-\pi, \pi]$, there exists $\bar{x}(x, t)$ in between x and x_* such that

$$\begin{aligned} \eta_x(x, t) &= \eta_x(x_*(t), t) + \frac{\eta_{xxx}(\bar{x}(x, t), t)}{2}(x - x_*(t))^2 \\ &\geq \frac{\eta_{xxx}(\bar{x}(x, t), t)}{2}(x - x_*(t))^2. \end{aligned}$$

Since $|x_*| \lesssim \varepsilon^{2+\mu}$, if $\varepsilon^2 \leq |x| \leq \varepsilon^{3/2}$, then $(x - x_*(t))^2 \gtrsim \varepsilon^4$ and $\eta_{xxx}(\bar{x}, t) \gtrsim (\varepsilon + t)\varepsilon^{-4}$, so we have

$$\eta_x(x, t) \gtrsim (\varepsilon + t).$$

It follows that for $\varepsilon^2 \leq |x| \leq \varepsilon^{3/2}$, $-\frac{\varepsilon}{2} \leq t \leq T_*$ we have $\eta_x \gtrsim \varepsilon$. We already know (see § 4.4) that

$$\eta_x \geq -\frac{t}{\varepsilon} + \mathcal{O}(\varepsilon^\mu)$$

for all $(x, t) \in \mathbb{T} \times [-\varepsilon, T_*]$, so we conclude that

$$\eta_x(x, t) \gtrsim \varepsilon \quad \text{for } \varepsilon^2 \leq |x| \leq \varepsilon^{3/2}. \quad (10.12)$$

Lemma 10.2 (Improved estimates for η_x and η_{xx}). *There exist constants A, c, C such that for all $(x, t) \in [-\varepsilon^2, \varepsilon^2] \times [-\varepsilon, T_*]$, we have*

$$\frac{1}{2\varepsilon}(T_* - t) + c(\varepsilon + t)\varepsilon^{-4}(x - x_*)^2 \leq \eta_x(x, t) \leq \frac{3}{2\varepsilon}(T_* - t) + C\varepsilon^{-3}(x - x_*)^2 \quad (10.13)$$

$$-A\varepsilon^{-2}(T_* - t) + c(\varepsilon + t)\varepsilon^{-4}(x - x_*) \leq \eta_{xx}(x, t) \leq A\varepsilon^{-2}(T_* - t) + C\varepsilon^{-3}(x - x_*) \text{ if } x \geq x_* \quad (10.14)$$

$$-A\varepsilon^{-2}(T_* - t) + C\varepsilon^{-3}(x - x_*) \leq \eta_{xx}(x, t) \leq A\varepsilon^{-2}(T_* - t) + c(\varepsilon + t)\varepsilon^{-4}(x - x_*) \text{ if } x \leq x_* \quad (10.15)$$

Proof of Lemma 10.2. Fix a point $(x, t) \in [-\varepsilon^2, \varepsilon^2] \times [-\varepsilon, T^*]$. We know that η_x is C^1 on $\mathbb{T} \times [-\varepsilon, T^*]$ and is C^2 on $\mathbb{T} \times [-\varepsilon, T^*]$. Therefore, Taylor's theorem tells us that there exists a point (x_1, t_1) on the segment connecting (x_*, T_*) to (x, t) such that

$$\begin{aligned} \eta_x(x, t) &= \eta_{xt}(x_*, T_*)(t - T_*) + \frac{1}{2}\eta_{xxx}(x_1, t_1)(x - x_*)^2 \\ &\quad + \eta_{xxt}(x_1, t_1)(t - T_*)(x - x_*) + \frac{1}{2}\eta_{xtt}(x_1, t_1)(t - T_*)^2. \end{aligned} \quad (10.16)$$

Similarly, there exists a point (x_2, t_2) on the segment such that

$$\eta_{xx}(x, t) = \eta_{xxx}(x_2, t_2)(x - x_*) + \eta_{txx}(x_2, t_2)(t - T_*). \quad (10.17)$$

We know that

$$\eta_{xt} = w'_0 e^{-\frac{1}{8}k_0} I_t + \mathcal{O}(\varepsilon^{\beta_1}).$$

We also know that since $w''_0(0) = 0$, $|x_*| \lesssim \varepsilon^{2+\mu}$, and $|w'''_0| \lesssim \varepsilon^{-4}$ we have

$$-\frac{1}{\varepsilon} \leq w'_0(x_*) \leq -\frac{1+C\varepsilon^{1+2\mu}}{\varepsilon}.$$

$$-\frac{1}{\varepsilon} - \mathcal{O}(\varepsilon^{\gamma_1}) \leq \eta_{xt}(x_*, T_*) \leq -\frac{1+C\varepsilon^{1+2\mu}}{\varepsilon} + \mathcal{O}(\varepsilon^{\gamma_1}).$$

We also know that for $i = 1, 2$

$$\eta_{xxx}(x_i, t_i) \sim (\varepsilon + t_i)\varepsilon^{-4}.$$

$$\eta_{txx} = \partial_x(\eta_x q^w \circ \eta) + \frac{1}{4}\partial_x(c \circ \eta)\partial_x(k \circ \eta) + \frac{1}{4}c \circ \eta \partial_x^2(k \circ \eta) + \frac{1}{3}\partial_x^2(z \circ \eta).$$

So for $i = 1, 2$

$$|\eta_{txx}(x_i, t_i)| \lesssim \varepsilon^{-2}.$$

Also

$$\begin{aligned} \eta_{xtt} &= (w'_0 - \frac{1}{4}c_0 k'_0)\partial_t(I_t e^{-\frac{1}{8}k_0}) + (\frac{1}{12}c\partial_\theta q^z - \frac{8}{3}\partial_\theta a w) \circ \eta \\ &\quad + \eta_{xt}(\frac{1}{4}\partial_\theta k + \frac{1}{3}\partial_\theta z) \circ \eta + \eta_x(\frac{1}{4}\partial_t(\partial_\theta k \circ \eta) + \frac{1}{3}\partial_t(\partial_\theta z \circ \eta)). \\ \implies |\eta_{xtt}(x_1, t_1)| &\lesssim \varepsilon^{\gamma_2 \wedge \beta_1 - 1}. \end{aligned}$$

Our result now follows. □

Using Lemma 10.2, we can now conclude that

$$\frac{1}{\eta_x} \leq \mathcal{B}([\frac{1}{2\varepsilon}(T_* - t) + c(\varepsilon + t)\varepsilon^{-4}(x - x_*)^2]^{-1}; \varepsilon^{-1}), \quad (10.18)$$

$$\frac{\eta_{xx}^2}{\eta_x} \leq \mathcal{B}(\varepsilon^{-3}; \varepsilon^{-4}). \quad (10.19)$$

The bound (10.18) will let us deduce (2.19), and (10.19) will be used frequently in § 10.4 and § 10.5.

10.4. Fourth derivative estimates along η

We know that

$$\eta_x^4 \partial_\theta^3 w \circ \eta = \eta_x \partial_x^3 (w \circ \eta) - 3\eta_{xx} \partial_x^2 (w \circ \eta) + (3\frac{\eta_{xx}^2}{\eta_x} - \eta_{xxx}) \partial_x (w \circ \eta). \quad (10.20)$$

Therefore,

$$\eta_x^4 |\partial_\theta^3 w \circ \eta| \lesssim \varepsilon^{-4} + \varepsilon^{-1} \frac{\eta_{xx}^2}{\eta_x} \leq \mathcal{B}(\varepsilon^{-4}; \varepsilon^{-5}).$$

It now follows that

$$\eta_x^4 |\partial_\theta^4 k \circ \eta| \leq \mathcal{B}(\varepsilon^{\gamma_2 - 5/2 \wedge \mu - 4}; \varepsilon^{\mu - 5}), \quad (10.21)$$

$$\eta_x^4 |\partial_\theta^4 a \circ \eta| \leq \mathcal{B}(\varepsilon^{-4}; \varepsilon^{-5}), \quad (10.22)$$

$$\eta_x^4 |\partial_\theta^4 z \circ \eta| \leq \mathcal{B}(\varepsilon^{\mu - 5}; \varepsilon^{\mu - 6}) \quad (10.23)$$

$$\eta_x^4 |\partial_\theta^4 \varpi \circ \eta| \leq \mathcal{B}(\varepsilon^{-5}; \varepsilon^{-6}). \quad (10.24)$$

The usual argument for bounding derivatives of η_x and $\eta_x q^w \circ \eta$ now gives

$$|\partial_x^4 \eta| \leq \mathcal{B}((\varepsilon + t)(|\partial_x^4 w_0| + \varepsilon^{\mu - 5}); (\varepsilon + t)(|\partial_x^4 w_0| + \varepsilon^{\mu - 6}))$$

and

$$|\partial_x^3 (\eta_x q^w \circ \eta)| \leq \mathcal{B}(\varepsilon^{\mu - 5}; \varepsilon^{-\frac{11}{2}}).$$

In the end, we obtain that

$$|\partial_x^4 (k \circ \eta)| \leq \mathcal{B}(\varepsilon^{\gamma_2 - 3 \wedge \mu - 4}; \varepsilon^{\gamma_2 - 4 \wedge \mu - 5}), \quad (10.25)$$

$$|\partial_x^4 (a \circ \eta)| \leq \mathcal{B}(\varepsilon^{-4}; \varepsilon^{-5}), \quad (10.26)$$

$$|\partial_x^4 (z \circ \eta)| \leq \mathcal{B}(\varepsilon^{\mu - 5}; \varepsilon^{\mu - 6}), \quad (10.27)$$

$$|\partial_x^4 (\varpi \circ \eta)| \leq \mathcal{B}(\varepsilon^{-5}; \varepsilon^{-6}), \quad (10.28)$$

$$|\partial_x^4 (w \circ \eta)| \leq \mathcal{B}(\varepsilon^{\mu - 5}; \varepsilon^{-\frac{11}{2}}). \quad (10.29)$$

10.5. Fifth Derivative Estimates

These estimates are different from the previous sections because they require more algebra and hinge on admittedly unexpected cancellation. First, note that

$$\eta_x^2 \partial_\theta^2 c \circ \eta = -\partial_x (c \circ \eta) \frac{\eta_{xx}}{\eta_x} + \partial_x^2 (c \circ \eta)$$

$$\eta_x^3 \partial_\theta^3 c \circ \eta = (3\frac{\eta_{xx}^2}{\eta_x} - \eta_{xxx}) \partial_x (c \circ \eta) \frac{1}{\eta_x} - 3\partial_x^2 (c \circ \eta) \frac{\eta_{xx}}{\eta_x} + \partial_x^3 (c \circ \eta)$$

$$\eta_{xx} \eta_x^3 (\partial_\theta c \partial_\theta^2 c) \circ \eta = \partial_x (c \circ \eta) [\eta_{xx} \partial_x^2 (c \circ \eta) - \partial_x (c \circ \eta) \frac{\eta_{xx}^2}{\eta_x}]$$

$$\eta_{xx} \eta_x^3 \partial_\theta^3 c \circ \eta = (3\frac{\eta_{xx}^2}{\eta_x} - \eta_{xxx}) \partial_x (c \circ \eta) \frac{\eta_{xx}}{\eta_x} - 3\partial_x^2 (c \circ \eta) \frac{\eta_{xx}^2}{\eta_x} + \partial_x^3 (c \circ \eta) \eta_{xx}$$

$$\eta_x^5 \partial_\theta^2 c^2 \circ \eta = \partial_x (c \circ \eta)^2 \frac{\eta_{xx}}{\eta_x} - 2\partial_x^2 (c \circ \eta) \partial_x (c \circ \eta) \eta_{xx} + \partial_x^2 (c \circ \eta)^2 \eta_x$$

$$\eta_x^5 (\partial_\theta c^2 \partial_\theta^2 c) \circ \eta = \partial_x (c \circ \eta)^2 [-\partial_x (c \circ \eta) \eta_{xx} + \partial_x^2 (c \circ \eta) \eta_x]$$

$$\eta_x^5 (\partial_\theta c \partial_\theta^3 c) \circ \eta = \partial_x (c \circ \eta) [(3\frac{\eta_{xx}^2}{\eta_x} - \eta_{xxx}) \partial_x (c \circ \eta) - 3\partial_x^2 (c \circ \eta) \eta_{xx} + \partial_x^3 (c \circ \eta) \eta_x]$$

$$\begin{aligned} \eta_x^5 \partial_\theta^4 c \circ \eta &= (10\eta_{xxx} - 15 \frac{\eta_{xx}^2}{\eta_x}) \partial_x(c \circ \eta) \frac{\eta_{xx}}{\eta_x} - \partial_x^4 \eta \partial_x(c \circ \eta) + (15 \frac{\eta_{xx}^2}{\eta_x} - 4\eta_{xxx}) \partial_x^2(c \circ \eta) \\ &\quad - 6\eta_{xx} \partial_x^3(c \circ \eta) + \eta_x \partial_x^4(c \circ \eta). \end{aligned}$$

Next, note that

$$\begin{aligned} \partial_\theta^2 a &= 2[\partial_\theta a - c + 2z]c^{-1} \partial_\theta c + \mathcal{O}(\varepsilon^{-1}), \\ \partial_\theta^3 a &= 2[\partial_\theta a - c + 2z]c^{-1} \partial_\theta^2 c + \mathcal{O}(\varepsilon^{-\frac{5}{2}} + \varepsilon^{-1}|\partial_\theta w| + |\partial_\theta w|^2), \\ \partial_\theta^4 a &= 2[\partial_\theta a - c + 2z]c^{-1} \partial_\theta^3 c + \mathcal{O}(\varepsilon^{-4} + \varepsilon^{-\frac{5}{2}}|\partial_\theta w| + \varepsilon^{-1}|\partial_\theta w|^2 + \varepsilon^{-1}|\partial_\theta^2 w| + |\partial_\theta w||\partial_\theta^2 w|), \\ \partial_\theta^5 a &= 2[\partial_\theta a - c - 2z]c^{-1} \partial_\theta^4 c + \mathcal{O}(\varepsilon^{-\frac{11}{2}} + \varepsilon^{-4}|\partial_\theta w| + \varepsilon^{-5/2}(|\partial_\theta w|^2 + |\partial_\theta^2 w|) \\ &\quad + \varepsilon^{-1}(|\partial_\theta w|^3 + |\partial_\theta w||\partial_\theta^2 w| + |\partial_\theta^3 w|) \\ &\quad + |\partial_\theta w||\partial_\theta^3 w| + |\partial_\theta w|^2|\partial_\theta^2 w| + |\partial_\theta^2 w|^2). \end{aligned}$$

Combining these identities and our estimates gives us

$$\begin{aligned} 10\eta_{xxx} \frac{\eta_{xx}}{\eta_x} \eta_x \partial_\theta^2 a \circ \eta &= 2[\partial_\theta a c^{-1} - 1 - 2c^{-1}z] \circ \eta (10\eta_{xxx} + 0) \partial_x(c \circ \eta) \frac{\eta_{xx}}{\eta_x} + \mathcal{B}(\varepsilon^{-5}; \varepsilon^{-\frac{11}{2}}), \\ (10\eta_{xxx} + 15 \frac{\eta_{xx}^2}{\eta_x}) \eta_x^2 \partial_\theta^3 a \circ \eta &= 2[\partial_\theta a c^{-1} - 1 - 2c^{-1}z] \circ \eta (-10\eta_{xxx} - 15 \frac{\eta_{xx}^2}{\eta_x}) \partial_x(c \circ \eta) \frac{\eta_{xx}}{\eta_x} + \mathcal{B}(\varepsilon^{-\frac{11}{2}}; \varepsilon^{-\frac{13}{2}}), \\ 10\eta_{xx} \eta_x^3 \partial_\theta^4 a \circ \eta &= 2[\partial_\theta a c^{-1} - 1 - 2c^{-1}z] \circ \eta (-10\eta_{xxx} + 30 \frac{\eta_{xx}^2}{\eta_x}) \partial_x(c \circ \eta) \frac{\eta_{xx}}{\eta_x} + \mathcal{B}(\varepsilon^{-5}; \varepsilon^{-\frac{13}{2}}), \\ \eta_x^5 \partial_\theta^5 a \circ \eta &= 2[\partial_\theta a c^{-1} - 1 - 2c^{-1}z] \circ \eta (10\eta_{xxx} - 15 \frac{\eta_{xx}^2}{\eta_x}) \partial_x(c \circ \eta) \frac{\eta_{xx}}{\eta_x} + \mathcal{B}(\varepsilon^{-\frac{11}{2}}; \varepsilon^{-\frac{13}{2}}). \end{aligned}$$

Therefore,

$$\begin{aligned} \partial_x^5(a \circ \eta) &= \eta_x^5 \partial_\theta^5 a \circ \eta + 10\eta_{xx} \eta_x^3 \partial_\theta^4 a \circ \eta + (10\eta_{xxx} + 15 \frac{\eta_{xx}^2}{\eta_x}) \eta_x^2 \partial_\theta^3 a \circ \eta \\ &\quad + (5\partial_x^4 \eta + 10\eta_{xxx} \frac{\eta_{xx}}{\eta_x}) \eta_x \partial_\theta^2 a \circ \eta + \partial_x^5 \eta \partial_\theta a \circ \eta \\ &= 2[\partial_\theta a c^{-1} - 1 - 2c^{-1}z] \circ \eta ([10 - 10 - 10 + 10]\eta_{xxx} + [0 - 15 + 30 - 15] \frac{\eta_{xx}^2}{\eta_x}) \partial_x(c \circ \eta) \frac{\eta_{xx}}{\eta_x} \\ &\quad + \partial_x^5 \eta \partial_\theta a \circ \eta + \mathcal{B}(\varepsilon^{-\frac{11}{2}}; \varepsilon^{-\frac{13}{2}}) \\ &= \partial_x^5 \eta \partial_\theta a \circ \eta + \mathcal{B}(\varepsilon^{-\frac{11}{2}}; \varepsilon^{-\frac{13}{2}}). \end{aligned}$$

The exact same cancellation occurs for the other two variables to give us

$$\begin{aligned} \partial_x^5(k \circ \eta) &= \partial_x^5 \eta \partial_\theta k \circ \eta + \mathcal{B}(\varepsilon^{\gamma_2 - 4 \wedge \mu - \frac{11}{2}}; \varepsilon^{\mu - \frac{13}{2}}), \\ \partial_x^5(z \circ \eta) &= \partial_x^5 \eta \partial_\theta z \circ \eta + \mathcal{B}(\varepsilon^{\mu - \frac{13}{2}}; \varepsilon^{\mu - \frac{15}{2}}). \end{aligned}$$

Similar computations prove that

$$\partial_x^4(\eta_x(\partial_\theta k \partial_\theta z) \circ \eta) = \partial_x^5(k \circ \eta) \partial_\theta z \circ \eta + \partial_\theta k \circ \eta \partial_x^5(z \circ \eta) + \mathcal{B}(\varepsilon^{\gamma_2 + \mu - 5 \wedge 2\mu - \frac{13}{2}}; \varepsilon^{2\mu - \frac{15}{2}}).$$

Now the usual method for bounding the derivatives of η_x and $\eta_x q^w \circ \eta$ produces

$$|\partial_x^5 \eta| \leq \mathcal{B}((\varepsilon + t)(|\partial_x^5 w_0| + \varepsilon^{\mu - \frac{13}{2}}); (\varepsilon + t)(|\partial_x^5 w_0| + \varepsilon^{\mu - \frac{15}{2}})). \quad (10.30)$$

In the end, we obtain that

$$|\partial_x^5(k \circ \eta)| \lesssim \mathcal{B}(\varepsilon^{\mu - 6}; \varepsilon^{\mu - \frac{13}{2}}),$$

$$\begin{aligned} |\partial_x^5(a \circ \eta)| &\lesssim \mathcal{B}(\varepsilon^{-6}; \varepsilon^{-\frac{13}{2}}), \\ |\partial_x^5(z \circ \eta)| &\lesssim \mathcal{B}(\varepsilon^{\mu-7}; \varepsilon^{\mu-\frac{15}{2}}), \\ |\partial_x^5(w \circ \eta)| &\lesssim \varepsilon^{-7}. \end{aligned}$$

Using similar computations to those in this section, one can compute that

$$\eta_x^7 |\partial_\theta^5 w \circ \eta| \leq \mathcal{B}(\varepsilon^{-7}; \varepsilon^{-9}).$$

This bound, together with similar bounds that we proved for $\partial_\theta^n z \circ \eta$, $\partial_\theta^n k \circ \eta$, and $\partial_\theta^n a \circ \eta$, combines with (10.18) to establish (2.19).

11. Inversion of η

In this section, we will confine our attention to labels $x \in (-\pi, \pi]$ with $|x| \leq \varepsilon^2$.

Since $w \circ \eta(\cdot, T_*)$ is $C^{4,1}$, it has the following Taylor expansion about x_* :

$$w \circ \eta = B_0^w + B_1^w(x - x_*) + B_2^w(x - x_*)^2 + B_3^w(x - x_*)^3 + R_0^w(x)(x - x_*)^4. \tag{11.1}$$

Here

$$|B_0^w| \lesssim 1, |B_1^w| \lesssim \varepsilon^{-1}, |B_2^w| \lesssim \varepsilon^{-2}, |B_3^w| \lesssim \varepsilon^{-4}, |R_0^w| \lesssim \varepsilon^{\mu-5}. \tag{11.2}$$

The flow $\eta(\cdot, T_*)$ also has the Taylor expansion

$$\begin{aligned} \eta(x, T_*) - \xi_* &= a_3(x - x_*)^3 + a_4(x)(x - x_*)^4 \\ &= a_3(x - x_*)^3 + a_4(x_*)(x - x_*)^4 + a_5(x)(x - x_*)^5, \end{aligned} \tag{11.3}$$

where $\xi_* := \eta(x_*, T_*)$, $a_3 := \frac{1}{6}\eta_{xxx}(x_*, T_*)$, $a_4(x_*) = \frac{1}{24}\partial_x^4\eta(x_*, T_*)$,

$$a_4(x) := \frac{\int_{x_*}^x \partial_x^4 \eta(y, T_*) (x - y)^3 dy}{3!(x - x_*)^4} \quad \text{and} \quad a_5(x) := \frac{\int_{x_*}^x \partial_x^5 \eta(y, T_*) (x - y)^4 dy}{4!(x - x_*)^5}. \tag{11.4}$$

Here $a_3 \sim \varepsilon^{-3}$, $|a_4(x)| \lesssim \varepsilon^{\mu-4}$, and $|a_5(x)| \lesssim \varepsilon^{-6}$. Note that $|a_3^{-4/3} a_4| \lesssim \varepsilon^\mu$.

Let $\theta = \eta(x, T_*)$. Lemma A.3 implies that there exists a constant C such that for all $x \in [-\varepsilon^2, \varepsilon^2]$ such that $|\theta - \xi_*| \leq C\varepsilon^{-3\mu}$ we have

$$\begin{aligned} (x - x_*) &= a_3^{-1/3}(\theta - \xi_*)^{1/3} \left[1 + \frac{1}{3}(-a_3^{-4/3} a_4(\theta - \xi_*)^{1/3}) + \frac{1}{3}(-a_3^{-4/3} a_4(\theta - \xi_*)^{1/3})^2 + \mathcal{O}(\varepsilon^{3\mu}|\theta - \xi_*|) \right] \\ &= a_3^{-1/3}(\theta - \xi_*)^{1/3} \left[1 + \frac{1}{3}(-a_3^{-4/3} a_4(\theta - \xi_*)^{1/3}) + \mathcal{O}(\varepsilon^{2\mu}|\theta - \xi_*|^{2/3}) \right] \end{aligned} \tag{11.5}$$

$$= a_3^{-1/3}(\theta - \xi_*)^{1/3} [1 + \mathcal{O}(\varepsilon^\mu|\theta - \xi_*|^{1/3})]. \tag{11.6}$$

A quick bootstrap argument lets us conclude that this formula holds for all $x \in [-\varepsilon^2, \varepsilon^2]$. Furthermore, it is easy to show that there exist two constants $0 < c < C$ such that

$$\{\theta : |\theta - \xi_*| \leq c\varepsilon^3\} \subset \{\theta : |x| \leq \varepsilon^2\} \subset \{\theta : |\theta - \xi_*| \leq C\varepsilon^3\}.$$

So we are working in a neighborhood of radius $\sim \varepsilon^3$ around ξ_* .

If we define

$$\begin{aligned}\hat{a}_0^w &:= B_0^w, \\ \hat{a}_1^w &:= a_3^{-1/3} B_1^w, \\ \hat{a}_2^w &:= a_3^{-2/3} B_2^w - \frac{1}{3} a_3^{-5/3} a_4(x_*) B_1^w,\end{aligned}$$

then we have

$$|\hat{a}_0^w| \lesssim 1, \quad |\hat{a}_1^w| \lesssim 1, \quad |\hat{a}_2^w| \lesssim 1,$$

and

$$w(\theta, T_*) = \hat{a}_0^w + \hat{a}_1^w (\theta - \xi_*)^{1/3} + \hat{a}_2^w (\theta - \xi_*)^{2/3} + \mathcal{O}(\varepsilon^{-1} |\theta - \xi_*|). \quad (11.7)$$

Squaring (11.5) and cubing (11.6) gives us

$$\begin{aligned}(x - x_*)^2 &= a_3^{-2/3} (\theta - \xi_*)^{2/3} - \frac{2}{3} a_3^{-2} a_4 (\theta - \xi_*) + \mathcal{O}(\varepsilon^{2\mu+2} |\theta - \xi_*|^{4/3}), \\ (x - x_*)^3 &= a_3^{-1} (\theta - \xi_*) + \mathcal{O}(\varepsilon^{\mu+3} |\theta - \xi_*|^{4/3}).\end{aligned}$$

Therefore,

$$\begin{aligned}\eta_x(x, T_*) &= 3a_3(x - x_*)^2 + [4a_4(x) + \partial_x a_4(x)(x - x_*)](x - x_*)^3 \\ &=: 3a_3(x - x_*)^2 + \tilde{a}_4(x - x_*)^3 \\ &= 3a_3^{1/3} (\theta - \xi_*)^{2/3} + a_3^{-1} (\tilde{a}_4 - 2a_4)(\theta - \xi_*) + \mathcal{O}(\varepsilon^{2\mu-1} |\theta - \xi_*|^{4/3}).\end{aligned} \quad (11.8)$$

Using this formula, one can compute that

$$\eta_x(x, T_*)^{-1} = \frac{1}{3} a_3^{-1/3} (\theta - \xi_*)^{-2/3} - \frac{1}{9} a_3^{-5/3} (\tilde{a}_4 - 2a_4)(\theta - \xi_*)^{-1/3} + \mathcal{O}(\varepsilon^{2\mu+1}).$$

Since $a_4(x) = a_4(x_*) + \mathcal{O}(\varepsilon^{-5} |\theta - \xi_*|^{1/3})$ and $\tilde{a}_4(x) = 4a_4(x_*) + \mathcal{O}(\varepsilon^{-5} |\theta - \xi_*|^{1/3})$, it follows that

$$\eta_x(x, T_*)^{-1} = \frac{1}{3} a_3^{-1/3} (\theta - \xi_*)^{-2/3} - \frac{2}{9} a_3^{-5/3} a_4(x_*) (\theta - \xi_*)^{-1/3} + \mathcal{O}(1). \quad (11.9)$$

Since $\partial_x(w \circ \eta) = B_1^w + 2B_2^w(x - x_*) + \mathcal{O}(\varepsilon^{-4} |x - x_*|^2)$, it follows that at time T_* we have

$$\begin{aligned}\partial_\theta w \circ \eta &= \eta_x^{-1} \partial_x(w \circ \eta) \\ &= \left[\frac{1}{3} a_3^{-1/3} (\theta - \xi_*)^{-2/3} - \frac{2}{9} a_3^{-5/3} a_4(x_*) (\theta - \xi_*)^{-1/3} + \mathcal{O}(1) \right] \\ &\quad \cdot \left[B_1^w + 2a_3^{-1/3} B_2^w (\theta - \xi_*)^{1/3} + \mathcal{O}(\varepsilon^{-2} |\theta - \xi_*|^2) \right] \\ &= \frac{1}{3} \hat{a}_1^w (\theta - \xi_*)^{-2/3} + \frac{2}{3} \hat{a}_2^w (\theta - \xi_*)^{-1/3} + \mathcal{O}(\varepsilon^{-1}).\end{aligned}$$

This is the expansion for $\partial_\theta w(\cdot, T_*)$ in Theorem 2.1.

Now consider

$$\partial_\theta^2 w \circ \eta = \eta_x(x, T_*)^{-2} [\partial_x^2(w \circ \eta) - \eta_{xx}(x, T_*) \partial_\theta w \circ \eta(x, T_*)].$$

Differentiating (11.3) twice and using our above expansions for $(x - x_*)^2$ and $(x - x_*)^3$ gives us

$$\begin{aligned}\eta_{xx}(x, T_*) &= 6a_3(x - x_*) + 12a_4(x_*)(x - x_*)^2 + [20a_5 + 10\partial_x a_5(x - x_*) + \partial_x^2 a_5(x - x_*)^2](x - x_*)^3 \\ &= 6a_3(x - x_*) + 12a_4(x_*)(x - x_*)^2 + \mathcal{O}(\varepsilon^{-6}|x - x_*|^3) \\ &= 6a_3^{2/3}(\theta - \xi_*)^{1/3} + 10a_3^{-2/3}a_4(x_*)(\theta - \xi_*)^{2/3} + \mathcal{O}(\varepsilon^{-3}|\theta - \xi_*|).\end{aligned}\tag{11.10}$$

Using the fact that $\partial_x^2(w \circ \eta) = 2B_2^w + 6B_3^w(x - x_*) + \mathcal{O}(\varepsilon^{\mu-5}|x - x_*|^2)$ along with our expansion for $\partial_\theta w \circ \eta$, (11.9), and (11.10) provide the expansion for $\partial_\theta^2 w(\cdot, T_*)$ as stated in Theorem 2.1.

Last, since

$$\partial_\theta^3 w \circ \eta = \eta_x^{-3}[\partial_x^3(w \circ \eta) - 3\eta_{xx}\eta_x\partial_\theta^2 w \circ \eta - \eta_{xxx}\partial_\theta w \circ \eta],$$

we can do similar computations to determine the expansion for $\partial_\theta^3 w(\cdot, T_*)$.

To get the expansions for the variables z, k , and a , similar computations can be made, except with the constants B_j^z, B_j^k , or B_j^a instead of B_j^w . The computations for these variables are nicer because $B_1^z = B_1^k = B_1^a = B_2^z = B_2^k = B_2^a = 0$, but one should use fifth-order expansions of $z \circ \eta, k \circ \eta$ and $a \circ \eta$. So we have

$$\begin{aligned}\mathring{a}_0^z &:= B_0^z, \\ \mathring{a}_3^z &:= a_3^{-1}B_3^z, \\ \mathring{a}_4^z &:= a_3^{-4/3}B_4^z - a_3^{-7/3}a_4(x_*)B_3^z,\end{aligned}$$

and $\mathring{a}_0^k, \mathring{a}_3^k, \mathring{a}_4^k, \mathring{a}_0^a, \mathring{a}_3^a, \mathring{a}_4^a$ are defined analogously. When one does the computations, one obtains the expansions for z, k , and a listed in Theorem 2.1.

Unlike the functions $w \circ \eta, z \circ \eta, k \circ \eta$, and $a \circ \eta$, which are in $C^{4,1}(\mathbb{T})$ at time T_* , the function $\varpi \circ \eta$ has only been proven to be in $C^{3,1}(\mathbb{T})$ at time T_* , so the Taylor expansion can only go to fourth order. However, we still have $B_1^\varpi = B_2^\varpi = 0$ which allows us to get constants in our expansion. \square

A. Appendix

A.1. Basic identities

The following equations are easy to compute from (2.5):

$$-\frac{3}{2}\partial_t(c \circ \psi) = (\partial_\theta w + 4a) \circ \psi(c \circ \psi).\tag{A.1a}$$

$$-\frac{3}{2}\partial_t(\partial_\theta c \circ \psi) = (c\partial_\theta^2 w) \circ \psi + \frac{3}{2}(\partial_\theta c \partial_\theta w) + \frac{3}{2}(\partial_\theta c \partial_\theta z) \circ \psi + 4(\partial_\theta a c + a \partial_\theta c) \circ \psi.\tag{A.1b}$$

$$-\frac{3}{2}\partial_t(k \circ \psi) = (c\partial_\theta k) \circ \psi.\tag{A.1c}$$

$$-\frac{3}{2}\partial_t(\partial_\theta k \circ \psi) = (c\partial_\theta^2 k) \circ \psi + (\partial_\theta k \partial_\theta w + \partial_\theta k \partial_\theta z) \circ \psi.\tag{A.1d}$$

$$-\frac{3}{2}\partial_t(z \circ \psi) = (4az - \frac{1}{4}c^2 \partial_\theta k) \circ \psi.\tag{A.1e}$$

$$-\frac{3}{2}\partial_t(\partial_\theta z \circ \psi) = (\frac{1}{2}\partial_\theta w \partial_\theta z + \frac{3}{2}\partial_\theta z^2 - \frac{1}{2}c\partial_\theta c \partial_\theta k - \frac{1}{4}c^2 \partial_\theta^2 k) \circ \psi + 4(\partial_\theta a z + a \partial_\theta z) \circ \psi.\tag{A.1f}$$

$$-\frac{3}{2}\partial_t(a \circ \psi) = (\partial_\theta a c + 2a^2 - c^2 - 4cz - 2z^2) \circ \psi.\tag{A.1g}$$

$$-\frac{3}{2}\partial_t(\partial_\theta a \circ \psi) = (\partial_\theta^2 a c + 2\partial_\theta a \partial_\theta c + 2\partial_\theta a \partial_\theta z + 4a \partial_\theta a) \circ \psi - ([2c + 4z]\partial_\theta c + [4c + 4z]\partial_\theta z) \circ \psi.\tag{A.1h}$$

$$-\frac{3}{2}\partial_t(c \circ \phi) = (4ac + c\partial_\theta c + c\partial_\theta z) \circ \phi.\tag{A.1i}$$

$$-\frac{3}{2}\partial_i(\partial_\theta c \circ \phi) = (c\partial_\theta^2 c) \circ \phi + (c\partial_\theta^2 z + 3\partial_\theta c^2) \circ \phi + 3(\partial_\theta c \partial_\theta z) \circ \phi + 4(\partial_\theta ac + a\partial_\theta c) \circ \phi. \quad (\text{A.1j})$$

$$-\frac{3}{2}\partial_i(\partial_\theta k \circ \phi) = (\partial_\theta k \partial_\theta w + \partial_\theta k \partial_\theta z) \circ \phi. \quad (\text{A.1k})$$

$$-\frac{3}{2}\partial_i(c \circ \eta) = (\partial_\theta z + 4a) \circ \eta(c \circ \eta). \quad (\text{A.1l})$$

$$-\frac{3}{2}\partial_i(k \circ \eta) = -(c\partial_\theta k) \circ \eta \quad (\text{A.1m})$$

$$-\frac{3}{2}\partial_i(\partial_\theta k \circ \eta) = (\partial_\theta w \partial_\theta k + \partial_\theta z \partial_\theta k - c\partial_\theta^2 k) \circ \eta. \quad (\text{A.1n})$$

A.2. Quartic Inversion

If K is a field, and $K((z))$ denotes the field of formal Laurent series ^{¶¶} in the variable z . The field of Puiseux series in the variable x is then defined to be the union $\bigcup_{n>0} K((x^{1/n}))$ which is itself a field. The most important result concerning Puiseux series is the following:

Theorem A.1 (Puiseux–Newton). *If K is an algebraically closed field of characteristic 0, then the field $\bigcup_{n>0} K((x^{1/n}))$ of Puiseux series with coefficients in K an algebraically closed field. Furthermore, given a polynomial $P(y) = \sum_{i=0}^N a_i(x)y^i$ with $a_i \in \bigcup_{n>0} K((x^{1/n}))$, the coefficients of the roots of P in y can be constructed using the method of Newton polygons.*

Proof of Theorem A.1. See [27, Chapter IV, Section 3] or [28, Section 8.3]. □

Of particular interest to us will be the following special case of the Puiseux–Newton theorem:

Theorem A.2 (Analytic Puiseux–Newton). *If $\mathbb{C}\{x\}$ denotes the ring of convergent power series in x , and $f(x, y) \in \mathbb{C}\{x\}[y]$ is a polynomial of degree $m > 0$, irreducible in $\mathbb{C}\{x\}[y]$, then there exists a convergent power series $y \in \mathbb{C}\{z\}$ such that the roots of f in $\bigcup_{n>0} \mathbb{C}((x^{1/n}))$ are all given by*

$$y(x^{1/m}), y(e^{2\pi i/m} x^{1/m}), \dots, y(e^{2\pi i \frac{m-1}{m}} x^{1/m}).$$

It follows that in general if $f(x, y) \in \mathbb{C}\{x\}[y]$, then for each Puiseux series solution \bar{y} of $f(x, \bar{y}(x)) = 0$, there exists some $y \in \mathbb{C}\{z\}$ and $m \leq \deg f$ such that $\bar{y}(x) = y(x^{1/m})$.

Proof of Theorem A.2. See [28, Section 8.3]. □

Lemma A.3 (Quartic Inversion). *There exists a constant $R > 0$ and a nonempty open interval I containing 0 such that for all $a_3 \in \mathbb{R}^\times, a_4 \in \mathbb{R}$ there exists a function $y(x)$ defined for x satisfying $|a_4^3 x| < R^3 a_3^4$ such that*

$$\{(x, y) \in \mathbb{R}^2 : |a_4^3 x| < R^3 a_3^4, a_4 y \in a_3 I, -x + a_3 y^3 + a_4 y^4 = 0\} = \{(x, y(x)) : |a_4^3 x| < R^3 a_3^4\}.$$

Furthermore, $y(x)$ is an analytic function of $x^{1/3}$ satisfying the bounds

$$|y(x) - a_3^{1/3} x^{1/3} + \frac{1}{3} a_3^{-5/3} a_4 x^{2/3} - \frac{1}{3} a_3^{-3} a_4^2 x| \lesssim a_3^{-13/3} a_4^3 x^{4/3}$$

for all $|a_4^3 x| < R^3 a_3^4$, with the constant in the inequality independent of a_3, a_4 .

^{¶¶}Formal Laurent series are formal power series which allow for finitely many terms of negative degree, not to be confused with the Laurent series in complex analysis, which may have infinitely many terms of negative degree but must converge in an annulus.

Proof of Lemma A.3. The case where $a_4 = 0$ is trivial, so we will prove our result in the case $a_4 \in \mathbb{R}^\times$. Define the recursive sequence $c_0 := 1$,

$$c_n := \sum_{\substack{\vec{k} \in (\mathbb{Z}_{\geq 0})^4 \\ k_1+k_2+k_3+k_4=n-1}} c_{k_1} c_{k_2} c_{k_3} c_{k_4} - \frac{1}{3} \sum_{\substack{\vec{m} \in (\mathbb{Z}_{\geq 0})^3 \\ m_1+m_2+m_3=n \\ 0 \leq m_i \leq n-1}} c_{m_1} c_{m_2} c_{m_3},$$

and define the formal power series $\bar{y} \in \mathbb{R}[[x]]$,

$$\bar{y}(x) = \sum_{n=0}^{\infty} (-1)^n \frac{c_n}{3^n} x^{n+1}.$$

It is easy to check that $y_0(x) := \bar{y}(x^{1/3})$ is a Puiseux series solution to the algebraic equation $-x + y_0^3 + y_0^4 = 0$. It follows from A.2 that \bar{y} must be convergent with some positive (possibly infinite) radius of convergence R . Now pick any $a_3 \in \mathbb{R}^\times, a_4 \in \mathbb{R}^\times$. If we define

$$y(x) := \frac{a_3}{a_4} \bar{y}(a_3^{-4/3} a_4 x^{1/3}),$$

then it is easy to check that y solves $-x + a_3 y^3 + a_4 y^4 = 0$.

Define the interval I to be the range of \bar{y} , thought of as a function on $(-R, R)$, and define $f(x, y) = -x + a_3 y^3 + a_4 y^4$. Because $\partial_x f = -1$ everywhere, we know that for each $y \in \mathbb{R}$, the equation $f(x, y) = 0$ has exactly one solution, x . Therefore, if (x, y) is a point such that $|x| < a_3^4 a_4^{-3} R^3, y \in a_3 a_4^{-1} I$, and $f(x, y) = 0$, then there exists x' with $|x'| < a_3^4 a_4^{-3} R^3$ such that $y(x') = y$ and since $f(x', y) = f(x', y(x')) = 0$ we conclude that $x = x'$ and $y = y(x)$.

The remaining expansion follows from the fact that $c_1 = 1$ and $c_2 = 3$, combined with the fact that the power series \bar{y} is convergent. □

Theorem A.4. *There exist universal constants C_1 and C_2 such that the following is true: Suppose that $I \subset \mathbb{R}$ is an interval, $x_0 \in I$, and $\theta \in C^{3,1}(I)$ is such that $L := \|\partial_x^4 \theta\|_{L^\infty}, a_3 \in \mathbb{R}^\times$, and θ has the Taylor expansion*

$$\theta(x) = \theta_0 + a_3(x - x_0)^3 + a_4(x)(x - x_0)^4$$

at x_0 . Then for all $x \in I$ such that $|\theta(x) - \theta_0| \leq C_1 \frac{a_3^4}{L^3}$, we have

$$(x - x_0) = a_3^{-1/3} (\theta(x) - \theta_0)^{1/3} - \frac{1}{3} a_3^{-5/3} a_4(x) (\theta(x) - \theta_0)^{2/3} + \frac{1}{3} a_3^{-3} a_4(x)^2 (\theta(x) - \theta_0) + R(\theta - \theta_0),$$

where R is a $C^{0, \frac{1}{3}}$ continuous function satisfying

$$|R(\theta - \theta_0)| \leq C_2 a_3^{-13/3} a_4(x)^3 (\theta(x) - \theta_0)^{4/3}.$$

Proof of Theorem A.4. Assume without loss of generality that $a_3 > 0$. We know that $a_4 = (x - x_0)^{-4} (\theta - \theta_0 - a_3(x - x_0)^3)$ is C^3 away from x_0 and that

$$a_4(x) = \frac{\int_{x_0}^x \partial_x^4 \theta(t) (x-t)^3 dt}{3!(x-x_0)^4}$$

for all $x \neq x_0$. It follows from this formula that

$$|a_4(x)| \leq \frac{L}{4!} \quad \text{and} \quad |\partial_x a_4(x)| \leq \frac{L}{3} \frac{1}{|x-x_0|}$$

for all $x \neq x_0$.

First define the function $f : \mathbb{R} \times (I - x_0) \rightarrow \mathbb{R}$,

$$f(x, y) := -x + a_3 y^3 + a_4 (y + x_0) y^4.$$

Using our bounds on $|a_4|$ and $|\partial_x a_4|$, we see that

$$\partial_y f(x, y) \geq y^2 (3a_3 - \frac{L}{2}|y|),$$

$$f(x, \frac{a_3}{L}) \geq \frac{23}{24} \frac{a_3^4}{L^3} - |x|,$$

$$f(x, -\frac{a_3}{L}) \leq |x| - \frac{23}{24} \frac{a_3^4}{L^3}.$$

Therefore, if we define $A := \{|x| < \frac{23}{24} \frac{a_3^4}{L^3}\}$ and $B := \{|y| < \frac{6a_3}{L}\}$, then for all $x \in A$ the function $f(x, \cdot) : B \rightarrow \mathbb{R}$ is strictly increasing and has a zero in the interior of B . It follows from Corollary 1.1 in [29] that there exists a unique continuous function $h : A \rightarrow B$ such that

$$\{(x, y) \in A \times B : f(x, y) = 0\} = \{(x, h(x)) : x \in A\}.$$

Now define the function $F : \mathbb{R}^3 \rightarrow \mathbb{R}$,

$$F(x, y, a) := -x + a_3 y^3 + a y^4.$$

It is easy to check that if $|a| \leq \frac{L}{4!}$ and $|x| < \frac{23}{24} \frac{a_3^4}{L^3}$ then

$$\partial_y F(x, y, a) \geq y^2 (3a_3 - \frac{L}{3!}|y|), \quad F(x, \frac{a_3}{L}, a) > 0, \quad \text{and} \quad F(x, -\frac{a_3}{L}, a) < 0.$$

Therefore, if $\tilde{A} := \{(x, a) : |x| < \frac{23}{24} \frac{a_3^4}{L^3}, |a| \leq \frac{L}{4!}\}$ and $\tilde{B} := (-18 \frac{a_3}{L}, 18 \frac{a_3}{L})$ then for all $(x, a) \in \tilde{A}$ the function $F(x, \cdot, a) : \tilde{B} \rightarrow \mathbb{R}$ is strictly increasing and contains a 0 in the interior of \tilde{B} . It follows from Corollary 1.1 of [29] that there exists a unique $H : \tilde{A} \rightarrow \tilde{B}$ continuous such that

$$\{(x, y, a) : |x| < \frac{23}{24} \frac{a_3^4}{L^3}, |y| < 18 \frac{a_3}{L}, |a| \leq \frac{L}{4!}, F(x, y, a) = 0\} = \{(x, H(x, a), a) : |x| < \frac{23}{24} \frac{a_3^4}{L^3}, |a| \leq \frac{L}{4!}\}.$$

Our previous lemma A.3 tells us that there exist constants $R, C_2 > 0$ independent of a_3 or L such that for all $|a| \leq \frac{L}{4!}$, $|x| < R^3 (4!)^3 \frac{a_3^4}{L^3}$ we have

$$H(x, a) = a_3^{-1/3} x^{1/3} - \frac{1}{3} a_3^{-5/3} a x^{2/3} + \frac{1}{3} a_3^{-3} a^2 x + \tilde{R}(x, a),$$

where $|\tilde{R}(x, a)| \leq C_2 a_3^{-13/3} a^3 x^{4/3}$. Now suppose that $|x| < \frac{23}{24} \frac{a_3^4}{L^3}$. Then $|h(x)| < 6 \frac{a_3}{L} < 18 \frac{a_3}{L}$ and

$$F(x, h(x), a_4(h(x) + x_0)) = f(x, h(x)) = 0,$$

so $h(x) = H(x, a_4(h(x)))$. It follows that if $C_1 := \min(\frac{23}{24}, (R4!)^3)$ then we have

$$\begin{aligned} h(x) &= a_3^{-1/3} x^{1/3} - \frac{1}{3} a_3^{-5/3} a_4 (h(x) + x_0) x^{2/3} + \frac{1}{3} a_3^{-3} a_4 (h(x) + x_0) x + \tilde{R}(x, a_4(h(x) + x_0)) \\ &=: a_3^{-1/3} x^{1/3} - \frac{1}{3} a_3^{-5/3} a_4 (h(x) + x_0) x^{2/3} + \frac{1}{3} a_3^{-3} a_4 (h(x) + x_0) x + R(x) \end{aligned}$$

for all $|x| < C_1 \frac{a_3^4}{L^3}$. Our result now follows. \square

Author contributions

All authors contributed equally.

Use of AI tools declaration

The authors declare they have not used Artificial Intelligence (AI) tools in the creation of this article.

Acknowledgments

Steve Shkoller was supported in part by the NSF grant DMS-2007606, the Collaborative NSF grant DMS-2307680, and the Department of Energy Advanced Simulation and Computing (ASC) Program. Isaac Neal and Vlad Vicol were supported in part by the NSF CAREER grant DMS-1911413. Vlad Vicol was in part supported by the Collaborative NSF grant DMS-2307681, and a Simons Investigator Award.

Conflict of interest

The authors declare there is no conflict of interest.

References

1. T. Buckmaster, T. D. Drivas, S. Shkoller, V. Vicol, Simultaneous development of shocks and cusps for 2D Euler with azimuthal symmetry from smooth data, *Ann. PDE*, **8** (2022). <https://doi.org/10.1007/s40818-022-00141-6>
2. T. Buckmaster, S. Shkoller, V. Vicol, Formation of shocks for 2D isentropic compressible Euler, *Commun. Pur. Appl. Math.*, **75** (2022), 2069–2120. <https://doi.org/10.1002/cpa.21956>
3. P. D. Lax, Development of singularities of solutions of nonlinear hyperbolic partial differential equations, *J. Math. Phys.*, **5** (1964), 611–613. <https://doi.org/10.1063/1.1704154>
4. T. P. Liu, Development of singularities in the nonlinear waves for quasi-linear hyperbolic partial differential equations, *J. Differ. Equations*, **33** (1979), 92–111. [https://doi.org/10.1016/0022-0396\(79\)90082-2](https://doi.org/10.1016/0022-0396(79)90082-2)
5. T. C. Sideris, Formation of singularities in three-dimensional compressible fluids, *Commun. Math. Phys.*, **101** (1985), 475–485. <https://doi.org/10.1007/BF01210741>
6. G. Chen, R. Pan, S. Zhu, Singularity formation for the compressible Euler equations, *SIAM J. Math. Anal.*, **49** (2017), 2591–2614. <https://doi.org/10.1137/16M1062818>
7. G. Chen, Optimal density lower bound on nonisentropic gas dynamics, *J. Differ. Equations*, **268** (2020), 4017–4028. <https://doi.org/10.1016/j.jde.2019.10.017>
8. C. Dafermos, *Hyperbolic Conservation Laws in Continuum Physics*, Springer, 2005. <https://doi.org/10.1007/978-3-662-49451-6>
9. G. Q. Chen, D. Wang, The Cauchy problem for the Euler equations for compressible fluids, in *Handbook of mathematical fluid dynamics, Vol. I* (eds. S. Friedlander, and D. Serre), North-Holland, Amsterdam (2002), 421–543. [https://doi.org/10.1016/s1874-5792\(02\)80012-x](https://doi.org/10.1016/s1874-5792(02)80012-x)

10. T. P. Liu, *Shock Waves*, American Mathematical Society, Providence, RI, 2021.
11. J. Rauch, BV estimates fail for most quasilinear hyperbolic systems in dimensions greater than one, *Commun. Math. Phys.*, **106** (1986), 481–484. <https://doi.org/10.1007/BF01207258>
12. M. P. Lebaud, Description de la formation d'un choc dans le p -système, *Journal de mathématiques pures et appliquées*, **73** (1994), 523–566.
13. S. Chen, L. Dong, Formation and construction of shock for p -system, *Sci. China Ser. A-Math.*, **44** (2001), 1139–1147. <https://doi.org/10.1007/BF02877431>
14. D. X. Kong, Formation and propagation of singularities for 2×2 quasilinear hyperbolic systems, *Trans. Amer. Math. Soc.*, **354** (2002), 3155–3179. <https://doi.org/10.1090/S0002-9947-02-02982-3>
15. T. Buckmaster, T. D. Drivas, S. Shkoller, V. Vicol, Formation and development of singularities for the compressible Euler equations, in *Proceedings of the International Congress of Mathematicians 2022*, EMS Press, Berlin, **5** (2023), 3636–3659. <https://doi.org/10.4171/ICM2022/87>
16. D. Christodoulou, The formation of shocks in 3-dimensional fluids, *European Mathematical Society*, 2007. <https://doi.org/10.4171/031>
17. D. Christodoulou, The shock development problem, *European Mathematical Society*, 2019. <https://doi.org/10.4171/192>
18. H. Yin, Formation and construction of a shock wave for 3-d compressible Euler equations with the spherical initial data, *Nagoya Math. J.*, **175** (2004), 125–164. <https://doi.org/10.1017/S002776300000893X>
19. D. Christodoulou, A. Lisibach, Shock development in spherical symmetry, *Ann. PDE*, **2** (2016). <https://doi.org/10.1007/s40818-016-0009-1>
20. J. Luk, J. Speck, Shock formation in solutions to the 2d compressible Euler equations in the presence of non-zero vorticity, *Invent. Mathe.*, **214** (2018), 1–169. <https://doi.org/10.1007/s00222-018-0799-8>
21. J. Luk, J. Speck, The stability of simple plane-symmetric shock formation for 3d compressible Euler flow with vorticity and entropy, *Anal PDE*, **17** (2024), 831–941. <https://doi.org/10.2140/apde.2024.17.831>
22. T. Buckmaster, S. Shkoller, V. Vicol, Formation of point shocks for 3D compressible Euler, *Comm. Pur. Appl. Math.*, **76** (2023), 2073–2191. <https://doi.org/10.1002/cpa.22068>
23. T. Buckmaster, S. Shkoller, V. Vicol, Shock formation and vorticity creation for 3D Euler, *Comm. Pur. Appl. Math.*, **76** (2023), 1965–2072. <https://doi.org/10.1002/cpa.22067>
24. S. Shkoller, V. Vicol, The geometry of maximal development and shock formation for the Euler equations in multiple space dimensions, *Invent. Math.*, **237** (2024), 871–1252. <https://doi.org/10.1007/s00222-024-01269-x>
25. L. Abbrescia, J. Speck, The emergence of the singular boundary from the crease in 3d compressible Euler flow, preprint, arXiv:2207.07107.
26. L. D. Landau, E. M. Lifshitz, *Fluid Mechanics*, Vol. 6, Elsevier, 1987. <https://doi.org/10.1016/C2013-0-03799-1>
27. R. J. Walker, *Algebraic Curves*, Springer-Verlag, 1950.

-
28. E. Brieskorn, H. Knörrer, *Plane algebraic curves*, Springer Basel, 1986.
<https://doi.org/10.1007/978-3-0348-5097-1>
29. S. Kumagai, An implicit function theorem: comment, *J Optim Theory Appl*, **31** (1980), 285–288.
<https://doi.org/10.1007/BF00934117>



AIMS Press

©2025 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)