

GLOBAL DYNAMICS OF SOME SYSTEM OF SECOND-ORDER DIFFERENCE EQUATIONS

TRAN HONG THAI*, NGUYEN ANH DAI AND PHAM TUAN ANH

Department of Mathematics, Hung Yen University of Technology and Education
Hung Yen 160000, Vietnam

(Communicated by Moxun Tang)

ABSTRACT. In this paper, we study the boundedness and persistence of positive solution, existence of invariant rectangle, local and global behavior, and rate of convergence of positive solutions of the following systems of exponential difference equations

$$\begin{aligned}x_{n+1} &= \frac{\alpha_1 + \beta_1 e^{-x_n-1}}{\gamma_1 + y_n}, & y_{n+1} &= \frac{\alpha_2 + \beta_2 e^{-y_n-1}}{\gamma_2 + x_n}, \\x_{n+1} &= \frac{\alpha_1 + \beta_1 e^{-y_n-1}}{\gamma_1 + x_n}, & y_{n+1} &= \frac{\alpha_2 + \beta_2 e^{-x_n-1}}{\gamma_2 + y_n},\end{aligned}$$

where the parameters $\alpha_i, \beta_i, \gamma_i$ for $i \in \{1, 2\}$ and the initial conditions x_{-1}, x_0, y_{-1}, y_0 are positive real numbers. Some numerical example are given to illustrate our theoretical results.

1. Introduction. Mathematical models of population dynamics are often described by difference equations and systems of difference equations. In particular, the population models involving exponential difference equations are quite popular, although their stability analysis can be complicated. In recent years, the global asymptotic behavior of the difference equations of exponential form has been one of the main topics in the theory of difference equations (see [2, 3, 4, 5, 7, 12, 13, 14, 15, 16, 17, 8, 20] and reference cited therein).

In [4], El-Metwally et al. investigated the following population model:

$$x_{n+1} = \alpha + \beta x_{n-1} e^{-x_n}, \quad (1)$$

where the parameters α and β are positive numbers and the initial conditions x_{-1} and x_0 are arbitrary non-negative numbers. Later in [5], Fotiades et al. studied the existence, uniqueness and attractivity of prime period two solution of this equation.

Papaschinopoulos et al. [15] and Papaschinopoulos and Schinas [17] investigated the dynamical properties of two-species model described by systems of difference equations, which is natural extension of single-species population model depicted in 1.

2020 *Mathematics Subject Classification.* Primary: 39A10, 39A30; Secondary: 40A05.

Key words and phrases. System of difference equations, boundedness, persistence, asymptotic behavior, rate of convergence.

The first author is supported by UTEHY grand number UTEHY.L.2020.11.

* Corresponding author.

Ozturk et al. [12] have investigated the following difference equation:

$$y_{n+1} = \frac{\alpha + \beta e^{-y_n}}{\gamma + y_{n-1}}, \quad (2)$$

where the parameters α, β, γ are positive numbers and the initial conditions are arbitrary non-negative numbers.

Papaschinopoulos et al. [16] have studied the following systems of two difference equations of exponential form:

$$\begin{aligned} x_{n+1} &= \frac{\alpha + \beta e^{-y_n}}{\gamma + y_{n-1}}, & y_{n+1} &= \frac{\delta + \epsilon e^{-x_n}}{\eta + x_{n-1}}, \\ x_{n+1} &= \frac{\alpha + \beta e^{-y_n}}{\gamma + x_{n-1}}, & y_{n+1} &= \frac{\delta + \epsilon e^{-x_n}}{\eta + y_{n-1}}, \\ x_{n+1} &= \frac{\alpha + \beta e^{-x_n}}{\gamma + y_{n-1}}, & y_{n+1} &= \frac{\delta + \epsilon e^{-y_n}}{\eta + x_{n-1}}, \end{aligned}$$

where $\alpha, \beta, \gamma, \delta, \epsilon, \eta$ are positive constants and the initial values x_{-1}, x_0, y_{-1}, y_0 are positive numbers.

In 2016, Wang and Feng [20] have investigated the dynamics of positive solution of the following difference equation which is naturally a new form of single-species model described in 1:

$$x_{n+1} = \alpha + \beta x_n e^{-x_{n-1}},$$

where the parameters α and β are positive numbers and the initial conditions x_{-1} and x_0 are arbitrary non-negative numbers.

Motivated by the aforementioned study, our goal in this paper is to investigate the qualitative behavior of positive solutions of some systems of exponential difference equations

$$x_{n+1} = \frac{\alpha_1 + \beta_1 e^{-x_{n-1}}}{\gamma_1 + y_n}, \quad y_{n+1} = \frac{\alpha_2 + \beta_2 e^{-y_{n-1}}}{\gamma_2 + x_n}, \quad (3)$$

$$x_{n+1} = \frac{\alpha_1 + \beta_1 e^{-y_{n-1}}}{\gamma_1 + x_n}, \quad y_{n+1} = \frac{\alpha_2 + \beta_2 e^{-x_{n-1}}}{\gamma_2 + y_n}, \quad (4)$$

where the parameters $\alpha_i, \beta_i, \gamma_i$ for $i \in \{1, 2\}$ and the initial conditions x_{-1}, x_0, y_{-1}, y_0 are positive real numbers.

More precisely, we investigate the boundedness character, persistence, existence of invariant rectangle, local asymptotic stability and global behavior of unique positive equilibrium point, and rate of convergence of positive solutions of system 3 and 4 which converges to its unique positive equilibrium point. For applications and basic theory of difference equations we refer to [1, 6, 10, 11, 19].

2. Preliminaries. In this section, we present some definitions and theorems which are used throughout this study.

Let us consider fourth-dimensional discrete dynamical system of the following form:

$$x_{n+1} = f(x_n, x_{n-1}, y_n, y_{n-1}), \quad y_{n+1} = g(x_n, x_{n-1}, y_n, y_{n-1}), \quad n = 0, 1, \dots \quad (5)$$

where $f : I^2 \times J^2 \rightarrow I$ and $g : I^2 \times J^2 \rightarrow J$ are continuously differentiable functions and I, J are some intervals of real numbers. Furthermore, a solution $\{x_n, y_n\}_{n=-1}^{\infty}$ of system 5 is uniquely determined by initial conditions $(x_i, y_i) \in I \times J$ for $i \in \{-1, 0\}$.

Along with system 5, we consider the corresponding vector map $F = (f, x_n, g, y_n)$. An equilibrium point of 5 is a point (\bar{x}, \bar{y}) that satisfies

$$\bar{x} = f(\bar{x}, \bar{x}, \bar{y}, \bar{y}), \quad \bar{y} = g(\bar{x}, \bar{x}, \bar{y}, \bar{y}).$$

The point $(\bar{x}, \bar{x}, \bar{y}, \bar{y})$ is also called a fixed point of the vector map F .

Definition 2.1. Let (\bar{x}, \bar{y}) be an equilibrium point of system 5.

(i) An equilibrium point (\bar{x}, \bar{y}) is called stable if for any $\varepsilon > 0$ there is $\delta > 0$ such that for every initial conditions (x_{-1}, y_{-1}) and (x_0, y_0) , if $\|(x_{-1}, y_{-1}) - (\bar{x}, \bar{y})\| + \|(x_0, y_0) - (\bar{x}, \bar{y})\| < \delta$ implies that $\|(x_n, y_n) - (\bar{x}, \bar{y})\| < \varepsilon$ for all $n > 0$, where $\|\cdot\|$ is usual Euclidean norm in \mathbb{R}^2 .

(ii) An equilibrium point (\bar{x}, \bar{y}) is called unstable if it is not stable.

(iii) An equilibrium point (\bar{x}, \bar{y}) is called locally asymptotically stable if it stable and if, in addition, there exists $r > 0$ such that $(x_n, y_n) \rightarrow (\bar{x}, \bar{y})$ as $n \rightarrow \infty$ for all (x_{-1}, y_{-1}) and (x_0, y_0) that satisfy $\|(x_{-1}, y_{-1}) - (\bar{x}, \bar{y})\| + \|(x_0, y_0) - (\bar{x}, \bar{y})\| < r$.

(iv) An equilibrium point (\bar{x}, \bar{y}) is called global attractor if $(x_n, y_n) \rightarrow (\bar{x}, \bar{y})$ as $n \rightarrow \infty$.

(v) An equilibrium point (\bar{x}, \bar{y}) is called globally asymptotically stable if it stable and a global attractor.

Definition 2.2. Let $(\bar{x}, \bar{x}, \bar{y}, \bar{y})$ be a fixed point of a map $F = (f, x_n, g, y_n)$ where f and g are continuously differentiable functions at (\bar{x}, \bar{y}) . The linearized system of 5 about the equilibrium point (\bar{x}, \bar{y}) is

$$X_{n+1} = J_F X_n,$$

where $X_n = \begin{pmatrix} x_n \\ x_{n-1} \\ y_n \\ y_{n-1} \end{pmatrix}$ and J_F is the Jacobian matrix of system 5 about the equilibrium point (\bar{x}, \bar{y}) .

Lemma 2.3. (see [19]) Assume that $X_{n+1} = F(X_n), n = 0, 1, \dots$, is a system of difference equations such that \bar{X} is a fixed point of F . If all eigenvalues of the Jacobian matrix J_F about \bar{X} lie inside the open unit disk $|\lambda| < 1$, then \bar{X} is locally asymptotically stable. If one of them has a modulus greater than one, then \bar{X} is unstable.

Definition 2.4. A positive solution $\{x_n, y_n\}_{n=-1}^\infty$ of system 5 is bounded and persists if there exist positive constants m, M and an interger $N \geq -1$ such that

$$m \leq x_n, y_n \leq M, \quad n \geq N.$$

In order to study the asymptotic behavior of positive equilibrium, we state the following lemma which is a slight modification of Theorem 1.16 of [6] and for readers convenience we state it without its proof.

Lemma 2.5. Assume that $f : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ and $g : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ be continuous functions and a, b, c, d are positive real numbers with $a < b, c < d$. Moreover, suppose that $f : [a, b] \times [c, d] \rightarrow [a, b]$ and $g : [a, b] \times [c, d] \rightarrow [c, d]$ such that following conditions are satisfied:

- (i) $f(x, y), g(x, y)$ are decreasing with respect to x (resp. y) for all y (resp. x);
- (ii) Let m_1, M_1, m_2, M_2 are real numbers such that

$$m_1 = f(M_1, M_2), M_1 = f(m_1, m_2), m_2 = g(M_1, M_2), M_2 = g(m_1, m_2) \quad (6)$$

then $m_1 = M_1$ and $m_2 = M_2$.

Then the systems of difference equations

$$x_{n+1} = f(x_{n-1}, y_n), \quad y_{n+1} = g(x_n, y_{n-1}), \quad (7)$$

$$x_{n+1} = f(x_n, y_{n-1}), \quad y_{n+1} = g(x_{n-1}, y_n) \quad (8)$$

have a unique equilibrium point (\bar{x}, \bar{y}) and every solution (x_n, y_n) of the system 7 (resp. 8) with $x_{-1}, x_0 \in [a, b], y_{-1}, y_0 \in [c, d]$ converges to the unique equilibrium (\bar{x}, \bar{y}) .

The following results give the rate of convergence of solutions of a system of difference equations

$$X_{n+1} = [A + B(n)]X_n \quad (9)$$

where X_n is a m -dimensional vector, $A \in \mathbf{C}^{m \times m}$ is a constant matrix, and $B : \mathbb{Z}^+ \rightarrow \mathbf{C}^{m \times m}$ is a matrix function satisfying

$$\|B(n)\| \rightarrow 0 \text{ when } n \rightarrow \infty, \quad (10)$$

where $\|\cdot\|$ denotes any matrix norm which is associated with the vector norm

$$\|(x, y)\| = \sqrt{x^2 + y^2}.$$

Proposition 2.6 (Perron's theorem [18]). *Assume that condition 10 holds. If X_n is a solution of system 9, then either $X_n = 0$ for all large n or*

$$\rho = \lim_{n \rightarrow \infty} \sqrt[n]{\|X_n\|} \quad (11)$$

exists and is equal to the modulus of one of the eigenvalues of matrix A .

Proposition 2.7 (See [18]). *Assume that condition 10 holds. If X_n is a solution of system 9, then either $X_n = 0$ for all large n or*

$$\rho = \lim_{n \rightarrow \infty} \frac{\|X_{n+1}\|}{\|X_n\|} \quad (12)$$

exists and is equal to the modulus of one of the eigenvalues of matrix A .

3. Main results.

3.1. Global behavior of solutions of system 3.

3.1.1. *Boundedness and persistence.* In this section, we show the boundedness and persistence of the positive solutions of system 3.

Lemma 3.1. *Every positive solution $\{(x_n, y_n)\}$ of system 3 is bounded and persists.*

Proof. For any positive solution $\{(x_n, y_n)\}$ of system 3, one has

$$x_{n+1} \leq \frac{\alpha_1 + \beta_1}{\gamma_1} = b_1, \quad y_{n+1} \leq \frac{\alpha_2 + \beta_2}{\gamma_2} = d_1, \quad n = 0, 1, 2, \dots \quad (13)$$

Furthermore, from system 3 and 13, we obtain that

$$x_{n+1} \geq \frac{\alpha_1 + \beta_1 e^{-b_1}}{\gamma_1 + d_1} = a_1, \quad y_{n+1} \geq \frac{\alpha_2 + \beta_2 e^{-d_1}}{\gamma_2 + b_1} = c_1, \quad n = 2, 3, 4, \dots \quad (14)$$

From 13 and 14, it follows that

$$a_1 \leq x_n \leq b_1, \quad c_1 \leq y_n \leq d_1, \quad n = 3, 4, 5, \dots$$

So the proof is complete. \square

Lemma 3.2. *Let $\{(x_n, y_n)\}$ be a positive solution of system 3. Then $[a_1, b_1] \times [c_1, d_1]$ is an invariant set for system 3.*

Proof. The proof follows by induction. □

3.1.2. *Stability analysis.* In this section, we shall investigate the asymptotic behavior of system 3. Similar method can be found in [9].

Let (\bar{x}, \bar{y}) be the equilibrium point of system 3 then

$$\bar{x} = \frac{\alpha_1 + \beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}}, \quad \bar{y} = \frac{\alpha_2 + \beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}}.$$

The linearized form of system 3 about the equilibrium point (\bar{x}, \bar{y}) is given by

$$X_{n+1} = J_F(\bar{x}, \bar{y})X_n,$$

where $X_n = \begin{pmatrix} x_n \\ x_{n-1} \\ y_n \\ y_{n-1} \end{pmatrix}$ and $J_F(\bar{x}, \bar{y}) = \begin{pmatrix} 0 & -\frac{\beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}} & -\frac{\bar{x}}{\gamma_1 + \bar{y}} & 0 \\ 1 & 0 & 0 & 0 \\ -\frac{\bar{y}}{\gamma_2 + \bar{x}} & 0 & 0 & -\frac{\beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}} \\ 0 & 0 & 1 & 0 \end{pmatrix}$.

In the following theorem, we show the asymptotic behavior of the positive solutions of system 3.

Theorem 3.3. *Suppose that the following relation holds true:*

$$\beta_1 < \gamma_1, \quad \beta_2 < \gamma_2. \tag{15}$$

Then system 3 has a unique positive equilibrium (\bar{x}, \bar{y}) and every positive solution of system 3 tends to the unique positive equilibrium as $n \rightarrow \infty$.

Proof. Consider the following functions:

$$f(x, y) = \frac{\alpha_1 + \beta_1 e^{-x}}{\gamma_1 + y}, \quad g(x, y) = \frac{\alpha_2 + \beta_2 e^{-y}}{\gamma_2 + x},$$

where $x \in I_1 = [a_1, b_1], y \in I_2 = [c_1, d_1]$ which implies that $f(x, y) \in I_1, g(x, y) \in I_2$ and so that $f : I_1 \times I_2 \rightarrow I_1, g : I_1 \times I_2 \rightarrow I_2$. Then, it is easy to see that $f(x, y), g(x, y)$ are decreasing with respect to x (resp. y) for all y (resp. x). Let (m, M, r, R) be a solution of the system

$$\begin{aligned} m &= f(M, R), & M &= f(m, r), \\ r &= g(M, R), & R &= g(m, r). \end{aligned}$$

Then, one has

$$m = \frac{\alpha_1 + \beta_1 e^{-M}}{\gamma_1 + R}, \quad M = \frac{\alpha_1 + \beta_1 e^{-m}}{\gamma_1 + r}, \quad r = \frac{\alpha_2 + \beta_2 e^{-R}}{\gamma_2 + M}, \quad R = \frac{\alpha_2 + \beta_2 e^{-r}}{\gamma_2 + m}. \tag{16}$$

Moreover arguing as in the proof of Theorem 1.16 of [6], it suffices to assume that

$$m \leq M, \quad r \leq R. \tag{17}$$

From 16, we get

$$\begin{aligned} \beta_1 e^{-m} &= M(\gamma_1 + r) - \alpha_1, & \beta_1 e^{-M} &= m(\gamma_1 + R) - \alpha_1, \\ \beta_2 e^{-r} &= R(\gamma_2 + m) - \alpha_2, & \beta_2 e^{-R} &= r(\gamma_2 + M) - \alpha_2, \end{aligned}$$

which imply that

$$\begin{aligned} \beta_1(e^{-m} - e^{-M}) &= \gamma_1(M - m) + Mr - mR, \\ \beta_2(e^{-r} - e^{-R}) &= \gamma_2(R - r) + mR - Mr. \end{aligned} \tag{18}$$

Then by adding the two relations 18, we obtain

$$\beta_1(e^{-m} - e^{-M}) + \beta_2(e^{-r} - e^{-R}) = \gamma_1(M - m) + \gamma_2(R - r).$$

Moreover, we get

$$\begin{aligned} e^R - e^r &= e^\xi(R - r), \quad \min\{R, r\} \leq \xi \leq \max\{R, r\}, \\ e^M - e^m &= e^\theta(M - m), \quad \min\{M, m\} \leq \theta \leq \max\{M, m\}. \end{aligned} \tag{19}$$

Then from 19, imply that

$$\beta_1 e^{-m-M+\theta}(M - m) + \beta_2 e^{-r-R+\xi}(R - r) = \gamma_1(M - m) + \gamma_2(R - r). \tag{20}$$

Hence from 20, we have

$$\gamma_1(M - m) \left(1 - \frac{\beta_1}{\gamma_1} e^{-m-M+\theta}\right) + \gamma_2(R - r) \left(1 - \frac{\beta_2}{\gamma_2} e^{-r-R+\xi}\right) = 0. \tag{21}$$

Finally, from 15, 17 and 21, it follows that $M = m$ and $R = r$. Therefore, from Lemma 2.5, it follows that system 3 has a unique positive equilibrium (\bar{x}, \bar{y}) and every positive solution of system 3 tends to the unique positive equilibrium as $n \rightarrow \infty$. This completes the proof of the theorem. \square

In the next theorem of this section, we will study the global asymptotic stability of the positive equilibrium of system 3.

Theorem 3.4. *Consider system 3 where 15 holds true. Also suppose that*

$$\frac{\beta_1 e^{-a_1}}{\gamma_1 + c_1} + \frac{\beta_2 e^{-c_1}}{\gamma_2 + a_1} + \frac{b_1 d_1 + \beta_1 \beta_2 e^{-a_1 - c_1}}{(\gamma_1 + c_1)(\gamma_2 + a_1)} < 1. \tag{22}$$

Then the unique positive equilibrium point (\bar{x}, \bar{y}) of system 3 is globally asymptotically stable.

Proof. First we will prove that (\bar{x}, \bar{y}) is locally asymptotically stable. The characteristic equation of the Jacobian matrix $J_F(\bar{x}, \bar{y})$ about (\bar{x}, \bar{y}) is given by

$$\lambda^4 + p_2 \lambda^2 + p_4 = 0, \tag{23}$$

where

$$\begin{aligned} p_2 &= \frac{\beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}} + \frac{\beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}} - \frac{\bar{x} \cdot \bar{y}}{(\gamma_1 + \bar{y})(\gamma_2 + \bar{x})}, \\ p_4 &= \frac{\beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}} \cdot \frac{\beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}}. \end{aligned}$$

From condition 22, we get

$$\begin{aligned} |p_2| + |p_4| &= \frac{\beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}} + \frac{\beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}} + \frac{\bar{x} \cdot \bar{y}}{(\gamma_1 + \bar{y})(\gamma_2 + \bar{x})} + \frac{\beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}} \cdot \frac{\beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}} \\ &\leq \frac{\beta_1 e^{-a_1}}{\gamma_1 + c_1} + \frac{\beta_2 e^{-c_1}}{\gamma_2 + a_1} + \frac{b_1 d_1 + \beta_1 \beta_2 e^{-a_1 - c_1}}{(\gamma_1 + c_1)(\gamma_2 + a_1)} < 1. \end{aligned}$$

Therefore, follows Remark 1.3.1 of reference [10], all the roots of equation 23 are of modulus less than 1, and it follows from Lemma 2.3 that the unique positive equilibrium point (\bar{x}, \bar{y}) of system 3 is locally asymptotically stable. Using Theorem

3.3, we obtain that (\bar{x}, \bar{y}) is globally asymptotically stable. This completes the proof of the theorem. \square

3.1.3. *Rate of convergence.* In this section, we give the rate of convergence of a solution that converges to the equilibrium of the systems 3. Similar method can be found in [8, 9].

Let $\{(x_n, y_n)\}$ be an arbitrary solution of system 3 such that $\lim_{n \rightarrow \infty} x_n = \bar{x}$, and $\lim_{n \rightarrow \infty} y_n = \bar{y}$, where $\bar{x} \in [a_1, b_1]$, and $\bar{y} \in [c_1, d_1]$. To find the error terms, one has from the system 3

$$\begin{aligned} x_{n+1} - \bar{x} &= \frac{\alpha_1 + \beta_1 e^{-x_{n-1}}}{\gamma_1 + y_n} - \frac{\alpha_1 + \beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}} \\ &= \frac{(\alpha_1 + \beta_1 e^{-x_{n-1}})(\gamma_1 + \bar{y}) - (\alpha_1 + \beta_1 e^{-\bar{x}})(\gamma_1 + y_n)}{(\gamma_1 + y_n)(\gamma_1 + \bar{y})} \\ &= \frac{-\alpha_1(y_n - \bar{y}) + \beta_1 \gamma_1 (e^{-x_{n-1}} - e^{-\bar{x}}) + \beta_1 (e^{-x_{n-1}} \bar{y} - e^{-\bar{x}} y_n)}{(\gamma_1 + y_n)(\gamma_1 + \bar{y})} \\ &= \frac{-\alpha_1(y_n - \bar{y}) - \beta_1 \gamma_1 e^{-x_{n-1}} (e^{x_{n-1} - \bar{x}} - 1)}{(\gamma_1 + y_n)(\gamma_1 + \bar{y})} \\ &\quad + \frac{\beta_1 (e^{-x_{n-1}} \bar{y} - e^{-x_{n-1}} y_n + e^{-x_{n-1}} y_n - e^{-\bar{x}} y_n)}{(\gamma_1 + y_n)(\gamma_1 + \bar{y})} \\ &= -\frac{\beta_1 e^{-x_{n-1}} (e^{x_{n-1} - \bar{x}} - 1)}{(\gamma_1 + \bar{y})(x_{n-1} - \bar{x})} (x_{n-1} - \bar{x}) - \frac{\alpha_1 + \beta_1 e^{-x_{n-1}}}{(\gamma_1 + y_n)(\gamma_1 + \bar{y})} (y_n - \bar{y}), \end{aligned}$$

and

$$\begin{aligned} y_{n+1} - \bar{y} &= \frac{\alpha_2 + \beta_2 e^{-y_{n-1}}}{\gamma_2 + x_n} - \frac{\alpha_2 + \beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}} \\ &= \frac{(\alpha_2 + \beta_2 e^{-y_{n-1}})(\gamma_2 + \bar{x}) - (\alpha_2 + \beta_2 e^{-\bar{y}})(\gamma_2 + x_n)}{(\gamma_2 + x_n)(\gamma_2 + \bar{x})} \\ &= \frac{-\alpha_2(x_n - \bar{x}) + \beta_2 \gamma_2 (e^{-y_{n-1}} - e^{-\bar{y}}) + \beta_2 (e^{-y_{n-1}} \bar{x} - e^{-\bar{y}} x_n)}{(\gamma_2 + x_n)(\gamma_2 + \bar{x})} \\ &= \frac{-\alpha_2(x_n - \bar{x}) - \beta_2 \gamma_2 e^{-y_{n-1}} (e^{y_{n-1} - \bar{y}} - 1)}{(\gamma_2 + x_n)(\gamma_2 + \bar{x})} \\ &\quad + \frac{\beta_2 (e^{-y_{n-1}} \bar{x} - e^{-y_{n-1}} x_n + e^{-y_{n-1}} x_n - e^{-\bar{y}} x_n)}{(\gamma_2 + x_n)(\gamma_2 + \bar{x})} \\ &= -\frac{\alpha_2 + \beta_2 e^{-y_{n-1}}}{(\gamma_2 + x_n)(\gamma_2 + \bar{x})} (x_n - \bar{x}) - \frac{\beta_2 e^{-y_{n-1}} (e^{y_{n-1} - \bar{y}} - 1)}{(\gamma_2 + \bar{x})(y_{n-1} - \bar{y})} (y_{n-1} - \bar{y}). \end{aligned}$$

Let $e_n^1 = x_n - \bar{x}$, and $e_n^2 = y_n - \bar{y}$, then one has

$$\begin{aligned} e_{n+1}^1 &= a_n e_{n-1}^1 + b_n e_n^2, \\ e_{n+1}^2 &= c_n e_n^1 + d_n e_{n-1}^2, \end{aligned}$$

where

$$\begin{aligned} a_n &= -\frac{\beta_1 e^{-x_{n-1}} (e^{x_{n-1} - \bar{x}} - 1)}{(\gamma_1 + \bar{y})(x_{n-1} - \bar{x})}, & b_n &= -\frac{\alpha_1 + \beta_1 e^{-x_{n-1}}}{(\gamma_1 + y_n)(\gamma_1 + \bar{y})}, \\ c_n &= -\frac{\alpha_2 + \beta_2 e^{-y_{n-1}}}{(\gamma_2 + x_n)(\gamma_2 + \bar{x})}, & d_n &= -\frac{\beta_2 e^{-y_{n-1}} (e^{y_{n-1} - \bar{y}} - 1)}{(\gamma_2 + \bar{x})(y_{n-1} - \bar{y})}. \end{aligned}$$

Moreover,

$$\begin{aligned} \lim_{n \rightarrow \infty} a_n &= -\frac{\beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}}, & \lim_{n \rightarrow \infty} b_n &= -\frac{\bar{x}}{\gamma_1 + \bar{y}}, \\ \lim_{n \rightarrow \infty} c_n &= -\frac{\bar{y}}{\gamma_2 + \bar{x}}, & \lim_{n \rightarrow \infty} d_n &= -\frac{\beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}}. \end{aligned}$$

So, the limiting system of the error terms can be written as

$$\begin{pmatrix} e_{n+1}^1 \\ e_n^1 \\ e_{n+1}^2 \\ e_n^2 \end{pmatrix} = \begin{pmatrix} 0 & -\frac{\beta_1 e^{-\bar{x}}}{\gamma_1 + \bar{y}} & -\frac{\bar{x}}{\gamma_1 + \bar{y}} & 0 \\ 1 & 0 & 0 & 0 \\ -\frac{\bar{y}}{\gamma_2 + \bar{x}} & 0 & 0 & -\frac{\beta_2 e^{-\bar{y}}}{\gamma_2 + \bar{x}} \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} e_n^1 \\ e_{n-1}^1 \\ e_n^2 \\ e_{n-1}^2 \end{pmatrix}$$

which similar to the linearized system of 3 about the equilibrium point (\bar{x}, \bar{y}) . Using Proposition 2.6 and 2.7, one has the following result.

Theorem 3.5. *Assume that $\{(x_n, y_n)\}$ be a positive solution of system 3 such that $\lim_{n \rightarrow \infty} x_n = \bar{x}$, and $\lim_{n \rightarrow \infty} y_n = \bar{y}$, where $\bar{x} \in [a_1, b_1]$ and $\bar{y} \in [c_1, d_1]$. Then the*

error vector $e_n = \begin{pmatrix} e_n^1 \\ e_{n-1}^1 \\ e_n^2 \\ e_{n-1}^2 \end{pmatrix}$ of every solution of (3) satisfies both of the following asymptotic relations:

$$\lim_{n \rightarrow \infty} (||e_n||)^{\frac{1}{n}} = |\lambda_i|, \quad \lim_{n \rightarrow \infty} \frac{||e_{n+1}||}{||e_n||} = |\lambda_i|, \quad i = 1, 2, 3, 4,$$

where λ_i is one of the charateristic roots of Jacobian matrix $J_F(\bar{x}, \bar{y})$.

3.2. Global behavior of solutions of system 4.

3.2.1. *Boundedness and persistence.* In the following lemma, we study the boundedness and persistence of the positive solutions of system 4.

Lemma 3.6. *Every positive solution $\{(x_n, y_n)\}$ of system 4 is bounded and persists.*

Proof. Let $\{(x_n, y_n)\}$ be a positive solution of system 4. Similarly as Lemma 3.1, for $n = 3, 4, 5, \dots$ by induction, we get

$$x_n \in [a_2, b_2], \quad y_n \in [c_2, d_2],$$

where

$$\begin{aligned} a_2 &= \frac{\alpha_1 + \beta_1 e^{-\frac{\alpha_2 + \beta_2}{\gamma_2}}}{\gamma_1 + \frac{\alpha_1 + \beta_1}{\gamma_1}}, & b_2 &= \frac{\alpha_1 + \beta_1}{\gamma_1}, \\ c_2 &= \frac{\alpha_2 + \beta_2 e^{-\frac{\alpha_1 + \beta_1}{\gamma_1}}}{\gamma_2 + \frac{\alpha_2 + \beta_2}{\gamma_2}}, & d_2 &= \frac{\alpha_2 + \beta_2}{\gamma_2}. \end{aligned}$$

So the proof is complete. □

Corollary 3.7. *Let $\{(x_n, y_n)\}$ be a positive solution of system 4. Then $[a_2, b_2] \times [c_2, d_2]$ is an invariant set for system 4.*

3.2.2. *Stability analysis.* In this section, we shall investigate the asymptotic behavior of system 4. Let (\bar{x}, \bar{y}) be the equilibrium point of system 4 then

$$\bar{x} = \frac{\alpha_1 + \beta_1 e^{-\bar{y}}}{\gamma_1 + \bar{x}}, \quad \bar{y} = \frac{\alpha_2 + \beta_2 e^{-\bar{x}}}{\gamma_2 + \bar{y}}.$$

The linearized form of system 4 about the equilibrium point (\bar{x}, \bar{y}) is given by

$$X_{n+1} = J_F(\bar{x}, \bar{y})X_n,$$

where $X_n = \begin{pmatrix} x_n \\ x_{n-1} \\ y_n \\ y_{n-1} \end{pmatrix}$ and $J_F(\bar{x}, \bar{y}) = \begin{pmatrix} -\frac{\bar{x}}{\gamma_1 + \bar{x}} & 0 & 0 & -\frac{\beta_1 e^{-\bar{y}}}{\gamma_1 + \bar{x}} \\ 1 & 0 & 0 & 0 \\ 0 & -\frac{\beta_2 e^{-\bar{x}}}{\gamma_2 + \bar{y}} & -\frac{\bar{y}}{\gamma_2 + \bar{y}} & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$.

In the following theorem, we show the asymptotic behavior of the positive solutions of system 4.

Theorem 3.8. *Suppose that the following relation holds true:*

$$\beta_1 \beta_2 < \gamma_1 \gamma_2. \tag{24}$$

Then system 4 has a unique positive equilibrium (\bar{x}, \bar{y}) and every positive solution of system 4 tends to the unique positive equilibrium as $n \rightarrow \infty$.

Proof. Consider the following functions:

$$f(x, y) = \frac{\alpha_1 + \beta_1 e^{-y}}{\gamma_1 + x}, \quad g(x, y) = \frac{\alpha_2 + \beta_2 e^{-x}}{\gamma_2 + y},$$

where $x \in I_3 = [a_2, b_2], y \in I_4 = [c_2, d_2]$ which implies that $f(x, y) \in I_3, g(x, y) \in I_4$ and so that $f : I_3 \times I_4 \rightarrow I_3, g : I_3 \times I_4 \rightarrow I_4$. Then, it is easy to see that $f(x, y), g(x, y)$ are decreasing with respect to x (resp. y) for all y (resp. x). Let (m, M, r, R) be a solution of the system

$$\begin{aligned} m &= f(M, R), & M &= f(m, r), \\ r &= g(M, R), & R &= g(m, r). \end{aligned}$$

Then, one has

$$m = \frac{\alpha_1 + \beta_1 e^{-R}}{\gamma_1 + M}, \quad M = \frac{\alpha_1 + \beta_1 e^{-r}}{\gamma_1 + m}, \quad r = \frac{\alpha_2 + \beta_2 e^{-M}}{\gamma_2 + R}, \quad R = \frac{\alpha_2 + \beta_2 e^{-m}}{\gamma_2 + r}. \tag{25}$$

From 25, we get

$$\begin{aligned} \beta_1 e^{-r} &= M(\gamma_1 + m) - \alpha_1, & \beta_1 e^{-R} &= m(\gamma_1 + M) - \alpha_1, \\ \beta_2 e^{-m} &= R(\gamma_2 + r) - \alpha_2, & \beta_2 e^{-M} &= r(\gamma_2 + R) - \alpha_2, \end{aligned}$$

which imply that

$$\begin{aligned} \beta_1(e^{-r} - e^{-R}) &= \gamma_1(M - m), \\ \beta_2(e^{-m} - e^{-M}) &= \gamma_2(R - r). \end{aligned} \tag{26}$$

Moreover, we get

$$\begin{aligned} e^R - e^r &= e^\xi(R - r), \quad \min\{R, r\} \leq \xi \leq \max\{R, r\}, \\ e^M - e^m &= e^\theta(M - m), \quad \min\{M, m\} \leq \theta \leq \max\{M, m\}. \end{aligned} \tag{27}$$

Then from 26 and 27, we have

$$\begin{aligned} M - m &= \frac{\beta_1}{\gamma_1}(e^{-r} - e^{-R}) = \frac{\beta_1}{\gamma_1}e^{-r-R}(e^R - e^r) = \frac{\beta_1}{\gamma_1}e^{-r-R+\xi}(R - r), \\ R - r &= \frac{\beta_2}{\gamma_2}(e^{-m} - e^{-M}) = \frac{\beta_2}{\gamma_2}e^{-m-M}(e^M - e^m) = \frac{\beta_2}{\gamma_2}e^{-m-M+\theta}(M - m), \end{aligned} \tag{28}$$

and so

$$|M - m| \leq \frac{\beta_1}{\gamma_1}|R - r|, \quad |R - r| \leq \frac{\beta_2}{\gamma_2}|M - m|.$$

Therefore, we get

$$\left(1 - \frac{\beta_1\beta_2}{\gamma_1\gamma_2}\right)|M - m| \leq 0, \quad \left(1 - \frac{\beta_1\beta_2}{\gamma_1\gamma_2}\right)|R - r| \leq 0. \tag{29}$$

Finally, from 24 and 29, it follows that $M = m$ and $R = r$. Therefore, from Lemma 2.5, it follows that system 4 has a unique positive equilibrium (\bar{x}, \bar{y}) and every positive solution of system 4 tends to the unique positive equilibrium as $n \rightarrow \infty$. This completes the proof of the theorem. \square

In the next theorem of this section, we will study the global asymptotic stability of the positive equilibrium of system 4.

Theorem 3.9. *Consider system 4 where 24 holds true. Also suppose that*

$$\frac{b_2}{\gamma_1 + a_2} + \frac{d_2}{\gamma_2 + c_2} + \frac{b_2d_2 + \beta_1\beta_2e^{-a_2-c_2}}{(\gamma_1 + a_2)(\gamma_2 + c_2)} < 1. \tag{30}$$

Then the unique positive equilibrium point (\bar{x}, \bar{y}) of system 4 is globally asymptotically stable.

Proof. First we will prove that (\bar{x}, \bar{y}) is locally asymptotically stable. The characteristic equation of the Jacobian matrix $J_F(\bar{x}, \bar{y})$ about (\bar{x}, \bar{y}) is given by

$$\lambda^4 + q_1\lambda^3 + q_2\lambda^2 + q_4 = 0, \tag{31}$$

where

$$\begin{aligned} q_1 &= \frac{\bar{x}}{\gamma_1 + \bar{x}} + \frac{\bar{y}}{\gamma_2 + \bar{y}}, \\ q_2 &= \frac{\bar{x}\bar{y}}{(\gamma_1 + \bar{x})(\gamma_2 + \bar{y})}, \\ q_4 &= -\frac{\beta_1e^{-\bar{y}}}{\gamma_1 + \bar{x}} \cdot \frac{\beta_2e^{-\bar{x}}}{\gamma_2 + \bar{y}}. \end{aligned}$$

From condition 30, we get

$$\begin{aligned} |q_1| + |q_2| + |q_4| &= \frac{\bar{x}}{\gamma_1 + \bar{x}} + \frac{\bar{y}}{\gamma_2 + \bar{y}} + \frac{\bar{x}\bar{y}}{(\gamma_1 + \bar{x})(\gamma_2 + \bar{y})} + \frac{\beta_1e^{-\bar{y}}}{\gamma_1 + \bar{x}} \cdot \frac{\beta_2e^{-\bar{x}}}{\gamma_2 + \bar{y}} \\ &\leq \frac{b_2}{\gamma_1 + a_2} + \frac{d_2}{\gamma_2 + c_2} + \frac{b_2d_2 + \beta_1\beta_2e^{-a_2-c_2}}{(\gamma_1 + a_2)(\gamma_2 + c_2)} < 1. \end{aligned}$$

Therefore, follows Remark 1.3.1 of reference [10], all the roots of equation 31 are of modulus less than 1, and it follows from Lemma 2.3 that the unique positive equilibrium point (\bar{x}, \bar{y}) of system 4 is locally asymptotically stable. Using Theorem 3.8, we obtain that (\bar{x}, \bar{y}) is globally asymptotically stable. This completes the proof of the theorem. \square

3.2.3. *Rate of convergence.* In this section, we give the rate of convergence of a solution that converges to the equilibrium of the systems 4.

Let $\{(x_n, y_n)\}$ be an arbitrary solution of system 4 such that $\lim_{n \rightarrow \infty} x_n = \bar{x}$, and $\lim_{n \rightarrow \infty} y_n = \bar{y}$, where $\bar{x} \in [a_2, b_2]$, and $\bar{y} \in [c_2, d_2]$. To find the error terms, one has from the system 4

$$\begin{aligned} x_{n+1} - \bar{x} &= \frac{\alpha_1 + \beta_1 e^{-y_{n-1}}}{\gamma_1 + x_n} - \frac{\alpha_1 + \beta_1 e^{-\bar{y}}}{\gamma_1 + \bar{x}} \\ &= \frac{(\alpha_1 + \beta_1 e^{-y_{n-1}})(\gamma_1 + \bar{x}) - (\alpha_1 + \beta_1 e^{-\bar{y}})(\gamma_1 + x_n)}{(\gamma_1 + x_n)(\gamma_1 + \bar{x})} \\ &= \frac{-\alpha_1(x_n - \bar{x}) + \beta_1 \gamma_1 (e^{-y_{n-1}} - e^{-\bar{y}}) + \beta_1 (e^{-y_{n-1}} \bar{x} - e^{-\bar{y}} x_n)}{(\gamma_1 + x_n)(\gamma_1 + \bar{x})} \\ &= \frac{-\alpha_1(x_n - \bar{x}) - \beta_1 \gamma_1 e^{-y_{n-1}} (e^{y_{n-1} - \bar{y}} - 1)}{(\gamma_1 + x_n)(\gamma_1 + \bar{x})} \\ &\quad + \frac{\beta_1 (e^{-y_{n-1}} \bar{x} - e^{-y_{n-1}} x_n + e^{-y_{n-1}} x_n - e^{-\bar{y}} x_n)}{(\gamma_1 + x_n)(\gamma_1 + \bar{x})} \\ &= -\frac{\alpha_1 + \beta_1 e^{-y_{n-1}}}{(\gamma_1 + x_n)(\gamma_1 + \bar{x})} (x_n - \bar{x}) - \frac{\beta_1 e^{-y_{n-1}} (e^{y_{n-1} - \bar{y}} - 1)}{(\gamma_1 + \bar{x})(y_{n-1} - \bar{y})} (y_{n-1} - \bar{y}), \end{aligned}$$

and

$$\begin{aligned} y_{n+1} - \bar{y} &= \frac{\alpha_2 + \beta_2 e^{-x_{n-1}}}{\gamma_2 + y_n} - \frac{\alpha_2 + \beta_2 e^{-\bar{x}}}{\gamma_2 + \bar{y}} \\ &= \frac{(\alpha_2 + \beta_2 e^{-x_{n-1}})(\gamma_2 + \bar{y}) - (\alpha_2 + \beta_2 e^{-\bar{x}})(\gamma_2 + y_n)}{(\gamma_2 + y_n)(\gamma_2 + \bar{y})} \\ &= \frac{-\alpha_2(y_n - \bar{y}) + \beta_2 \gamma_2 (e^{-x_{n-1}} - e^{-\bar{x}}) + \beta_2 (e^{-x_{n-1}} \bar{y} - e^{-\bar{x}} y_n)}{(\gamma_2 + y_n)(\gamma_2 + \bar{y})} \\ &= \frac{-\alpha_2(y_n - \bar{y}) - \beta_2 \gamma_2 e^{-x_{n-1}} (e^{x_{n-1} - \bar{x}} - 1)}{(\gamma_2 + y_n)(\gamma_2 + \bar{y})} \\ &\quad + \frac{\beta_2 (e^{-x_{n-1}} \bar{y} - e^{-x_{n-1}} y_n + e^{-x_{n-1}} y_n - e^{-\bar{x}} y_n)}{(\gamma_2 + y_n)(\gamma_2 + \bar{y})} \\ &= -\frac{\beta_2 e^{-x_{n-1}} (e^{x_{n-1} - \bar{x}} - 1)}{(\gamma_2 + \bar{y})(x_{n-1} - \bar{x})} (x_{n-1} - \bar{x}) - \frac{\alpha_2 + \beta_2 e^{-x_{n-1}}}{(\gamma_2 + y_n)(\gamma_2 + \bar{y})} (y_n - \bar{y}). \end{aligned}$$

Let $e_n^1 = x_n - \bar{x}$, and $e_n^2 = y_n - \bar{y}$, then one has

$$\begin{aligned} e_{n+1}^1 &= f_n e_n^1 + g_n e_{n-1}^2, \\ e_{n+1}^2 &= h_n e_{n-1}^1 + k_n e_n^2, \end{aligned}$$

where

$$\begin{aligned} f_n &= -\frac{\alpha_1 + \beta_1 e^{-y_{n-1}}}{(\gamma_1 + x_n)(\gamma_1 + \bar{x})}, & g_n &= -\frac{\beta_1 e^{-y_{n-1}} (e^{y_{n-1} - \bar{y}} - 1)}{(\gamma_1 + \bar{x})(y_{n-1} - \bar{y})}, \\ h_n &= -\frac{\beta_2 e^{-x_{n-1}} (e^{x_{n-1} - \bar{x}} - 1)}{(\gamma_2 + \bar{y})(x_{n-1} - \bar{x})}, & k_n &= -\frac{\alpha_2 + \beta_2 e^{-x_{n-1}}}{(\gamma_2 + y_n)(\gamma_2 + \bar{y})}. \end{aligned}$$

Moreover,

$$\lim_{n \rightarrow \infty} f_n = -\frac{\bar{x}}{\gamma_1 + \bar{x}}, \quad \lim_{n \rightarrow \infty} g_n = -\frac{\beta_1 e^{-\bar{y}}}{\gamma_1 + \bar{x}},$$

$$\lim_{n \rightarrow \infty} h_n = -\frac{\beta_2 e^{-\bar{x}}}{\gamma_2 + \bar{y}}, \quad \lim_{n \rightarrow \infty} k_n = -\frac{\bar{y}}{\gamma_2 + \bar{y}}.$$

So, the limiting system of the error terms can be written as

$$\begin{pmatrix} e_{n+1}^1 \\ e_n^1 \\ e_{n+1}^2 \\ e_n^2 \end{pmatrix} = \begin{pmatrix} -\frac{\bar{x}}{\gamma_1 + \bar{x}} & 0 & 0 & -\frac{\beta_1 e^{-\bar{y}}}{\gamma_1 + \bar{x}} \\ 1 & 0 & 0 & 0 \\ 0 & -\frac{\beta_2 e^{-\bar{x}}}{\gamma_2 + \bar{y}} & -\frac{\bar{y}}{\gamma_2 + \bar{y}} & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} e_n^1 \\ e_{n-1}^1 \\ e_n^2 \\ e_{n-1}^2 \end{pmatrix}$$

which similar to the linearized system of 4 about the equilibrium point (\bar{x}, \bar{y}) . Using Proposition 2.6 and 2.7, one has the following result.

Theorem 3.10. Assume that $\{(x_n, y_n)\}$ be a positive solution of system 4 such that $\lim_{n \rightarrow \infty} x_n = \bar{x}$, and $\lim_{n \rightarrow \infty} y_n = \bar{y}$, where $\bar{x} \in [a_2, b_2]$ and $\bar{y} \in [c_2, d_2]$. Then

the error vector $e_n = \begin{pmatrix} e_n^1 \\ e_{n-1}^1 \\ e_n^2 \\ e_{n-1}^2 \end{pmatrix}$ of every solution of 4 satisfies both of the following asymptotic relations:

$$\lim_{n \rightarrow \infty} (||e_n||)^{\frac{1}{n}} = |\lambda_i|, \quad \lim_{n \rightarrow \infty} \frac{||e_{n+1}||}{||e_n||} = |\lambda_i|, \quad i = 1, 2, 3, 4,$$

where λ_i is one of the charateristic roots of Jacobian matrix $J_F(\bar{x}, \bar{y})$.

4. Numerical simulations. In an effort to affirm our theoretical dialogue, we consider several numerical examples. These examples represent different types of qualitative behavior of solutions of the systems 3 and 4. All plots in this section are drawn with MATLAB.

Example 4.1. Let $\alpha_1 = 30, \beta_1 = 1.4, \gamma_1 = 1.5, \alpha_2 = 45, \beta_2 = 2.5$, and $\gamma_2 = 2.8$. Then system 3 can be written as

$$x_{n+1} = \frac{30 + 1.4e^{-x_{n-1}}}{1.5 + y_n}, \quad y_{n+1} = \frac{45 + 2.5e^{-y_{n-1}}}{2.8 + x_n}, \tag{32}$$

with initial conditions $x_{-1} = 0.59, x_0 = 0.61, y_{-1} = 0.96$, and $y_0 = 0.94$.




FIGURE 1. Plot of x_n for the system 32

In this case, the unique positive equilibrium point of the system 32 is given by $(\bar{x}, \bar{y}) = (3.455959, 7.193442)$. Moreover, the plot of x_n is shown in Figure 1, the plot of y_n is shown in Figure 2, and an attractor of the system 32 is shown in Figure 3.




FIGURE 2. Plot of y_n for the system 32




FIGURE 3. An attractor of the system 32

Example 4.2. Let $\alpha_1 = 0.2, \beta_1 = 19, \gamma_1 = 4, \alpha_2 = 0.3, \beta_2 = 20,$ and $\gamma_2 = 2.$ Then system 3 can be written as

$$x_{n+1} = \frac{0.2 + 19e^{-x_{n-1}}}{4 + y_n}, \quad y_{n+1} = \frac{0.3 + 20e^{-y_{n-1}}}{2 + x_n}, \tag{33}$$

with initial conditions $x_{-1} = 1, x_0 = 1, y_{-1} = 3,$ and $y_0 = 3.$

In this case, the unique positive equilibrium point of the system 33 is unstable. Moreover, the plot of x_n is shown in Figure 4, the plot of y_n is shown in Figure 5, and a phase portrait of system 33 is shown in Figure 6.

Example 4.3. Let $\alpha_1 = 401, \beta_1 = 1, \gamma_1 = 1.75, \alpha_2 = 395, \beta_2 = 1.5,$ and $\gamma_2 = 1.$ Then system 4 can be written as

$$x_{n+1} = \frac{401 + 1e^{-y_{n-1}}}{1.75 + x_n}, \quad y_{n+1} = \frac{395 + 1.5e^{-x_{n-1}}}{1 + y_n}, \tag{34}$$

with initial conditions $x_{-1} = 15, x_0 = 24.5, y_{-1} = 15,$ and $y_0 = 25.$




FIGURE 4. Plot of x_n for the system 33FIGURE 5. Plot of y_n for the system 33

FIGURE 6. Phase portrait of system 33

In this case, the unique positive equilibrium point of the system 34 is given by $(\bar{x}, \bar{y}) = (19.169092, 19.380895)$. Moreover, the plot of x_n is shown in Figure 7, the plot of y_n is shown in Figure 8, and an attractor of the system 34 is shown in Figure 9.

Example 4.4. Let $\alpha_1 = 4, \beta_1 = 10, \gamma_1 = 1.6, \alpha_2 = 5.5, \beta_2 = 10.8$, and $\gamma_2 = 1$. Then system 4 can be written as

$$x_{n+1} = \frac{4 + 10e^{-y_{n-1}}}{1.6 + x_n}, \quad y_{n+1} = \frac{5.5 + 10.8e^{-x_{n-1}}}{1 + y_n}, \quad (35)$$




FIGURE 7. Plot of x_n for the system 34




FIGURE 8. Plot of y_n for the system 34




FIGURE 9. An attractor of the system 34

with initial conditions $x_{-1} = 1.3, x_0 = 1.5, y_{-1} = 2$, and $y_0 = 1.7$.

In this case, the unique positive equilibrium point of the system (35) is unstable. Moreover, the plot of x_n is shown in Figure 10, the plot of y_n is shown in Figure 11, and a phase portrait of system 35 is shown in Figure 12.

5. Conclusion. In this study, we investigate the qualitative behavior of some systems of exponential difference equations. We have proved the boundedness and




FIGURE 10. Plot of x_n for the system 35FIGURE 11. Plot of y_n for the system 35

FIGURE 12. Phase portrait of system 35

persistence of positive solutions of system 3 and 4. Moreover, we have shown that unique positive equilibrium point of system 3 and 4 is locally as well as globally asymptotically stable under certain parametric conditions. Furthermore, the rate of convergence of positive solutions of 3 and 4 which converges to its unique positive equilibrium point is demonstrated. Finally, some illustrative numerical examples are provided.

Acknowledgments. This research is funded by Hung Yen University of Technology and Education under grand number UTEHY.L.2020.11.

REFERENCES

- [1] R. P. Agarwal, *Difference Equations and Inequalities*, 2nd edition, Dekker, New York, 2000.
- [2] Q. Din, [Global stability of a population models](#), *Chaos, Solitons & Fractals*, **59** (2014), 119–128.
- [3] Q. Din and E. M. Elsayed, Stability analysis of a discrete ecological model, *Computational Ecology and Software*, **4** (2014), 89–103.
- [4] H. El-Metwally, E. A. Grove, G. Ladas, R. Levins and M. Radin, [On the difference equation \$x_{n+1} = \alpha + \beta x_{n-1} e^{-x_n}\$](#) , *Nonlinear Anal.*, **47** (2001), 4623–4634.
- [5] N. Fotiades and G. Papaschinopoulos, [Existence, uniqueness and attractivity of prime period two solution for a difference equation of exponential form](#), *Appl. Math. Comput.*, **218** (2012), 11648–11653.
- [6] E. A. Grove and G. Ladas, *Periodicities in Nonlinear Difference Equations*, Chapman and Hall/CRC, Boca Raton, Fla, 2005.
- [7] E. A. Grove, G. Ladas, N. R. Prokup and R. Levins, On the global behavior of solutions of a biological model, *Comm. Appl. Nonlinear Anal.*, **7** (2000), 33–46.
- [8] T. Hong Thai, Asymptotic behavior of the solution of a system of difference equations, *Int. J. Difference Equ.*, **13** (2018), 157–171.
- [9] T. Hong Thai and V. Van Khuong, [Stability analysis of a system of second-order difference equations](#), *Math. Methods Appl. Sci.*, **39** (2016), 3691–3700.
- [10] V. L. Kocić and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1993.
- [11] M. R. S. Kulenović and G. Ladas, *Dynamics of Second Order Rational Difference Equations*, Chapman and Hall/CRC, Boca Raton, Fla, 2002.
- [12] I. Ozturk, F. Bozkurt and S. Ozen, [On the difference equation \$y_{n+1} = \frac{\alpha + \beta e^{-y_n}}{\gamma + y_{n-1}}\$](#) , *Appl. Math. Comput.*, **181** (2006), 1387–1393.
- [13] G. Papaschinopoulos, G. Ellina and K. B. Papadopoulos, [Asymptotic behavior of the positive solutions of an exponential type system of difference equations](#), *Appl. Math. Comput.*, **245** (2014), 181–190.
- [14] G. Papaschinopoulos, N. Fotiades and C. J. Schinas, [On a system of difference equations including negative exponential terms](#), *J. Difference Equ. Appl.*, **20** (2014), 717–732.
- [15] G. Papaschinopoulos, M. A. Radin and C. J. Schinas, [On the system of two difference equations of exponential form: \$x_{n+1} = a + bx_{n-1} e^{-y_n}\$, \$y_{n+1} = c + dy_{n-1} e^{-x_n}\$](#) , *Math. Comput. Modelling*, **54** (2011), 2969–2977.
- [16] G. Papaschinopoulos, M. Radin and C. J. Schinas, [Study of the asymptotic behavior of the solutions of three systems of difference equations of exponential form](#), *Appl. Math. Comput.*, **218** (2012), 5310–5318.
- [17] G. Papaschinopoulos and C. J. Schinas, [On the dynamics of two exponential type systems of difference equations](#), *Comput. Math. Appl.*, **64** (2012), 2326–2334.
- [18] M. Pituk, [More on Poincaré’s and Peron’s theorems for difference equations](#), *J. Difference Equ. Appl.*, **8** (2002), 201–216.
- [19] H. Sedaghat, *Nonlinear Difference Equations: Theory with Applications to Social Science Models*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2003.
- [20] W. Wang and H. Feng, [On the dynamics of positive solutions for the difference equation in a new population model](#), *J. Nonlinear Sci. Appl.*, **9** (2016), 1748–1754.

Received for publication June 2021; early access October 2021.

E-mail address: tranhongthai@utehy.edu.vn

E-mail address: nguyenanhhdai2@gmail.com

E-mail address: anhpt@hus.edu.vn