



Research article

Sign-changing solutions of critical quasilinear Kirchhoff-Schrödinger-Poisson system with logarithmic nonlinearity

Hui Jian¹, Shenghao Feng² and Li Wang^{1,*}

¹ College of Science, East China Jiaotong University, Nanchang 330013, Jiangxi, China

² Department of Mathematics, Nanchang University, Nanchang 330031, Jiangxi, China

* Correspondence: Email: wangli.423@163.com.

Abstract: In the present paper, we study the following Kirchhoff-Schrödinger-Poisson system with logarithmic and critical nonlinearity:

-(a + b integral |nabla u|^2 dx) Delta u + V(x)u - 1/2 u Delta (u^2) + phi u = lambda |u|^{q-2} u ln |u|^2 + |u|^4 u, x in Omega,
-Delta phi = u^2, x in Omega,
u = phi = 0, x in partial Omega,

where lambda, b > 0, a > 1/4, 4 < q < 6, V(x) is a smooth potential function and Omega is a bounded domain in R^3 with Lipschitz boundary. Combining constraint variational method and perturbation method, we prove that the above problem has a least energy sign-changing solution u_0 which has precisely two nodal domains. Moreover, we show that the energy of u_0 is strictly larger than twice the ground state energy.

Keywords: quasilinear Kirchhoff-Schrödinger-Poisson; critical problem; logarithmic nonlinearity

Mathematics Subject Classification: 35A15, 35J60, 47G20

1. Introduction and main results

In this paper, we consider the existence of a least energy sign-changing solution of the following quasilinear Kirchhoff-Schrödinger-Poisson type system:

-(a + b integral |nabla u|^2 dx) Delta u + V(x)u - 1/2 u Delta (u^2) + phi u = lambda |u|^{q-2} u ln |u|^2 + |u|^4 u, x in Omega,
-Delta phi = u^2, x in Omega,
u = phi = 0, x in partial Omega, (1.1)

where $\lambda, b > 0, a > \frac{1}{4}, 4 < q < 6$, $V(x)$ is a smooth potential function and Ω is a bounded domain in \mathbb{R}^3 with Lipschitz boundary. After the pioneer work of Lions [17], some researchers began to pay attention to the following Kirchhoff Dirichlet problem:

$$\begin{cases} -(a + b \int_{\Omega} |\nabla u|^2 dx) \Delta u = f(x, u), & x \in \Omega, \\ u|_{\partial\Omega} = 0. \end{cases} \quad (1.2)$$

Problem (1.2) is related to a model firstly proposed by Kirchhoff [13] as an existension of the classical D'Alembert's wave equations for free vibration of elastic strings, which is related to the stationary analogue of the equation:

$$\rho \frac{\partial^2 u}{\partial t^2} - \left(\frac{P_0}{h} + \frac{E}{2L} \int_0^L \left| \frac{\partial u}{\partial x} \right|^2 dx \right) \frac{\partial^2 u}{\partial x^2} = 0,$$

where ρ, P_0, h, E, L are constants. Because problem (1.2) has nonlocal term $(\int_{\Omega} |\nabla u|^2 dx) \Delta u$, there are some difficulties in the study of the nonlocal problems by means of variational method.

In recent years, many interesting results on the existence of positive solutions, multiple solutions, bound state solutions, semiclassical state solutions and sign-changing solutions for (1.2) can be found in [1, 2, 4, 6, 7, 22, 24] and the references therein.

By using the constraint variational method and the quantitative deformation lemma, Wang [26] obtained the existence of a least energy sign-changing solution for the following Kirchhoff-type equation with critical growth:

$$\begin{cases} -\left(a + b \int_{\Omega} |\nabla u|^2 dx\right) \Delta u = |u|^4 u + \lambda f(x, u), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases} \quad (1.3)$$

where λ is large enough and f satisfies suitable conditions. Lately, Li and Wang [14] studied ground state sign-changing solutions for Kirchhoff equations with logarithmic nonlinearity:

$$\begin{cases} -\left(a + b \int_{\Omega} |\nabla u|^2 dx\right) \Delta u + V(x)u = |u|^{p-2} u \ln u^2, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases} \quad (1.4)$$

where $4 < p < 2^*$, they used constraint variational method, topological degree theory and some new energy estimate inequalities to prove the existence of ground state solutions and ground state sign-changing solutions. Recently, Liang and Rădulescu [16] got a more general result about problem (1.4) with critical growth.

Nevertheless, there are relatively few studies on quasilinear Schrödinger-Poisson system. Illner [12] first studied quasilinear Schrödinger-Poisson system. This quasilinear version of the nonlinear Schrödinger equation arises in several models of different physical phenomena, such as superfluid films, plasma physics, condensed matter theory, etc. (see [21, 23]). For quasilinear problem, by using the methods of perturbation and the Mountain Pass theorem, Feng [8] proved the existence of non-trivial solution to the following quasilinear Schrödinger-Poisson equation:

$$\begin{cases} -\Delta u + V(x)u + \phi u - \frac{1}{2}u\Delta(u^2) = f(x, u), & \text{in } \mathbb{R}^3, \\ -\Delta \phi = u^2, & \text{in } \mathbb{R}^3, \end{cases} \quad (1.5)$$

where $V \in C(\mathbb{R}^3, \mathbb{R})$, $\lim_{|x| \rightarrow \infty} V(x) = \infty$ and $V(x) \geq m > 0$ for some constant m to overcome the lack of compactness. Lately, under suitable condition of f , Chen and Tang [5] applied some new analytical techniques and non-Nehari manifold method investigated the existence of ground state sign-changing solutions for the following quasilinear Schrödinger equations with a Kirchhoff-type perturbation:

$$\begin{aligned} & \left(1 + b \int_{\mathbb{R}^3} g^2(u) |\nabla u|^2 dx\right) \left[-\operatorname{div}(g^2(u) \nabla u) + g(u)g'(u) |\nabla u|^2\right] \\ & + V(x)u = K(x)f(u). \end{aligned} \quad (1.6)$$

Figueiredo and Siciliano in [9, 10] paid close attention to two different critical systems with 4-Laplacian operator in \mathbb{R}^3 and a bounded domain in \mathbb{R}^2 , they obtained the existence and asymptotic behavior of nontrivial solutions. In [19], Massar studied a nonlocal Schrödinger-Poisson system with critical exponent, Wang [27] investigated nontrivial solutions of quasilinear Schrödinger-Kirchhoff-type equation with radial potentials. Fu and Zhu [15] considered the multiple solutions to a class of generalized quasilinear Schrödinger equations with a Kirchhoff-type perturbation. However, there are relatively few achievements on the so called quasilinear Kirchhoff-Schrödinger-Poisson type systems with critical growth, furthermore, few studies have included logarithmic terms about quasilinear problem. It is quite natural to ask: What is going to happen with logarithmic nonlinear terms for the critical quasilinear Kirchhoff Schrödinger-Poisson system? In this paper, we will show that there exists a least energy sign-changing solution.

According to the Lax-Milgram Theorem, for $u \in H_0^1(\Omega)$, there is a unique $\phi_u \in D_0^{1,2}(\Omega)$ that satisfies

$$-\Delta \phi_u = u^2.$$

Therefore, $(u, \phi) \in H_0^1(\Omega) \times H_0^1(\Omega)$ is a solution of (1.1) if, and only if, $\phi = \phi_u$ and $u \in H_0^1(\Omega)$ is a weak solution of the nonlocal problem

$$\begin{cases} -(a + b \int_{\Omega} |\nabla u|^2 dx) \Delta u + V(x)u - \frac{1}{2} u \Delta(u^2) + \phi_u u = \lambda |u|^{q-2} u \ln |u|^2 + |u|^4 u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega. \end{cases}$$

Let $H_V^1(\Omega) := \{u \in H_0^1(\Omega) \mid \int_{\Omega} V(x)u^2 dx < +\infty\}$, and $V \in C(\Omega, \mathbb{R})$, $V(x) \geq m > 0$ for some constant m . So, the functional associated with system (1.1) can be defined by

$$\begin{aligned} I_b^\lambda(u) &= \frac{1}{2} \int_{\Omega} (a |\nabla u|^2 + V(x)u^2) dx + \frac{b}{4} \left(\int_{\Omega} |\nabla u|^2 dx \right)^2 + \frac{1}{2} \int_{\Omega} |\nabla u|^2 u^2 dx \\ &+ \frac{1}{4} \int_{\Omega} \phi_u u^2 dx + \frac{2\lambda}{q^2} \int_{\Omega} |u|^q dx - \frac{\lambda}{q} \int_{\Omega} |u|^q \ln |u|^2 dx - \frac{1}{6} \int_{\Omega} |u|^6 dx. \end{aligned}$$

It is easy to see that $I_b^\lambda(u) \in C^1(H_V^1(\Omega), \mathbb{R})$. Moreover, for any $u, \varphi \in H_V^1(\Omega)$, we have

$$\begin{aligned} \langle (I_b^\lambda)'(u), \varphi \rangle &= \int_{\Omega} a \nabla u \nabla \varphi dx + b \int_{\Omega} |\nabla u|^2 dx \int_{\Omega} \nabla u \nabla \varphi dx + \int_{\Omega} V(x)u \varphi dx + \int_{\Omega} \phi_u u \varphi dx \\ &+ \int_{\Omega} (|\nabla u|^2 u \varphi + |u|^2 \nabla u \nabla \varphi) dx - \lambda \int_{\Omega} |u|^{q-2} u \varphi \ln |u|^2 dx - \int_{\Omega} |u|^4 u \varphi dx. \end{aligned}$$

There are some difficulties in applying variational method directly to the problem (1.1) because of the quasilinear term $\int_{\Omega} u^2 |\nabla u|^2 dx$. It appears that finding a suitable space where the matching functional has both smoothness and compactness qualities is unattainable. On the other hand, it is difficult to apply the dual approach since problem (1.1) exists nonlocal term. In order to overcome the lack of compactness caused by the critical term, we would employ the method from [3, 18]. In fact, we will use the approximation method by adding a 4-Laplacian operator, i.e., we consider the sign-changing critical point of the perturbed functional:

$$I_{b,\mu}^{\lambda}(u) = I_b^{\lambda}(u) + \frac{\mu}{4} \int_{\Omega} (|\nabla u|^4 + u^4) dx, \quad (1.7)$$

where $\mu \in (0, 1]$. Then by using the approximation technique, we get the existence of sign-changing solution of problem (1.1).

We first try to seek a minimizer of energy functional $I_{b,\mu}^{\lambda}$ over the following constraint:

$$\mathcal{M}_{b,\mu}^{\lambda} = \{u \in H_V^1(\Omega), u^{\pm} \neq 0 \text{ and } \langle (I_{b,\mu}^{\lambda})'(u), u^+ \rangle = \langle (I_{b,\mu}^{\lambda})'(u), u^- \rangle = 0\},$$

and consider a minimization problem of $I_{b,\mu}^{\lambda}$ on $\mathcal{M}_{b,\mu}^{\lambda}$, here

$$u(x) = u^+(x) + u^-(x), \quad u^+(x) = \max\{u(x), 0\} \text{ and } u^-(x) = \min\{u(x), 0\}.$$

We will prove that the minimizer is a critical point of $I_{b,\mu}^{\lambda}$ and obtain the convergence property as $\mu \rightarrow 0$, thus we get the least energy sign-changing solution of problem (1.1). Since problem (1.1) has nonlocal term and logarithmic nonlinearity, it is difficult to prove $\mathcal{M}_{b,\mu}^{\lambda} \neq \emptyset$. Inspired by [26], we combine modified Miranda's theorem [20], quantitative lemma, topological degree theory and perturbation method to prove that the minimizer of the constrained problem is also a least energy sign-changing solution.

Our main results of this paper are as follows:

Theorem 1.1. *Suppose that $V \in C(\Omega, \mathbb{R})$, $V(x) \geq m > 0$ for some constant m . Then there exists $\lambda^* > 0$ such that, for all $\lambda > \lambda^*$, problem (1.1) possesses one least energy sign-changing solution u_0 which has precisely two nodal domains.*

Theorem 1.2. *Suppose that $V \in C(\Omega, \mathbb{R})$, $V(x) \geq m > 0$ for some constant m . Then there exists $\lambda^{**} > 0$ such that, for all $\lambda > \max\{\lambda^*, \lambda^{**}\}$, $c^* := \inf_{u \in \mathcal{N}_b^{\lambda}} I_b^{\lambda}(u) > 0$ is achieved either by a positive or a negative function and $I_b^{\lambda}(u_0) > 2c^*$, where $\mathcal{N}_b^{\lambda} = \{u \in H_V^1(\Omega) \setminus \{0\} | \langle (I_b^{\lambda})'(u), u \rangle = 0\}$ and u_0 is the least energy sign-changing solution obtained in Theorem 1.1.*

2. Framework

In this section, we introduce the variational framework associated with problem (1.1). We first describe the working space. Let $L^p(\Omega)$ be the usual Lebesgue space with the norm $\|u\|_p = \left(\int_{\Omega} |u|^p dx\right)^{1/p}$ and $H_0^1(\Omega)$ be the completion of $C_0^{\infty}(\Omega)$ with respect to the norm:

$$\|u\|_{H_0^1(\Omega)}^2 = \int_{\Omega} (|\nabla u|^2 + u^2) dx.$$

Moreover, we denote the completion of $C_0^\infty(\Omega)$ with respect to the norm:

$$\|u\|_1^2 := \|u\|_{D_0^{1,2}(\Omega)}^2 = \int_{\Omega} |\nabla u|^2 dx.$$

In order to use perturbation method, we will use the space

$$E = W^{1,4}(\Omega) \cap H_V^1(\Omega),$$

where

$$H_V^1(\Omega) = \left\{ u \in H_0^1(\Omega) : \int_{\Omega} V(x)u^2 dx < \infty \right\},$$

which is a Hilbert space endowed with the norm:

$$\|u\|^2 = \int_{\Omega} (a|\nabla u|^2 + V(x)u^2) dx,$$

and $W^{1,4}(\Omega)$ endowed with the norm:

$$\|u\|_W := \left(\int_{\Omega} |\nabla u|^4 + u^4 dx \right)^{\frac{1}{4}}.$$

Moreover, according to Hölder inequality

$$\int_{\Omega} |\nabla u|^2 u^2 dx \leq \left(\int_{\Omega} |\nabla u|^4 dx \right)^{\frac{1}{2}} \left(\int_{\Omega} |u|^4 dx \right)^{\frac{1}{2}} \leq \|u\|_W^4.$$

The norm of E is denoted by

$$\|u\|_E = \left(\|u\|_W^2 + \|u\|^2 \right)^{\frac{1}{2}}.$$

Since $I_{b,\mu}^\lambda(u) = I_b^\lambda(u) + \frac{\mu}{4} \int_{\Omega} (|\nabla u|^4 + u^4) dx$, we can easily get $I_{b,\mu}^\lambda \in C^1(E, \mathbb{R})$ for all $\varphi \in E$ and

$$\begin{aligned} \langle (I_{b,\mu}^\lambda)'(u), \varphi \rangle &= \mu \int_{\Omega} (|\nabla u|^2 \nabla u \nabla \varphi + |u|^2 u \varphi) dx + \int_{\Omega} a \nabla u \nabla \varphi dx + b \int_{\Omega} |\nabla u|^2 dx \int_{\Omega} \nabla u \nabla \varphi dx \\ &+ \int_{\Omega} V(x) u \varphi dx + \int_{\Omega} \phi_u u \varphi dx + \int_{\Omega} (|\nabla u|^2 u \varphi + |u|^2 \nabla u \nabla \varphi) dx \\ &- \lambda \int_{\Omega} |u|^{q-2} u \varphi \ln |u|^2 dx - \int_{\Omega} |u|^4 u \varphi dx. \end{aligned}$$

It is noticed that if $u^\pm \neq 0$, we have

$$\begin{aligned} I_{b,\mu}^\lambda &= I_{b,\mu}^\lambda(u^+) + I_{b,\mu}^\lambda(u^-) + \frac{b}{2} \|u^+\|_1^2 \|u^-\|_1^2 + \frac{1}{4} \int_{\Omega} \phi_{u^-} (u^+)^2 dx + \frac{1}{4} \int_{\Omega} \phi_{u^+} (u^-)^2 dx, \\ \langle (I_{b,\mu}^\lambda)'(u), u^+ \rangle &= \langle (I_{b,\mu}^\lambda)'(u^+), u^+ \rangle + b \|u^+\|_1^2 \|u^-\|_1^2 + \int_{\Omega} \phi_{u^-} (u^+)^2 dx, \\ \langle (I_{b,\mu}^\lambda)'(u), u^- \rangle &= \langle (I_{b,\mu}^\lambda)'(u^-), u^- \rangle + b \|u^+\|_1^2 \|u^-\|_1^2 + \int_{\Omega} \phi_{u^+} (u^-)^2 dx. \end{aligned}$$

Our goal in this paper is to seek the least energy sign-changing solutions of problem (1.1).

Now, fixed $u \in E$ with $u^\pm \neq 0$, we denote $\psi_u : [0, \infty) \times [0, \infty) \rightarrow \mathbb{R}$ and mapping $T_u : [0, \infty) \times [0, \infty) \rightarrow \mathbb{R}^2$ by

$$\psi_u(\alpha, \beta) = I_{b,\mu}^\lambda(\alpha u^+ + \beta u^-), \quad (2.1)$$

and

$$T_u(\alpha, \beta) = (\langle (I_{b,\mu}^\lambda)'(\alpha u^+ + \beta u^-), \alpha u^+ \rangle, \langle (I_{b,\mu}^\lambda)'(\alpha u^+ + \beta u^-), \beta u^- \rangle). \quad (2.2)$$

At last of this section, we give some properties of ϕ_u .

Lemma 2.1. ([25]) For any $u \in H_0^1(\Omega)$, we have

(1) there exist $C > 0$ such that

$$\int_{\Omega} \phi_u u^2 dx \leq C \|u\|_1^4, \quad \forall u \in H_0^1(\Omega);$$

(2) $\phi_u > 0$, $\forall u \in H_0^1(\Omega)$;

(3) $\phi_{\tau u} = \tau^2 \phi_u$, $\forall \tau > 0$ and $u \in H_0^1(\Omega)$;

(4) If $u_n \rightharpoonup u$ in $H_0^1(\Omega)$, then $\phi_{u_n} \rightharpoonup \phi_u$ in $D_0^{1,2}(\Omega)$.

3. Some technical lemmas

In this section, we give some useful lemmas as which are critical to the proof of Theorem 1.1.

Lemma 3.1. For any $u \in E$ with $u^\pm \neq 0$, then there exists a unique maximum point pair (α_u, β_u) of the function ψ_u such that $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\mu}^1$.

Proof. Our proof will be divided into three steps.

Step 1: For any $u \in E$ with $u^\pm \neq 0$, in the following, we will prove the existence of α_u and β_u . From sample computation, we have

$$\lim_{\tau \rightarrow 0} \frac{|\tau|^{q-1} \ln |\tau|^2}{|\tau|} = 0 \quad \text{and} \quad \lim_{\tau \rightarrow \infty} \frac{|\tau|^{q-1} \ln |\tau|^2}{|\tau|^{r-1}} = 0 \quad (3.1)$$

for all $r \in (q, 6)$. Then for any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that

$$|\tau|^{q-1} \ln |\tau|^2 \leq \varepsilon |\tau| + C_\varepsilon |\tau|^{r-1}. \quad (3.2)$$

Since $4 < q < 6$, it follows from (3.2) and the Sobolev embedding theorem that

$$\begin{aligned}
& \langle (I_{b,\mu}^\lambda)'(\alpha u^+ + \beta u^-), \alpha u^+ \rangle \\
&= \mu \alpha^4 \|u^+\|_W^4 + \alpha^2 \|u^+\|^2 + b \alpha^4 \|u^+\|_1^4 + b \alpha^2 \beta^2 \|u^+\|_1^2 \|u^-\|_1^2 + 2\alpha^2 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx \\
&\quad + \alpha^4 \int_\Omega \phi_{u^+}(u^+)^2 dx + \alpha^2 \beta^2 \int_\Omega \phi_{u^-}(u^+)^2 dx - \lambda \int_\Omega |\alpha u^+|^q \ln |\alpha u^+|^2 dx - \alpha^6 \int_\Omega |u^+|^6 dx \\
&\geq \alpha^2 \|u^+\|^2 + b \alpha^4 \|u^+\|_1^4 + b \alpha^2 \beta^2 \|u^+\|_1^2 \|u^-\|_1^2 + 2\alpha^2 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx \\
&\quad + \alpha^4 \int_\Omega \phi_{u^+}(u^+)^2 dx + \alpha^2 \beta^2 \int_\Omega \phi_{u^-}(u^+)^2 dx - \lambda \alpha^2 \varepsilon \int_\Omega |u^+|^2 dx \\
&\quad - \lambda C_\varepsilon \alpha^r \int_\Omega |u^+|^r dx - \alpha^6 \int_\Omega |u^+|^6 dx \\
&\geq \alpha^2 \|u^+\|^2 + b \alpha^4 \|u^+\|_1^4 - \lambda \alpha^2 \varepsilon C_1 \|u^+\|^2 - \lambda C_\varepsilon \alpha^r C_2 \|u^+\|^r - C_3 \alpha^6 \|u^+\|^6 \\
&= (1 - \lambda \varepsilon C_1) \alpha^2 \|u^+\|^2 + b \alpha^4 \|u^+\|_1^4 - \lambda C_\varepsilon \alpha^r C_2 \|u^+\|^r - C_3 \alpha^6 \|u^+\|^6,
\end{aligned}$$

where C_1, C_2, C_3 are positive constants. Choosing $\varepsilon > 0$ such that $1 - \lambda \varepsilon C_1 > 0$. Since $4 < r < 6$, we have $\langle (I_{b,\mu}^\lambda)'(\alpha u^+ + \beta u^-), \alpha u^+ \rangle > 0$ for α small enough and all $\beta > 0$.

Similarly, we obtain that $\langle (I_{b,\mu}^\lambda)'(\alpha u^+ + \beta u^-), \beta u^- \rangle > 0$ for β small enough and all $\alpha > 0$.

Therefore, there exists $\alpha_1 > 0$ such that

$$\langle (I_{b,\mu}^\lambda)'(\alpha_1 u^+ + \beta u^-), \alpha_1 u^+ \rangle > 0, \quad \langle (I_{b,\mu}^\lambda)'(\alpha u^+ + \alpha_1 u^-), \alpha_1 u^- \rangle > 0 \quad (3.3)$$

for all $\alpha, \beta > 0$.

On the other hand, since $u^+ \neq 0$, there exists a constant $\theta > 0$ such that $\text{meas}\{x \in \Omega, u^+ > \theta\} > 0$. Since $q > 4$, we deduce that, for any $M > 1$, there exists $T > 0$ such that $\frac{|\tau|^q \ln |\tau|^2}{\tau^4} > M$ for all $\tau > T$. Therefore, for $\alpha > \frac{T}{\theta}$, we have

$$\lambda \int_\Omega |\alpha u^+|^q \ln |\alpha u^+|^2 dx \geq M \alpha^4 \int_{\{u^+ > \theta\}} (u^+)^4 dx.$$

We can choose $\alpha = \alpha_2^* > \alpha_1$, if $\beta \in [\alpha_1, \alpha_2^*]$ and α_2^* is large enough, it follows that

$$\begin{aligned}
& \langle (I_{b,\mu}^\lambda)'(\alpha_2^* u^+ + \beta u^-), \alpha_2^* u^+ \rangle \\
&\leq \mu (\alpha_2^*)^4 \|u^+\|_W^4 + (\alpha_2^*)^2 \|u^+\|^2 + b (\alpha_2^*)^4 \|u^+\|_1^4 + b (\alpha_2^*)^2 \beta^2 \|u^+\|_1^2 \|u^-\|_1^2 \\
&\quad + 2(\alpha_2^*)^4 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx + (\alpha_2^*)^4 \int_\Omega \phi_{u^+}(u^+)^2 dx \\
&\quad + (\alpha_2^*)^2 \beta^2 \int_\Omega \phi_{u^-}(u^+)^2 dx - M (\alpha_2^*)^4 \int_{\{u^+ > \theta\}} (u^+)^4 dx - (\alpha_2^*)^6 \int_\Omega |u^+|^6 dx \\
&\leq 0.
\end{aligned}$$

Similarly, we get

$$\langle (I_{b,\mu}^\lambda)'(\alpha u^+ + \alpha_2^* u^-), \alpha_2^* u^- \rangle \leq 0.$$

Let $\alpha_2 > \alpha_2^*$ be large enough, we obtain that, for all $\alpha, \beta \in [\alpha_1, \alpha_2]$, we have

$$\langle (I_{b,\mu}^\lambda)'(\alpha_2 u^+ + \beta u^-), \alpha_2 u^+ \rangle < 0, \quad \langle (I_{b,\mu}^\lambda)'(\alpha u^+ + \alpha_2 u^-), \alpha_2 u^- \rangle < 0. \quad (3.4)$$

Combining (3.3) and (3.4) with Miranda's theorem, there exist $(\alpha_u, \beta_u) \in (0, +\infty) \times (0, +\infty)$ such that $T_u(\alpha_u, \beta_u) = (0, 0)$, i.e., $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\mu}^\lambda$.

Step 2: In this step, we will prove the uniqueness of the pair (α_u, β_u) .

Case 1: $u \in \mathcal{M}_{b,\mu}^\lambda$.

If $u \in \mathcal{M}_{b,\mu}^\lambda$, we have

$$\begin{aligned} & \mu \|u^+\|_W^4 + \|u^+\|^2 + b \|u^+\|_1^4 + b \|u^+\|_1^2 \|u^-\|_1^2 + 2 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx \\ & + \int_\Omega \phi_{u^+}(u^+)^2 dx + \int_\Omega \phi_{u^-}(u^+)^2 dx \\ & = \lambda \int_\Omega |u^+|^q \ln |u^+|^2 dx + \int_\Omega |u^+|^6 dx \end{aligned} \quad (3.5)$$

and

$$\begin{aligned} & \mu \|u^-\|_W^4 + \|u^-\|^2 + b \|u^-\|_1^4 + b \|u^+\|_1^2 \|u^-\|_1^2 + 2 \int_\Omega |\nabla u^-|^2 |u^-|^2 dx \\ & + \int_\Omega \phi_{u^-}(u^-)^2 dx + \int_\Omega \phi_{u^+}(u^-)^2 dx \\ & = \lambda \int_\Omega |u^-|^q \ln |u^-|^2 dx + \int_\Omega |u^-|^6 dx. \end{aligned} \quad (3.6)$$

In the following we show that $(\alpha_u, \beta_u) = (1, 1)$.

Let (α_u, β_u) be a pair of numbers such that $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\mu}^\lambda$ with $0 < \alpha_u \leq \beta_u$. Hence, one has that

$$\begin{aligned} & (\alpha_u)^4 \mu \|u^+\|_W^4 + (\alpha_u)^2 \|u^+\|^2 + b (\alpha_u)^4 \|u^+\|_1^4 + b (\alpha_u)^2 (\beta_u)^2 \|u^+\|_1^2 \|u^-\|_1^2 \\ & + 2 (\alpha_u)^4 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx + (\alpha_u)^4 \int_\Omega \phi_{u^+}(u^+)^2 dx + (\alpha_u)^2 (\beta_u)^2 \int_\Omega \phi_{u^-}(u^+)^2 dx \\ & = \lambda \int_\Omega |\alpha_u u^+|^q \ln |\alpha_u u^+|^2 dx + \int_\Omega |\alpha_u u^+|^6 dx \end{aligned} \quad (3.7)$$

and

$$\begin{aligned} & (\beta_u)^4 \mu \|u^-\|_W^4 + (\beta_u)^2 \|u^-\|^2 + b (\beta_u)^4 \|u^-\|_1^4 + b (\alpha_u)^2 (\beta_u)^2 \|u^+\|_1^2 \|u^-\|_1^2 \\ & + 2 (\beta_u)^4 \int_\Omega |\nabla u^-|^2 |u^-|^2 dx + (\beta_u)^4 \int_\Omega \phi_{u^-}(u^-)^2 dx + (\alpha_u)^2 (\beta_u)^2 \int_\Omega \phi_{u^+}(u^-)^2 dx \\ & = \lambda \int_\Omega |\beta_u u^-|^q \ln |\beta_u u^-|^2 dx + \int_\Omega |\beta_u u^-|^6 dx. \end{aligned} \quad (3.8)$$

According to $0 < \alpha_u \leq \beta_u$ and (3.8), we have that

$$\begin{aligned} & \mu \|u^-\|_W^4 + \frac{\|u^-\|^2}{\beta_u^2} + b \|u^-\|_1^4 + b \|u^+\|_1^2 \|u^-\|_1^2 \\ & + 2 \int_{\Omega} |\nabla u^-|^2 |u^-|^2 dx + \int_{\Omega} \phi_{u^-} (u^-)^2 dx + \int_{\Omega} \phi_{u^+} (u^-)^2 dx \\ & \geq \lambda \int_{\Omega} \frac{|\beta_u u^-|^q \ln |\beta_u u^-|^2}{(\beta_u)^4} dx + (\beta_u)^2 \int_{\Omega} |u^-|^6 dx. \end{aligned} \quad (3.9)$$

If $\beta_u > 1$, by (3.6) and (3.9), one has that

$$\begin{aligned} & \left(\frac{1}{(\beta_u)^2} - 1 \right) \|u^-\|^2 \\ & \geq \lambda \int_{\Omega} \left[\frac{|\beta_u u^-|^q \ln |\beta_u u^-|^2}{(\beta_u)^4} - |u^-|^q \ln |u^-|^2 \right] dx + ((\beta_u)^2 - 1) \int_{\Omega} |u^-|^6 dx. \end{aligned}$$

The left side of above inequality is negative, which is a contradiction because the right side is positive. Therefore, we conclude that $0 < \alpha_u \leq \beta_u \leq 1$.

Similarly, by (3.5), (3.7) and $0 < \alpha_u \leq \beta_u$, we have that

$$\begin{aligned} & \left(\frac{1}{(\alpha_u)^2} - 1 \right) \|u^+\|^2 \\ & \leq \lambda \int_{\Omega} \left[\frac{|\alpha_u u^+|^q \ln |\alpha_u u^+|^2}{(\alpha_u)^4} - |u^+|^q \ln |u^+|^2 \right] dx + ((\alpha_u)^2 - 1) \int_{\Omega} |u^+|^6 dx. \end{aligned}$$

This fact implies that $\alpha_u \geq 1$. Consequently, $\alpha_u = \beta_u = 1$.

Case 2: $u \notin \mathcal{M}_{b,\mu}^\lambda$.

Suppose that there exists $(\tilde{\alpha}_1, \tilde{\beta}_1), (\tilde{\alpha}_2, \tilde{\beta}_2)$ such that

$$u_1 = \tilde{\alpha}_1 u^+ + \tilde{\beta}_1 u^- \in \mathcal{M}_{b,\mu}^\lambda \quad \text{and} \quad u_2 = \tilde{\alpha}_2 u^+ + \tilde{\beta}_2 u^- \in \mathcal{M}_{b,\mu}^\lambda.$$

Hence

$$u_2 = \left(\frac{\tilde{\alpha}_2}{\tilde{\alpha}_1} \right) \tilde{\alpha}_1 u^+ + \left(\frac{\tilde{\beta}_2}{\tilde{\beta}_1} \right) \tilde{\beta}_1 u^- = \left(\frac{\tilde{\alpha}_2}{\tilde{\alpha}_1} \right) u_1^+ + \left(\frac{\tilde{\beta}_2}{\tilde{\beta}_1} \right) u_1^- \in \mathcal{M}_{b,\mu}^\lambda.$$

By $u_1 \in \mathcal{M}_{b,\mu}^\lambda$, one has that

$$\frac{\tilde{\alpha}_2}{\tilde{\alpha}_1} = \frac{\tilde{\beta}_2}{\tilde{\beta}_1} = 1.$$

Hence, $\tilde{\alpha}_1 = \tilde{\alpha}_2, \tilde{\beta}_1 = \tilde{\beta}_2$.

Step 3: In this step we will prove that (α_u, β_u) is the unique maximum point of ψ_u on $[0, +\infty) \times [0, +\infty)$.

Firstly, it is easy to see that

$$2\rho^q - q\rho^q \ln |\rho|^2 \leq 2 \quad \text{for all } \rho \in (0, \infty). \quad (3.10)$$

Let $\Omega^+ = \{x \in \Omega : u(x) > 0\}$ and $\Omega^- = \{x \in \Omega : u(x) < 0\}$, $u \in H$ with $u^\pm \neq 0$, we have

$$\int_{\Omega} |\alpha u^+ + \beta u^-|^q \ln |\alpha u^+ + \beta u^-|^2 dx = \int_{\Omega} (|\alpha u^+|^q \ln |\alpha u^+|^2 + |\beta u^-|^q \ln |\beta u^-|^2) dx. \quad (3.11)$$

Combining (3.10) and (3.11), we get

$$\begin{aligned}
c &= I_{b,\mu}^{\lambda}(\alpha u^+ + \beta u^-) \\
&= \frac{\mu\alpha^4}{4}\|u^+\|_W^4 + \frac{\mu\beta^4}{4}\|u^-\|_W^4 + \frac{\alpha^2}{2}\|u^+\|^2 + \frac{\beta^2}{2}\|u^-\|^2 + \frac{\alpha^4}{4}b\|u^+\|_1^4 + \frac{\beta^4}{4}b\|u^-\|_1^4 \\
&\quad + \frac{\alpha^2\beta^2}{2}b\|u^+\|_1^2\|u^-\|_1^2 + \frac{\alpha^4}{2}\int_{\Omega}|\nabla u^+|^2|u^+|^2dx + \frac{\beta^4}{2}\int_{\Omega}|\nabla u^-|^2|u^-|^2dx \\
&\quad + \frac{\alpha^4}{4}\int_{\Omega}\phi_{u^+}(u^+)^2dx + \frac{\alpha^2\beta^2}{4}\int_{\Omega}\phi_{u^-}(u^+)^2dx + \frac{\alpha^2\beta^2}{4}\int_{\Omega}\phi_{u^+}(u^-)^2dx + \frac{\beta^4}{4}\int_{\Omega}\phi_{u^-}(u^-)^2dx \\
&\quad + \frac{\lambda}{q^2}\int_{\Omega}(2|\alpha u^+|^q - q|\alpha u^+|^q \ln|\alpha u^+|^2)dx + \frac{\lambda}{q^2}\int_{\Omega}(2|\beta u^-|^q - q|\beta u^-|^q \ln|\beta u^-|^2)dx \\
&\quad - \frac{\alpha^6}{6}\int_{\Omega}|u^+|^6dx - \frac{\beta^6}{6}\int_{\Omega}|u^-|^6dx \\
&\leq \frac{\mu\alpha^4}{4}\|u^+\|_W^4 + \frac{\mu\beta^4}{4}\|u^-\|_W^4 + \frac{\alpha^2}{2}\|u^+\|^2 + \frac{\beta^2}{2}\|u^-\|^2 + \frac{\alpha^4}{4}b\|u^+\|_1^4 + \frac{\beta^4}{4}b\|u^-\|_1^4 \\
&\quad + \frac{\alpha^2\beta^2}{2}b\|u^+\|_1^2\|u^-\|_1^2 + \frac{\alpha^4}{2}\int_{\Omega}|\nabla u^+|^2|u^+|^2dx + \frac{\beta^4}{2}\int_{\Omega}|\nabla u^-|^2|u^-|^2dx \\
&\quad + \frac{\alpha^4}{4}\int_{\Omega}\phi_{u^+}(u^+)^2dx + \frac{\alpha^2\beta^2}{4}\int_{\Omega}\phi_{u^-}(u^+)^2dx + \frac{\alpha^2\beta^2}{4}\int_{\Omega}\phi_{u^+}(u^-)^2dx + \frac{\beta^4}{4}\int_{\Omega}\phi_{u^-}(u^-)^2dx \\
&\quad + \frac{4\lambda}{q^2}|\Omega| - \frac{\alpha^6}{6}\int_{\Omega}|u^+|^6dx - \frac{\beta^6}{6}\int_{\Omega}|u^-|^6dx,
\end{aligned}$$

which implies that $\lim_{(\alpha,\beta)\rightarrow\infty}\psi_u(\alpha,\beta) = -\infty$. So it is sufficient to check that a maximum point can not be achieved on the boundary of $[0, +\infty) \times [0, +\infty)$. By contradiction, we suppose that $(0, \beta_u)$ is a maximum point of $\psi_u(\alpha, \beta)$ with $\beta_u \geq 0$. Then, we have

$$\begin{aligned}
&\psi_u(\alpha, \beta_u) \\
&= \frac{\mu\alpha^4}{4}\|u^+\|_W^4 + \frac{\mu(\beta_u)^4}{4}\|u^-\|_W^4 + \frac{\alpha^2}{2}\|u^+\|^2 + \frac{(\beta_u)^2}{2}\|u^-\|^2 + \frac{\alpha^4}{4}b\|u^+\|_1^4 + \frac{(\beta_u)^4}{4}b\|u^-\|_1^4 \\
&\quad + \frac{\alpha^2(\beta_u)^2}{2}b\|u^+\|_1^2\|u^-\|_1^2 + \frac{\alpha^4}{2}\int_{\Omega}|\nabla u^+|^2|u^+|^2dx + \frac{(\beta_u)^4}{2}\int_{\Omega}|\nabla u^-|^2|u^-|^2dx \\
&\quad + \frac{\alpha^4}{4}\int_{\Omega}\phi_{u^+}(u^+)^2dx + \frac{(\beta_u)^4}{4}\int_{\Omega}\phi_{u^-}(u^-)^2dx + \frac{\alpha^2(\beta_u)^2}{4}\int_{\Omega}\phi_{u^-}(u^+)^2dx \\
&\quad + \frac{\alpha^2(\beta_u)^2}{4}\int_{\Omega}\phi_{u^+}(u^-)^2dx + \frac{\lambda}{q^2}\int_{\Omega}(2|\alpha u^+|^q - q|\alpha u^+|^q \ln|\alpha u^+|^2)dx \\
&\quad + \frac{\lambda}{q^2}\int_{\Omega}(2|\beta_u u^-|^q - q|\beta_u u^-|^q \ln|\beta_u u^-|^2)dx - \frac{\alpha^6}{6}\int_{\Omega}|u^+|^6dx - \frac{(\beta_u)^6}{6}\int_{\Omega}|u^-|^6dx.
\end{aligned}$$

Therefore, it is obvious that

$$\begin{aligned}
 (\psi_u(\alpha, \beta_u))'_\alpha &= \mu\alpha^3 \|u^+\|_W^4 + \alpha \|u^+\|^2 + b\alpha^3 \|u^+\|_1^4 + b\alpha\beta_u \|u^+\|_1^2 \|u^-\|_1^2 + 2\alpha^3 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx \\
 &\quad + \alpha^3 \int_\Omega \phi_{u^+}(u^+)^2 dx + \frac{\alpha(\beta_u)^2}{2} \int_\Omega \phi_{u^-}(u^+)^2 dx + \frac{\alpha(\beta_u)^2}{2} \int_\Omega \phi_{u^+}(u^-)^2 dx \\
 &\quad - \alpha^{q-1} \lambda \int_\Omega |u^+|^q \ln |\alpha u^+|^2 dx - \alpha^5 \int_\Omega |u^+|^6 dx \\
 &> 0,
 \end{aligned}$$

if α is small enough, that is, ψ_u is an increasing function with respect to α if α is small enough. This yields the contradiction. Similarly, ψ_u can not achieve its global maximum point at $(\alpha_u, 0)$ with $\alpha_u > 0$. \square

Lemma 3.2. For any $u \in E$ with $u^\pm \neq 0$, such that $\langle (I_{b,\mu}^\lambda)'(u), u^\pm \rangle \leq 0$, then the unique maximum point of ψ_u in $[0, +\infty) \times [0, +\infty)$ satisfies $0 < \alpha_u, \beta_u \leq 1$.

Proof. If $\alpha_u = 0$ or $\beta_u = 0$, according to Lemma 3.1, ψ_u can not achieve maximum. Without loss of generality, let $\alpha_u \geq \beta_u > 0$. On the one hand, by $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\mu}^\lambda$, we have

$$\begin{aligned}
 &(\alpha_u)^4 \mu \|u^+\|_W^4 + (\alpha_u)^2 \|u^+\|^2 + b(\alpha_u)^4 \|u^+\|_1^4 + b(\alpha_u)^2 (\beta_u)^2 \|u^+\|_1^2 \|u^-\|_1^2 \\
 &\quad + 2(\alpha_u)^4 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx + (\alpha_u)^4 \int_\Omega \phi_{u^+}(u^+)^2 dx + (\alpha_u)^2 (\beta_u)^2 \int_\Omega \phi_{u^-}(u^+)^2 dx \\
 &= \lambda \int_\Omega |\alpha_u u^+|^q \ln |\alpha_u u^+|^2 dx + (\alpha_u)^6 \int_\Omega |u^+|^6 dx.
 \end{aligned} \tag{3.12}$$

On the other hand, by $\langle (I_{b,\mu}^\lambda)'(u), u^\pm \rangle \leq 0$, we obtain

$$\begin{aligned}
 &\mu \|u^+\|_W^4 + \|u^+\|^2 + b \|u^+\|_1^4 + b \|u^+\|_1^2 \|u^-\|_1^2 \\
 &\quad + 2 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx + \int_\Omega \phi_{u^+}(u^+)^2 dx + \int_\Omega \phi_{u^-}(u^+)^2 dx \\
 &\leq \lambda \int_\Omega |u^+|^q \ln |u^+|^2 dx + \int_\Omega |u^+|^6 dx.
 \end{aligned} \tag{3.13}$$

So, it follows from (3.12) and (3.13), we get

$$\begin{aligned}
 &\left(\frac{1}{(\alpha_u)^2} - 1\right) \|u^+\|^2 \\
 &\geq \lambda \int_\Omega [(\alpha_u)^{q-4} |u^+|^q \ln |\alpha_u u^+|^2 - |u^+|^q \ln |u^+|^2] dx + ((\alpha_u)^2 - 1) \int_\Omega |u^+|^6 dx.
 \end{aligned} \tag{3.14}$$

Since $q > 4$, we conclude that $0 < \beta_u < \alpha_u \leq 1$, so $0 < \alpha_u, \beta_u \leq 1$. \square

Lemma 3.3. Let $c_{b,\mu}^\lambda = \inf_{u \in \mathcal{M}_b^\lambda} I_{b,\mu}^\lambda(u)$, then we get $\lim_{\lambda \rightarrow \infty} c_{b,\mu}^\lambda = 0$.

Proof. For any $u \in \mathcal{M}_{b,\mu}^\lambda$,

$$\begin{aligned} & \mu \|u^\pm\|_W^4 + \|u^\pm\|^2 + b \|u^\pm\|_1^4 + b \|u^\pm\|_1^2 \|u^\pm\|_1^2 \\ & + 2 \int_\Omega |\nabla u^\pm|^2 |u^\pm|^2 dx + \int_\Omega \phi_{u^\pm}(u^\pm)^2 dx + \int_\Omega \phi_{u^\mp}(u^\mp)^2 dx \\ & = \lambda \int_\Omega |u^\pm|^q \ln |u^\pm|^2 dx + \int_\Omega |u^\pm|^6 dx. \end{aligned}$$

Then by (3.2) and the Sobolev inequalities, we get

$$\|u^\pm\|^2 \leq \lambda \int_\Omega |u^\pm|^q \ln |u^\pm|^2 dx + \int_\Omega |u^\pm|^6 dx \leq \lambda \varepsilon C_1 \|u^\pm\|^2 + \lambda C_\varepsilon C_2 \|u^\pm\|^r + C_3 \|u^\pm\|^6.$$

Thus

$$(1 - \lambda \varepsilon C_1) \|u^\pm\|^2 \leq C_2 \|u^\pm\|^r + C_3 \|u^\pm\|^6.$$

Choosing ε small enough such that $1 - \lambda \varepsilon C_1 > 0$, since $r > 4$, there exists $\rho > 0$ such that

$$\|u^\pm\|^2 \geq \rho \text{ for all } u \in \mathcal{M}_{b,\mu}^\lambda. \quad (3.15)$$

Thanks to $u \in \mathcal{M}_{b,\mu}^\lambda$, we have $\langle (I_{b,\mu}^\lambda)'(u), u \rangle = 0$. Then

$$\begin{aligned} I_{b,\mu}^\lambda(u) &= I_{b,\mu}^\lambda(u) - \frac{1}{q} \langle (I_{b,\mu}^\lambda)'(u), u \rangle \\ &= \mu \left(\frac{1}{4} - \frac{1}{q}\right) \|u\|_W^4 + \left(\frac{1}{2} - \frac{1}{q}\right) \|u\|^2 + \left(\frac{1}{4} - \frac{1}{q}\right) b \|u\|_1^4 + 2 \left(\frac{1}{4} - \frac{1}{q}\right) \int_\Omega |\nabla u|^2 |u|^2 dx \\ &\quad + \left(\frac{1}{4} - \frac{1}{q}\right) \int_\Omega \phi_u u^2 dx + \frac{2\lambda}{q^2} \int_\Omega |u|^q dx + \left(\frac{1}{q} - \frac{1}{6}\right) \int_\Omega |u|^6 dx \\ &\geq \left(\frac{1}{2} - \frac{1}{q}\right) \|u\|^2, \end{aligned}$$

thus $I_{b,\mu}^\lambda(u)$ is bounded below on $\mathcal{M}_{b,\mu}^\lambda$.

For any $u \in E$ with $u^\pm \neq 0$, by using Lemma 3.1, for each $\lambda > 0$, there exist $\{\alpha_\lambda, \beta_\lambda\}$ such that $\alpha_\lambda u^+ + \beta_\lambda u^- \in \mathcal{M}_{b,\mu}^\lambda$, we have

$$\begin{aligned} 0 &\leq c_b^\lambda = \inf I_{b,\mu}^\lambda(u) \leq I_{b,\mu}^\lambda(\alpha_\lambda u^+ + \beta_\lambda u^-) \\ &\leq \frac{\mu}{4} \|\alpha_\lambda u^+ + \beta_\lambda u^-\|_W^4 + \frac{1}{2} \|\alpha_\lambda u^+ + \beta_\lambda u^-\|^2 + \frac{b}{4} \|\alpha_\lambda u^+ + \beta_\lambda u^-\|_1^4 + \frac{2\lambda}{q^2} \int_\Omega |\alpha_\lambda u^+ + \beta_\lambda u^-|^q dx \\ &\quad + \frac{1}{2} \int_\Omega |\nabla(\alpha_\lambda u^+ + \beta_\lambda u^-)|^2 |\alpha_\lambda u^+ + \beta_\lambda u^-|^2 dx + \frac{1}{4} \int_\Omega \phi_{\alpha_\lambda u^+ + \beta_\lambda u^-} (\alpha_\lambda u^+ + \beta_\lambda u^-)^2 dx \\ &\leq \frac{\mu}{2} (\alpha_\lambda)^4 \|u^+\|_W^4 + \frac{\mu}{2} (\beta_\lambda)^4 \|u^-\|_W^4 + (\alpha_\lambda)^2 \|u^+\|^2 + (\beta_\lambda)^2 \|u^-\|^2 + 2b(\alpha_\lambda)^4 \|u^+\|_1^4 \\ &\quad + 2b(\beta_\lambda)^4 \|u^-\|_1^4 + (\alpha_\lambda)^4 \int_\Omega |\nabla u^+|^2 |u^+|^2 dx + (\beta_\lambda)^4 \int_\Omega |\nabla u^-|^2 |u^-|^2 dx \\ &\quad + 2C(\alpha_\lambda)^4 \|u^+\|_1^4 + 2C(\beta_\lambda)^4 \|u^-\|_1^4 + \frac{2\lambda}{q^2} \int_\Omega |\alpha_\lambda u^+|^q dx + \frac{2\lambda}{q^2} \int_\Omega |\beta_\lambda u^-|^q dx. \end{aligned}$$

Next we will prove that $\alpha_\lambda \rightarrow 0$ and $\beta_\lambda \rightarrow 0$ as $\lambda \rightarrow \infty$.

Let $G_u = \{(\alpha_\lambda, \beta_\lambda) \in [0, +\infty) \times [0, +\infty) : T_u(\alpha_\lambda, \beta_\lambda) = (0, 0), \lambda > 0\}$, we can calculate that

$$\begin{aligned} & (\alpha_\lambda)^6 \int_{\Omega} |u^+|^6 dx + (\beta_\lambda)^6 \int_{\Omega} |u^-|^6 dx \\ & + \lambda(\alpha_\lambda)^q \int_{\Omega} |u^+|^q \ln |\alpha_\lambda u^+|^2 dx + \lambda(\beta_\lambda)^q \int_{\Omega} |u^-|^q \ln |\beta_\lambda u^-|^2 dx \\ & = \mu \|\alpha_\lambda u^+ + \beta_\lambda u^-\|_W^4 + \|\alpha_\lambda u^+ + \beta_\lambda u^-\|^2 + b \|\alpha_\lambda u^+ + \beta_\lambda u^-\|_1^4 + 2(\alpha_\lambda)^4 \int_{\Omega} |\nabla u^+|^2 |u^+|^2 dx \\ & + 2(\beta_\lambda)^4 \int_{\Omega} |\nabla u^-|^2 |u^-|^2 dx + \int_{\Omega} \phi_{\alpha_\lambda u^+ + \beta_\lambda u^-} (\alpha_\lambda u^+ + \beta_\lambda u^-)^2 dx \\ & \leq 2\mu(\alpha_\lambda)^4 \|u^+\|_W^4 + 2\mu(\beta_\lambda)^4 \|u^-\|_W^4 + 2(\alpha_\lambda)^2 \|u^+\|^2 \\ & + 2(\beta_\lambda)^2 \|u^-\|^2 + 4b(\alpha_\lambda)^4 \|u^+\|_1^4 + 4b(\beta_\lambda)^4 \|u^-\|_1^4 \\ & + 2(\alpha_\lambda)^4 \int_{\Omega} |\nabla u^+|^2 |u^+|^2 dx + 2(\beta_\lambda)^4 \int_{\Omega} |\nabla u^-|^2 |u^-|^2 dx \\ & + 2C(\alpha_\lambda)^4 \|u^+\|_1^4 + 2C(\beta_\lambda)^4 \|u^-\|_1^4. \end{aligned}$$

Hence G_u is bounded. Let $\{\lambda_n\} \subset (0, \infty)$ such that $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$. Then there exist α_0 and β_0 such that $(\alpha_{\lambda_n}, \beta_{\lambda_n}) \rightarrow (\alpha_0, \beta_0)$ as $n \rightarrow \infty$.

Now, we claim $\alpha_0 = \beta_0 = 0$. Suppose, by contradiction, if $\alpha_0 > 0$ or $\beta_0 > 0$, by $\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^- \in \mathcal{M}_{b,\mu}^{\lambda_n}$, for any $n \in \mathbb{N}$, we have

$$\begin{aligned} & \mu \|\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-\|_W^4 + \|\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-\|^2 + b \|\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-\|_1^4 \\ & + \int_{\Omega} |\nabla(\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-)|^2 |\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-|^2 dx + \int_{\Omega} \phi_{\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-} (\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-)^2 dx. \quad (3.16) \\ & = \lambda_n \int_{\Omega} |\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-|^q \ln |\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-|^2 dx + \int_{\Omega} |\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-|^6 dx. \end{aligned}$$

Thanks to $\alpha_{\lambda_n} u^+ \rightarrow \alpha_0 u^+$ and $\beta_{\lambda_n} u^- \rightarrow \beta_0 u^-$ in E , (3.2) and the Lebesgue dominated convergence theorem, we get

$$\int_{\Omega} |\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-|^q \ln |\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-|^2 dx \rightarrow \int_{\Omega} |\alpha_0 u^+ + \beta_0 u^-|^q \ln |\alpha_0 u^+ + \beta_0 u^-|^2 dx > 0$$

as $n \rightarrow \infty$. It follows from $\lambda_n \rightarrow \infty$ and the boundness of $\{\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-\}$ in E that we have a contradiction with equality (3.16). Hence, $\alpha_0 = \beta_0 = 0$, we conclude that $\lim_{\lambda \rightarrow \infty} c_{b,\mu}^\lambda = 0$. \square

Lemma 3.4. There exists $\lambda^* > 0$ such that for all $\lambda \geq \lambda^*$, the infimum $c_{b,\mu}^\lambda$ is achieved.

Proof. By the definition of $c_{b,\mu}^\lambda = \inf_{u \in \mathcal{M}_{b,\mu}^\lambda} I_{b,\mu}^\lambda(u)$, there exists a sequence $\{u_n\} \subset \mathcal{M}_{b,\mu}^\lambda$ such that

$$\lim_{\lambda \rightarrow \infty} I_{b,\mu}^\lambda(u_n) = c_{b,\mu}^\lambda.$$

Obviously, $\{u_n\}$ is bounded in E . Then, up to subsequence, still denoted by $\{u_n\}$, there exists $u \in E$ such that $u_n \rightharpoonup u$. Since the embedding $E \hookrightarrow L^p(\Omega)$ is compact for all $p \in [2, 6)$, we have

$$u_n \rightarrow u \text{ in } L^p(\Omega) \text{ and } u_n \rightarrow u \text{ a.e. } x \in \Omega.$$

Hence

$$\begin{aligned} u_n^\pm &\rightharpoonup u^\pm \text{ in } E, \\ u_n^\pm &\rightarrow u^\pm \text{ in } L^p(\Omega), \\ u_n^\pm &\rightarrow u^\pm \text{ in a.e. } x \in \Omega. \end{aligned}$$

By Lemma 3.1, we have

$$I_{b,\mu}^\lambda(\alpha u_n^+ + \beta u_n^-) \leq I_{b,\mu}^\lambda(u_n)$$

for all $\alpha, \beta \geq 0$. On the one hand, the Vitali convergence theorem yields that

$$\lim_{n \rightarrow \infty} \int_{\Omega} |u_n|^q \ln |u_n|^2 dx \rightarrow \int_{\Omega} |u|^q \ln |u|^2 dx. \quad (3.17)$$

Then, by (3.17), Brézis-Lieb lemma [28] and the weak semicontinuity of norm, we get

$$\begin{aligned} &\liminf_{n \rightarrow \infty} I_{b,\mu}^\lambda(\alpha u_n^+ + \beta u_n^-) \\ &\geq \mu \frac{\alpha^4}{4} \lim_{n \rightarrow \infty} \left(\|u_n^+ - u^+\|_W^4 + \|u^+\|_W^4 \right) + \mu \frac{\beta^4}{4} \lim_{n \rightarrow \infty} \left(\|u_n^- - u^-\|_W^4 + \|u^-\|_W^4 \right) \\ &\quad + \frac{\alpha^2}{2} \lim_{n \rightarrow \infty} (\|u_n^+ - u^+\|^2 + \|u^+\|^2) + \frac{\beta^2}{2} (\|u_n^- - u^-\|^2 + \|u^-\|^2) \\ &\quad + \frac{b\alpha^4}{4} \left(\lim_{n \rightarrow \infty} \|u_n^+ - u^+\|_1^2 + \|u^+\|_1^2 \right)^2 + \frac{b\beta^4}{4} \left(\lim_{n \rightarrow \infty} \|u_n^- - u^-\|_1^2 + \|u^-\|_1^2 \right)^2 \\ &\quad + \frac{\alpha^4}{2} \liminf_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n^+|^2 |u_n^+|^2 dx + \frac{\beta^4}{2} \liminf_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n^-|^2 |u_n^-|^2 dx + \frac{\alpha^4}{4} \lim_{n \rightarrow \infty} \int_{\Omega} \phi_{u_n^+}(u_n^+)^2 dx \\ &\quad + \frac{\alpha^2 \beta^2}{4} \lim_{n \rightarrow \infty} \int_{\Omega} \phi_{u_n^+}(u_n^+)^2 dx + \frac{\alpha^2 \beta^2}{4} \lim_{n \rightarrow \infty} \int_{\Omega} \phi_{u_n^+}(u_n^-)^2 dx + \frac{\beta^4}{4} \lim_{n \rightarrow \infty} \int_{\Omega} \phi_{u_n^-}(u_n^-)^2 dx \\ &\quad - \frac{\alpha^6}{6} \left(\lim_{n \rightarrow \infty} \int_{\Omega} |u_n^+ - u^+|^6 dx + \int_{\Omega} |u^+|^6 dx \right) - \frac{\beta^6}{6} \left(\lim_{n \rightarrow \infty} \int_{\Omega} |u_n^- - u^-|^6 dx + \int_{\Omega} |u^-|^6 dx \right) \\ &\quad + \frac{2\lambda}{q^2} \int_{\Omega} |\alpha u_n^+ + \beta u_n^-|^q dx - \frac{\lambda}{q} \int_{\Omega} |\alpha u_n^+ + \beta u_n^-|^q \ln |\alpha u_n^+ + \beta u_n^-|^2 dx \\ &\geq I_{b,\mu}^\lambda(\alpha u^+ + \beta u^-) + \frac{\mu\alpha^4}{4} A_1 + \frac{\mu\beta^4}{4} A_2 + \frac{\alpha^2}{2} A_3 + \frac{b\alpha^4}{2} A_5 \|u^+\|_1^2 + \frac{b\alpha^4}{4} A_5^2 - \frac{\alpha^6}{6} B_1 \\ &\quad + \frac{\beta^2}{2} A_4 + \frac{\beta^4}{2} A_6^2 \|u^-\|_1^2 + \frac{b\beta^4}{4} A_6^2 - \frac{\beta^6}{6} B_2, \end{aligned}$$

where

$$\begin{aligned} A_1 &= \lim_{n \rightarrow \infty} \|u_n^+ - u^+\|_W^4, A_2 = \lim_{n \rightarrow \infty} \|u_n^- - u^-\|_W^4, A_3 = \lim_{n \rightarrow \infty} \|u_n^+ - u^+\|^2, A_4 = \lim_{n \rightarrow \infty} \|u_n^- - u^-\|^2, \\ A_5 &= \lim_{n \rightarrow \infty} \|u_n^+ - u^+\|_1^2, A_6 = \lim_{n \rightarrow \infty} \|u_n^- - u^-\|_1^2, B_1 = \lim_{n \rightarrow \infty} |u_n^+ - u^+|_6^6, B_2 = \lim_{n \rightarrow \infty} |u_n^- - u^-|_6^6 \end{aligned}$$

for all $\alpha \geq 0$ and $\beta \geq 0$. So,

$$\begin{aligned} c_{b,\mu}^\lambda &\geq I_{b,\mu}^\lambda(\alpha u^+ + \beta u^-) + \frac{\mu\alpha^4}{4} A_1 + \frac{\mu\beta^4}{4} A_2 + \frac{\alpha^2}{2} A_3 + \frac{b\alpha^4}{2} A_5 \|u^+\|_1^2 + \frac{b\alpha^4}{4} A_5^2 \\ &\quad - \frac{\alpha^6}{6} B_1 + \frac{\beta^2}{2} A_4 + \frac{\beta^4}{2} A_6^2 \|u^-\|_1^2 + \frac{b\beta^4}{4} A_6^2 - \frac{\beta^6}{6} B_2. \end{aligned} \quad (3.18)$$

Denote $\sigma := \frac{1}{3}S^{\frac{3}{2}}$, where $S = \inf_{u \in H^1 \setminus \{0\}} \frac{\|u\|^2}{(\int_{\Omega} |u|^6 dx)^{\frac{1}{3}}}$. According to Lemma 3.3, there is $\lambda^* > 0$ such that

$c_{b,\mu}^{\lambda} < \sigma$ for all $\lambda \geq \lambda^*$.

Step 1: we prove that $u^{\pm} \neq 0$. By contradiction, we suppose $u^+ = 0$ ($u^- = 0$ is similar).

Case 1: $B_1 = 0$. If $A_1 = A_3 = 0$, that is, $u_n^+ \rightarrow u^+$ in E . According to (3.15), we obtain $\|u^+\| > 0$, which contradicts $u^+ = 0$. If $A_1 > 0$, $A_3 > 0$, by (3.18), we get $\frac{\alpha^2}{2}A_3 < c_{b,\mu}^{\lambda}$ for all $\alpha \geq 0$, which is a contradiction.

Case 2: $B_1 > 0$. According to the definition of S , we get $\sigma := \frac{1}{3}S^{\frac{3}{2}} \leq \frac{1}{3}(\frac{A_3}{(B_1)^{\frac{1}{3}}})^{\frac{3}{2}}$, by direct calculation, we obtain

$$\frac{1}{3}\left(\frac{A_3}{(B_1)^{\frac{1}{3}}}\right)^{\frac{3}{2}} = \max_{\alpha \geq 0} \left\{ \frac{\alpha^2}{2}A_3 - \frac{\alpha^6}{6}B_1 \right\} \leq \max_{\alpha \geq 0} \left\{ \frac{\mu\alpha^4}{4}A_1 + \frac{\alpha^2}{2}A_3 + \frac{b\alpha^4}{2}A_5\|u^+\|_1^2 + \frac{b\alpha^4}{4}A_5^2 - \frac{\alpha^6}{6}B_1 \right\}.$$

Since $c_{b,\mu}^{\lambda} \rightarrow 0$ as $\lambda \rightarrow \infty$, there exists $\lambda^* > 0, C > 0$ such that for all $\lambda > \lambda^*, c_{b,\mu}^{\lambda} \leq C$. Then, without loss of generality, we can assume $c_{b,\mu}^{\lambda} < \sigma$, choose $\beta = 0$, it follows from (3.18) that

$$\sigma \leq \max_{\alpha \geq 0} \left\{ \frac{\alpha^2}{2}A_3 - \frac{\alpha^6}{6}B_1 \right\} \leq \max_{\alpha \geq 0} \left\{ \frac{\mu\alpha^4}{4}A_1 + \frac{\alpha^2}{2}A_3 + \frac{b\alpha^4}{2}A_5\|u^+\|_1^2 + \frac{b\alpha^4}{4}A_5^2 - \frac{\alpha^6}{6}B_1 \right\} < \sigma.$$

It is a contradiction, then we obtain $u^+ \neq 0$. Similarly, we obtain $u^- \neq 0$.

Step 2: we prove that $B_1 = 0, B_2 = 0$. We just prove $B_1 = 0$. By contradiction, we suppose that $B_1 > 0$.

Case 1: $B_2 > 0$. Let $\widehat{\alpha}_1$ and $\widehat{\beta}_1$ satisfy

$$\begin{aligned} & \frac{\mu(\widehat{\alpha}_1)^4}{4}A_1 + \frac{(\widehat{\alpha}_1)^2}{2}A_3 + \frac{b(\widehat{\alpha}_1)^4}{2}A_5\|u^+\|_1^2 + \frac{b(\widehat{\alpha}_1)^4}{4}A_5^2 - \frac{(\widehat{\alpha}_1)^6}{6}B_1 \\ &= \max_{\alpha \geq 0} \left\{ \frac{\mu\alpha^4}{4}A_1 + \frac{\alpha^2}{2}A_3 + \frac{b\alpha^4}{2}A_5\|u^+\|_1^2 + \frac{b\alpha^4}{4}A_5^2 - \frac{\alpha^6}{6}B_1 \right\}, \end{aligned}$$

and

$$\begin{aligned} & \frac{\mu(\widehat{\beta}_1)^4}{4}A_2 + \frac{(\widehat{\beta}_1)^2}{2}A_4 + \frac{b(\widehat{\beta}_1)^4}{2}A_6\|u^-\|_1^2 + \frac{b(\widehat{\beta}_1)^4}{4}A_6^2 - \frac{(\widehat{\beta}_1)^6}{6}B_2 \\ &= \max_{\beta \geq 0} \left\{ \frac{\mu\beta^4}{4}A_2 + \frac{\beta^2}{2}A_4 + \frac{b\beta^4}{2}A_6\|u^-\|_1^2 + \frac{b\beta^4}{4}A_6^2 - \frac{\beta^6}{6}B_2 \right\}. \end{aligned}$$

According to $[0, \widehat{\alpha}_1] \times [0, \widehat{\beta}_1]$ is compact, there exists $(\alpha_u, \beta_u) \in [0, \widehat{\alpha}_1] \times [0, \widehat{\beta}_1]$ such that $\psi_u(\alpha_u, \beta_u) = \max_{(\alpha, \beta) \in [0, \widehat{\alpha}_1] \times [0, \widehat{\beta}_1]} \psi_u(\alpha, \beta)$.

In the following, we will prove $(\alpha_u, \beta_u) \in (0, \widehat{\alpha}_1) \times (0, \widehat{\beta}_1)$. Obviously, if β small enough, we have

$$\psi_u(\alpha, 0) < I_{b,\mu}^{\lambda}(\alpha u^+) + I_{b,\mu}^{\lambda}(\beta u^-) \leq I_{b,\mu}^{\lambda}(\alpha u^+ + \beta u^-) = \psi_u(\alpha, \beta), \quad \forall \alpha \in [0, \widehat{\alpha}_1].$$

Hence, there exists $\beta_0 \in [0, \widehat{\beta}_1]$ such that $\psi_u(\alpha, 0) < \psi_u(\alpha, \beta_0)$ for all $\alpha \in [0, \widehat{\alpha}_1]$. That is, $(\alpha_u, \beta_u) \notin [0, \widehat{\alpha}_1] \times \{0\}$. By similar discussion, we conclude that $(\alpha_u, \beta_u) \notin \{0\} \times [0, \widehat{\beta}_1]$.

Obviously, we get

$$\frac{\mu\alpha^4}{4}A_1 + \frac{\alpha^2}{2}A_3 + \frac{b\alpha^4}{2}A_5\|u^+\|_1^2 + \frac{b\alpha^4}{4}A_5^2 - \frac{\alpha^6}{6}B_1 > 0, \quad \alpha \in (0, \widehat{\alpha}_1), \quad (3.19)$$

$$\frac{\mu\beta^4}{4}A_2 + \frac{\beta^2}{2}A_4 + \frac{b\beta^4}{2}A_6\|u^-\|_1^2 + \frac{b\beta^4}{4}A_6^2 - \frac{\beta^6}{6}B_2 > 0, \beta \in (0, \widehat{\beta}_1]. \quad (3.20)$$

Then, for all $\alpha \in (0, \widehat{\alpha}_1]$ and $\beta \in (0, \widehat{\beta}_1]$, we get

$$\begin{aligned} \sigma &\leq \frac{\mu(\widehat{\alpha}_1)^4}{4}A_1 + \frac{(\widehat{\alpha}_1)^2}{2}A_3 + \frac{b(\widehat{\alpha}_1)^4}{2}A_5\|u^+\|_1^2 + \frac{b(\widehat{\alpha}_1)^4}{4}A_5^2 - \frac{(\widehat{\alpha}_1)^6}{6}B_1 \\ &\quad + \frac{\mu\beta^4}{4}A_2 + \frac{\beta^2}{2}A_4 + \frac{b\beta^4}{2}A_6\|u^-\|_1^2 + \frac{b\beta^4}{4}A_6^2 - \frac{\beta^6}{6}B_2 \end{aligned}$$

and

$$\begin{aligned} \sigma &\leq \frac{\mu\alpha^4}{4}A_1 + \frac{\alpha^2}{2}A_3 + \frac{b\alpha^4}{2}A_5\|u^+\|_1^2 + \frac{b\alpha^4}{4}A_5^2 - \frac{\alpha^6}{6}B_1 \\ &\quad + \frac{\mu(\widehat{\beta}_1)^4}{4}A_2 + \frac{\widehat{\beta}_1^2}{2}A_4 + \frac{b(\widehat{\beta}_1)^4}{2}A_6\|u^-\|_1^2 + \frac{b(\widehat{\beta}_1)^4}{4}A_6^2 - \frac{(\widehat{\beta}_1)^6}{6}B_2. \end{aligned}$$

Together with (3.18), we obtain $\psi_u(\alpha, \widehat{\beta}_1) \leq 0$, $\psi_u(\widehat{\alpha}_1, \beta) \leq 0$ for all $\alpha \in [0, \widehat{\alpha}_1]$ and $\beta \in [0, \widehat{\beta}_1]$. That is, $(\alpha_u, \beta_u) \notin [0, \widehat{\alpha}_1] \times \{\widehat{\beta}_1\}$ and $(\alpha_u, \beta_u) \notin \{0, \widehat{\alpha}_1\} \times [0, \widehat{\beta}_1]$.

In conclusion, we get $(\alpha_u, \beta_u) \in (0, \widehat{\alpha}_1) \times (0, \widehat{\beta}_1)$. Hence, $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\mu}^\lambda$. So, combining (3.18), (3.19) with (3.20), we have that

$$\begin{aligned} c_{b,\mu}^\lambda &\geq I_{b,\mu}^\lambda(\alpha_u u^+ + \beta_u u^-) + \frac{\mu(\alpha_u)^4}{4}A_1 + \frac{(\alpha_u)^2}{2}A_3 + \frac{b(\alpha_u)^4}{2}A_5\|u^+\|_1^2 + \frac{b(\alpha_u)^4}{4}A_5^2 \\ &\quad - \frac{(\alpha_u)^6}{6}B_1 + \frac{\mu(\beta_u)^4}{4}A_2 + \frac{\beta_u^2}{2}A_4 + \frac{b(\beta_u)^4}{2}A_6\|u^-\|_1^2 + \frac{b(\beta_u)^4}{4}A_6^2 - \frac{(\beta_u)^6}{6}B_2 \\ &> I_{b,\mu}^\lambda(\alpha_u u^+ + \beta_u u^-) \geq c_{b,\mu}^\lambda. \end{aligned}$$

Therefore, we have a contradiction.

Case 2: $B_2 = 0$. In this case, we can maximize in $(0, \widehat{\alpha}_1) \times (0, \infty)$. Indeed, it is possible to show that there exists $\beta_0 \in [0, \infty]$ such that $I_{b,\mu}^\lambda(\alpha u^+ + \beta u^-) < 0$ for all $(\alpha, \beta) \in [0, \widehat{\alpha}_1] \times [\beta_0, \infty)$. Hence, there exists $(\alpha_u, \beta_u) \in [0, \widehat{\alpha}_1] \times [0, \infty)$ such that

$$\psi_u(\alpha_u, \beta_u) = \max_{\alpha \in [0, \widehat{\alpha}_1] \times [0, \infty)} \psi_u(\alpha, \beta).$$

Following, we prove that $(\alpha_u, \beta_u) \in [0, \widehat{\alpha}_1] \times [0, \infty)$.

Since $\psi_u(\alpha, 0) \leq \psi_u(\alpha, \beta)$ for $\alpha \in [0, \widehat{\alpha}_1]$ and β is small enough, we have $(\alpha_u, \beta_u) \notin \{0\} \times [0, \infty)$. On the other hand, for all $\beta \in [0, \infty)$, it is obvious that

$$\begin{aligned} \sigma &\leq \frac{\mu(\widehat{\alpha}_1)^4}{4}A_1 + \frac{(\widehat{\alpha}_1)^2}{2}A_3 + \frac{b(\widehat{\alpha}_1)^4}{2}A_5\|u^+\|_1^2 + \frac{b(\widehat{\alpha}_1)^4}{4}A_5^2 - \frac{(\widehat{\alpha}_1)^6}{6}B_1 \\ &\quad + \frac{\mu\beta^4}{4}A_2 + \frac{\beta^2}{2}A_4 + \frac{b\beta^4}{2}A_6\|u^-\|_1^2 + \frac{b\beta^4}{4}A_6^2. \end{aligned}$$

Hence, we have that $\psi_u(\widehat{\alpha}_1, \beta) \leq 0$ for all $\beta \in [0, \infty)$. Thus, $(\alpha_u, \beta_u) \notin \{\widehat{\alpha}_1\} \times [0, \infty)$, and so $(\alpha_u, \beta_u) \in$

$[0, \widehat{\alpha}_1] \times [0, \infty)$. That is, $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\mu}^\lambda$, therefore, according to (3.19), we have

$$\begin{aligned} c_{b,\mu}^\lambda &\geq I_{b,\mu}^\lambda(\alpha_u u^+ + \beta_u u^-) + \frac{\mu(\alpha_u)^4}{4} A_1 + \frac{(\alpha_u)^2}{2} A_3 + \frac{b(\alpha_u)^4}{2} A_5 \|u^+\|_1^2 + \frac{b(\alpha_u)^4}{4} A_5^2 \\ &\quad - \frac{(\alpha_u)^6}{6} B_1 + \frac{\mu(\beta_u)^4}{4} A_2 + \frac{(\beta_u)^2}{2} A_4 + \frac{b(\beta_u)^4}{2} A_6 \|u^-\|_1^2 + \frac{b(\beta_u)^4}{4} A_6^2 \\ &> I_{b,\mu}^\lambda(\alpha_u u^+ + \beta_u u^-) \geq c_{b,\mu}^\lambda, \end{aligned}$$

which is a contradiction.

Therefore, from above discussion, we have $B_1 = B_2 = 0$.

Lastly, we prove that $c_{b,\mu}^\lambda$ is achieved.

Since $u^\pm \neq 0$, by Lemma 3.1, there exist $\alpha_u, \beta_u > 0$ such that

$$\widetilde{u} = \alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\mu}^\lambda.$$

Furthermore, the norm in E is lower semicontinuous, it is easy to see that

$$\langle (I_{b,\mu}^\lambda)'(u), u^\pm \rangle \leq 0.$$

By Lemma 3.2, we obtain $\alpha_u, \beta_u \leq 1$.

From $u_n \in \mathcal{M}_{b,\mu}^\lambda$, according to Lemma 3.1, we get

$$I_{b,\mu}^\lambda(\alpha_u u_n^+ + \beta_u u_n^-) \leq I_{b,\mu}^\lambda(u_n^+ + u_n^-) = I_{b,\mu}^\lambda(u_n).$$

Thanks to $B_1 = B_2 = 0$, we obtain

$$\begin{aligned} c_{b,\mu}^\lambda &\leq I_{b,\mu}^\lambda(\widetilde{u}) - \frac{1}{q} \langle (I_{b,\mu}^\lambda)'(\widetilde{u}), \widetilde{u} \rangle \\ &= \left(\frac{1}{4} - \frac{1}{q}\right) \|\widetilde{u}\|_W^4 + \left(\frac{1}{2} - \frac{1}{q}\right) \|\widetilde{u}\|^2 + \left(\frac{1}{4} - \frac{1}{q}\right) \|\widetilde{u}\|_1^4 + \left(\frac{1}{2} - \frac{1}{q}\right) \int_\Omega |\nabla \widetilde{u}|^2 |\widetilde{u}|^2 dx \\ &\quad + \left(\frac{1}{4} - \frac{1}{q}\right) \int_\Omega \phi_{\widetilde{u}} \widetilde{u}^2 dx + \frac{2\lambda}{q^2} \int_\Omega |\widetilde{u}|^q dx + \left(\frac{1}{q} - \frac{1}{6}\right) \int_\Omega |\widetilde{u}|^6 dx \\ &= \left(\frac{1}{4} - \frac{1}{q}\right) \|\alpha_u u^+ + \beta_u u^-\|_W^4 + \left(\frac{1}{2} - \frac{1}{q}\right) \|\alpha_u u^+ + \beta_u u^-\|^2 + \left(\frac{1}{4} - \frac{1}{q}\right) \|\alpha_u u^+ + \beta_u u^-\|_1^4 \\ &\quad + \left(\frac{1}{4} - \frac{1}{q}\right) (\alpha_u)^4 \int_\Omega \phi_{u^+} (u^+)^2 dx + \left(\frac{1}{4} - \frac{1}{q}\right) (\alpha_u)^2 (\beta_u)^2 \int_\Omega \phi_{u^-} (u^+)^2 dx \\ &\quad + \left(\frac{1}{4} - \frac{1}{q}\right) (\alpha_u)^2 (\beta_u)^2 \int_\Omega \phi_{u^+} (u^-)^2 dx + \left(\frac{1}{4} - \frac{1}{q}\right) (\beta_u)^4 \int_\Omega \phi_{u^-} (u^-)^2 dx \\ &\quad + \frac{2\lambda}{q^2} \left[\int_\Omega |\alpha_u u^+|^q dx + \int_\Omega |\beta_u u^-|^q dx \right] + \left(\frac{1}{q} - \frac{1}{6}\right) \left[\int_\Omega |\alpha_u u^+|^6 dx + \int_\Omega |\beta_u u^-|^6 dx \right] \\ &\leq \left(\frac{1}{4} - \frac{1}{q}\right) \|u\|_W^4 + \left(\frac{1}{2} - \frac{1}{q}\right) \|u\|^2 + \left(\frac{1}{4} - \frac{1}{q}\right) \|u\|_1^4 + \left(\frac{1}{2} - \frac{1}{q}\right) \int_\Omega |\nabla u|^2 |u|^2 dx \\ &\quad + \left(\frac{1}{4} - \frac{1}{q}\right) \int_\Omega \phi_u' u^2 dx + \frac{2\lambda}{q^2} \int_\Omega |u|^q dx + \left(\frac{1}{q} - \frac{1}{6}\right) \int_\Omega |u|^6 dx \\ &\leq \liminf_{n \rightarrow \infty} \left[I_{b,\mu}^\lambda(u_n) - \frac{1}{q} \langle (I_{b,\mu}^\lambda)'(u_n), u_n \rangle \right]. \end{aligned}$$

Therefore, $\alpha_u = \beta_u = 1$ and $c_{b,\mu}^\lambda$ is achieved by $u_{b,\mu} = u^+ + u^- \in \mathcal{M}_{b,\mu}^\lambda$. \square

4. Proof of Theorems

In order to obtain a sign-changing solution of problem (1.1), we firstly prove that $u_{b,\mu}$ is a sign-changing critical point of $I_{b,\mu}^\lambda$.

Lemma 4.1. *If $u_{b,\mu} \in \mathcal{M}_{b,\mu}^\lambda$ and $I_{b,\mu}^\lambda(u_{b,\mu}) = c_{b,\mu}^\lambda$ for $\mu \in (0, 1]$, then $u_{b,\mu}$ is a sign-changing critical point of $I_{b,\mu}^\lambda$. Moreover, $u_{b,\mu}$ has exactly two nodal domains.*

Proof. Since $u_{b,\mu} \in \mathcal{M}_{b,\mu}^\lambda$, for $(\alpha, \beta) \in (\mathbb{R}^+ \times \mathbb{R}^+) \setminus (1, 1)$, we have

$$I_{b,\mu}^\lambda(\alpha u_{b,\mu}^+ + \beta u_{b,\mu}^-) < I_{b,\mu}^\lambda(u_{b,\mu}^+ + u_{b,\mu}^-) = c_{b,\mu}^\lambda. \quad (4.1)$$

Arguing by contradiction, we assume that $(I_{b,\mu}^\lambda)'(u_{b,\mu}) \neq 0$, then there exist $\delta > 0$ and $\tau > 0$ such that

$$\|(I_{b,\mu}^\lambda)'(v)\| > \tau \text{ for all } \|v - u_{b,\mu}\| \geq 3\delta.$$

Choose $\tau \in (0, \min\{\frac{1}{2}, \frac{\delta}{\sqrt{2}\|u_{b,\mu}\|}\})$, let

$$D = (1 - \tau, 1 + \tau) \times (1 - \tau, 1 + \tau)$$

and

$$g(\alpha, \beta) = \alpha u_{b,\mu}^+ + \beta u_{b,\mu}^- \text{ for all } (\alpha, \beta) \in D.$$

According to (4.1), we have that

$$c_\lambda := \max_{\partial\Omega} (I_{b,\mu}^\lambda) \circ g < c_{b,\mu}^\lambda. \quad (4.2)$$

Let $\varepsilon := \min\{c_{b,\mu}^\lambda - c_\lambda, \frac{\tau\varepsilon}{8}\}$ and $S_\delta = B(u_{b,\mu}, \delta)$, according to Lemma 2.3 in [28], there exists a deformation $\eta \in C([0, 1] \times D, D)$ such that

$$(a) \quad \eta(1, v) = v \text{ if } v \notin (I_{b,\mu}^\lambda)^{-1}([c_{b,\mu}^\lambda - 2\varepsilon, c_{b,\mu}^\lambda + 2\varepsilon] \cap S_{2\delta}),$$

$$(b) \quad \eta(1, (I_{b,\mu}^\lambda)^{c_{b,\mu}^\lambda + \varepsilon} \cap S_{2\delta}) \subset (I_{b,\mu}^\lambda)^{c_{b,\mu}^\lambda - \varepsilon},$$

$$(c) \quad I_{b,\mu}^\lambda(\eta(1, v)) \leq I_{b,\mu}^\lambda(v) \text{ for all } v \in H.$$

Firstly, from (b) and Lemma 3.1, it is easy to see that

$$I_{b,\mu}^\lambda(g(\alpha, \beta)) < c_{b,\mu}^\lambda < c_{b,\mu}^\lambda + \varepsilon.$$

That is,

$$g(\alpha, \beta) \in (I_{b,\mu}^\lambda)^{c_{b,\mu}^\lambda + \varepsilon}.$$

On the other hand, we have

$$\begin{aligned} \|g(\alpha, \beta) - u_{b,\mu}\|^2 &= \|(\alpha - 1)u_{b,\mu}^+ + (\beta - 1)u_{b,\mu}^-\|^2 \\ &\leq 2[(\alpha - 1)^2\|u_{b,\mu}^+\|^2 + (\beta - 1)^2\|u_{b,\mu}^-\|^2] \\ &\leq 2\tau^2\|u_{b,\mu}\|^2 \leq \delta^2. \end{aligned}$$

Hence, by (b), we have

$$I_{b,\mu}^\lambda(\eta(1, g(\alpha, \beta))) \leq c_{b,\mu}^\lambda - \varepsilon < c_{b,\mu}^\lambda. \quad (4.3)$$

Next, we prove that $\eta(1, g(D)) \cap \mathcal{M}_{b,\mu}^\lambda \neq \emptyset$, which contradicts the definition of $c_{b,\mu}^\lambda$. Let $\gamma(\alpha, \beta) := \eta(1, g(\alpha, \beta))$ and

$$\begin{aligned} \Psi_0(\alpha, \beta) &:= \left(\langle (I_{b,\mu}^\lambda)'(g(\alpha, \beta)), u_{b,\mu}^+ \rangle, \langle (I_{b,\mu}^\lambda)'(g(\alpha, \beta)), u_{b,\mu}^- \rangle \right) \\ &= \left(\langle (I_{b,\mu}^\lambda)'(\alpha u_{b,\mu}^+ + \beta u_{b,\mu}^-), u_{b,\mu}^+ \rangle, \langle (I_{b,\mu}^\lambda)'(\alpha u_{b,\mu}^+ + \beta u_{b,\mu}^-), u_{b,\mu}^- \rangle \right) \\ &= (\varphi_u^1(\alpha, \beta), \varphi_u^2(\alpha, \beta)) \end{aligned}$$

and

$$\Psi_1(\alpha, \beta) := \left(\frac{1}{\alpha} \langle (I_{b,\mu}^\lambda)'(\gamma(\alpha, \beta)), (\gamma(\alpha, \beta))^+ \rangle, \frac{1}{\beta} \langle (I_{b,\mu}^\lambda)'(\gamma(\alpha, \beta)), (\gamma(\alpha, \beta))^- \rangle \right).$$

Since $u_{b,\mu} \in \mathcal{M}_{b,\mu}^\lambda$, by the direct calculation, we have

$$\begin{aligned} \left. \frac{\partial \varphi_u^1(\alpha, \beta)}{\partial \alpha} \right|_{(1,1)} &= 3\|u_{b,\mu}^+\|_W^4 + \|u_{b,\mu}^+\|^2 + 3b\|u_{b,\mu}^+\|_1^4 + b\|u_{b,\mu}^+\|_1^2 \|u_{b,\mu}^-\|_1^2 \\ &\quad + 2 \int_\Omega |\nabla u_{b,\mu}^+|^2 |u_{b,\mu}^+|^2 dx + 3 \int_\Omega \phi_{u_{b,\mu}^+}(u_{b,\mu}^+)^2 dx + \int_\Omega \phi_{u_{b,\mu}^-}(u_{b,\mu}^+)^2 dx \\ &\quad - \lambda(q-1) \int_\Omega |u_{b,\mu}^+|^q \ln |u_{b,\mu}^+|^2 dx - 2\lambda \int_\Omega |u_{b,\mu}^+|^q dx - 5 \int_\Omega |u_{b,\mu}^+|^6 dx \\ &= (4-q)\|u_{b,\mu}^+\|_W^4 + (2-q)\|u_{b,\mu}^+\|^2 + b(4-q)\|u_{b,\mu}^+\|_1^4 + (2-q)b\|u_{b,\mu}^+\|_1^2 \|u_{b,\mu}^-\|_1^2 \\ &\quad + (2-q) \int_\Omega \phi_{u_{b,\mu}^-}(u_{b,\mu}^+)^2 dx + (4-q) \int_\Omega \phi_{u_{b,\mu}^+}(u_{b,\mu}^+)^2 dx \\ &\quad + (4-2q) \int_\Omega |\nabla u_{b,\mu}^+|^2 |u_{b,\mu}^+|^2 dx - 2\lambda \int_\Omega |u_{b,\mu}^+|^q dx - (6-q) \int_\Omega |u_{b,\mu}^+|^6 dx, \end{aligned}$$

and

$$\left. \frac{\partial \varphi_u^1(\alpha, \beta)}{\partial \beta} \right|_{(1,1)} = 2b\|u_{b,\mu}^+\|_1^2 \|u_{b,\mu}^-\|_1^2 + 2 \int_\Omega \phi_{u_{b,\mu}^-}(u_{b,\mu}^+)^2 dx.$$

Similarly,

$$\begin{aligned} \left. \frac{\partial \varphi_u^2(\alpha, \beta)}{\partial \beta} \right|_{(1,1)} &= 3\|u_{b,\mu}^-\|_W^4 + \|u_{b,\mu}^-\|^2 + 3b\|u_{b,\mu}^-\|_1^4 + b\|u_{b,\mu}^+\|_1^2 \|u_{b,\mu}^-\|_1^2 \\ &\quad + 2 \int_\Omega |\nabla u_{b,\mu}^-|^2 |u_{b,\mu}^-|^2 dx + 3 \int_\Omega \phi_{u_{b,\mu}^-}(u_{b,\mu}^-)^2 dx + \int_\Omega \phi_{u_{b,\mu}^+}(u_{b,\mu}^-)^2 dx \\ &\quad - \lambda(q-1) \int_\Omega |u_{b,\mu}^-|^q \ln |u_{b,\mu}^-|^2 dx - 2\lambda \int_\Omega |u_{b,\mu}^-|^q dx - 5 \int_\Omega |u_{b,\mu}^-|^6 dx \\ &= (4-q)\|u_{b,\mu}^-\|_W^4 + (2-q)\|u_{b,\mu}^-\|^2 + b(4-q)\|u_{b,\mu}^-\|_1^4 + (2-q)b\|u_{b,\mu}^+\|_1^2 \|u_{b,\mu}^-\|_1^2 \\ &\quad + (2-q) \int_\Omega \phi_{u_{b,\mu}^+}(u_{b,\mu}^-)^2 dx + (4-q) \int_\Omega \phi_{u_{b,\mu}^-}(u_{b,\mu}^-)^2 dx \\ &\quad + (4-2q) \int_\Omega |\nabla u_{b,\mu}^-|^2 |u_{b,\mu}^-|^2 dx - 2\lambda \int_\Omega |u_{b,\mu}^-|^q dx - (6-q) \int_\Omega |u_{b,\mu}^-|^6 dx, \end{aligned}$$

and

$$\frac{\partial \varphi_u^2(\alpha, \beta)}{\partial \alpha} \Big|_{(1,1)} = 2b \|u_{b,\mu}^+\|_1^2 \|u_{b,\mu}^-\|_1^2 + 2 \int_{\Omega} \phi_{u_{b,\mu}^+} (u_{b,\mu}^-)^2 dx.$$

Let

$$M = \begin{bmatrix} \frac{\varphi_u^1(\alpha, \beta)}{\partial \alpha} \Big|_{(1,1)} & \frac{\varphi_u^2(\alpha, \beta)}{\partial \alpha} \Big|_{(1,1)} \\ \frac{\varphi_u^1(\alpha, \beta)}{\partial \beta} \Big|_{(1,1)} & \frac{\varphi_u^2(\alpha, \beta)}{\partial \beta} \Big|_{(1,1)} \end{bmatrix}.$$

Since $q > 4$, then,

$$\det M = \frac{\partial \varphi_u^1(\alpha, \beta)}{\partial \alpha} \Big|_{(1,1)} \times \frac{\partial \varphi_u^2(\alpha, \beta)}{\partial \beta} \Big|_{(1,1)} - \frac{\partial \varphi_u^1(\alpha, \beta)}{\partial \beta} \Big|_{(1,1)} \times \frac{\partial \varphi_u^2(\alpha, \beta)}{\partial \alpha} \Big|_{(1,1)} > 0.$$

Since $\Psi_0(\alpha, \beta)$ is a C^1 function and $(1,1)$ is the unique isolated zero point of Ψ_0 , by using the degree theory, we deduce that $\deg(\Psi_0, D, 0) = 1$.

Hence, combining (4.3) and (a), we obtain

$$g(\alpha, \beta) = \gamma(\alpha, \beta) \text{ on } \partial D.$$

Consequently, we obtain $\deg(\Psi_1, D, 0) = 1$. Therefore, $\Psi_1(\alpha_0, \beta_0) = 0$ for some $(\alpha_0, \beta_0) \in D$ such that

$$\eta(1, g(\alpha_0, \beta_0)) = \gamma(\alpha_0, \beta_0) \in \mathcal{M}_{b,\mu}^\lambda,$$

which is contradicted to (4.3).

Finally, we prove that $u_{b,\mu}$ has exactly two nodal domains. To this end, we assume by contradiction that

$$u_{b,\mu} = u_1 + u_2 + u_3$$

with

$$u_i \neq 0, u_1 > 0, u_2 < 0 \text{ and } \text{suppt}(u_i) \cap \text{suppt}(u_j) = \emptyset, \text{ for } i \neq j, i, j = 1, 2, 3$$

and

$$\langle (I_{b,\mu}^\lambda)'(u_{b,\mu}), u_i \rangle = 0, \text{ for } i = 1, 2, 3.$$

Setting $v := u_1 + u_2$, we see that $v^+ = u_1$ and $v^- = u_2$, i.e., $v^\pm \neq 0$. Then, there exists a unique pair of positive numbers (α_v, β_v) such that

$$\alpha_v u_1 + \beta_v u_2 \in \mathcal{M}_{b,\mu}^\lambda.$$

Hence

$$I_{b,\mu}^\lambda(\alpha_v u_1 + \beta_v u_2) \geq c_{b,\mu}^\lambda.$$

Moreover, using the fact $\langle (I_{b,\mu}^\lambda)'(u_{b,\mu}), u_i \rangle = 0$, we obtain

$$\langle (I_{b,\mu}^\lambda)'(v), v^\pm \rangle = -b \|v^\pm\|_1^2 \|u_3\|_1^2 - \int_{\Omega} \phi_{u_3} (v^\pm)^2 dx \leq 0.$$

From Lemma 3.2, we get

$$(\alpha_v, \beta_v) \in (0, 1] \times (0, 1].$$

On the other hand, we obtain

$$\begin{aligned}
 0 &= \frac{1}{4} \langle (I_{b,\mu}^\lambda)'(u_{b,\mu}), u_3 \rangle \\
 &= \frac{\mu}{4} \|u_3\|_W^4 + \frac{1}{4} \|u_3\|^2 + \frac{b}{4} \|u_1\|_1^2 \|u_3\|_1^2 + \frac{b}{4} \|u_2\|_1^2 \|u_3\|_1^2 + \int_{\Omega} |\nabla u_3|^2 |u_3|^2 dx \\
 &\quad + \frac{b}{4} \|u_3\|_1^4 + \frac{1}{4} \int_{\Omega} \phi_{u_b}(u_3)^2 dx - \frac{1}{4} \int_{\Omega} |u_3|^6 dx - \frac{\lambda}{4} \int_{\Omega} |u_3|^q \ln |u_3|^2 dx \\
 &\leq I_{b,\mu}^\lambda(u_3) + \frac{b}{4} \|u_1\|_1^2 \|u_3\|_1^2 + \frac{b}{4} \|u_2\|_1^2 \|u_3\|_1^2 + \frac{1}{4} \int_{\Omega} \phi_{u_1}(u_3)^2 dx + \frac{1}{4} \int_{\Omega} \phi_{u_2}(u_3)^2 dx.
 \end{aligned}$$

Hence

$$\begin{aligned}
 c_{b,\mu}^\lambda &\leq I_{b,\mu}^\lambda(\alpha_v u_1 + \beta_v u_2) \\
 &= I_{b,\mu}^\lambda(\alpha_v u_1 + \beta_v u_2) - \frac{1}{4} \langle (I_{b,\mu}^\lambda)'(\alpha_v u_1 + \beta_v u_2), \alpha_v u_1 + \beta_v u_2 \rangle \\
 &= \frac{1}{4} (\|\alpha_v u_1\|^2 + \|\beta_v u_2\|^2) + \frac{\lambda}{q^2} \left[\int_{\Omega} |\alpha_v u_1|^q dx + \int_{\Omega} |\beta_v u_2|^q dx \right] \\
 &\quad + \left(\frac{1}{4} - \frac{1}{q} \right) \lambda \left[\int_{\Omega} |\alpha_v u_1|^q \ln |\alpha_v u_1|^2 dx + \int_{\Omega} |\beta_v u_2|^q \ln |\beta_v u_2|^2 dx \right] \\
 &\quad + \frac{1}{12} \left[\int_{\Omega} |\alpha_v u_1|^6 dx + \int_{\Omega} |\beta_v u_2|^6 dx \right] \\
 &\leq \frac{1}{4} (\|u_1\|^2 + \|u_2\|^2) + \frac{\lambda}{q^2} \left[\int_{\Omega} |u_1|^q dx + \int_{\Omega} |u_2|^q dx \right] \\
 &\quad + \left(\frac{1}{4} - \frac{1}{q} \right) \lambda \left[\int_{\Omega} |u_1|^q \ln |u_1|^2 dx + \int_{\Omega} |u_2|^q \ln |u_2|^2 dx \right] \\
 &\quad + \frac{1}{12} \left[\int_{\Omega} |u_1|^6 dx + \int_{\Omega} |u_2|^6 dx \right] \\
 &= I_{b,\mu}^\lambda(u_1 + u_2) - \frac{1}{4} \langle (I_{b,\mu}^\lambda)'(u_1 + u_2), u_1 + u_2 \rangle \\
 &\leq I_{b,\mu}^\lambda(u_1) + I_{b,\mu}^\lambda(u_2) + I_{b,\mu}^\lambda(u_3) + \frac{b}{4} (\|u_2\|_1^2 + \|u_3\|_1^2) \|u_1\|_1^2 \\
 &\quad + \frac{b}{4} (\|u_1\|_1^2 + \|u_3\|_1^2) \|u_2\|_1^2 + \frac{b}{4} (\|u_1\|_1^2 + \|u_2\|_1^2) \|u_3\|_1^2 \\
 &\quad + \frac{1}{4} \int_{\Omega} \phi_{u_1}(u_3)^2 dx + \frac{1}{4} \int_{\Omega} \phi_{u_2}(u_3)^2 dx + \frac{1}{4} \int_{\Omega} \phi_{u_1}(u_2)^2 dx \\
 &\quad + \frac{1}{4} \int_{\Omega} \phi_{u_3}(u_2)^2 dx + \frac{1}{4} \int_{\Omega} \phi_{u_2}(u_1)^2 dx + \frac{1}{4} \int_{\Omega} \phi_{u_3}(u_1)^2 dx \\
 &= I_{b,\mu}^\lambda(u) = c_{b,\mu}^\lambda,
 \end{aligned}$$

which is a contradiction, that is, $u_3 = 0$ and $u_{b,\mu}$ has exactly two nodal domains. \square

Lemma 4.2. Let $\mu_n \rightarrow 0$ and $\{u_{\mu_n}\} \subset E$ be a sequence of critical points of I_{b,μ_n}^λ , and there exists C independent of n such that $(I_{b,\mu_n}^\lambda)'(u_{\mu_n}) = 0$ and $I_{b,\mu_n}^\lambda(u_{\mu_n}) \leq C$. Then up to a subsequence $u_{\mu_n} \rightarrow u_0$ in E as $n \rightarrow \infty$ and u_0 is a critical point of I_b^λ .

Proof. We prove the lemma in three parts:

Claim 1. $\{u_{\mu_n}\}$ is bounded in E .

$$\begin{aligned} C &\geq I_{b,\mu}^\lambda(u_{\mu_n}) - \frac{1}{q} \langle (I_{b,\mu}^\lambda)'(u_{\mu_n}), u_{\mu_n} \rangle \\ &= \mu \left(\frac{1}{4} - \frac{1}{q} \right) \|u_{\mu_n}\|_W^4 + \left(\frac{1}{2} - \frac{1}{q} \right) \|u_{\mu_n}\|^2 + \left(\frac{1}{4} - \frac{1}{q} \right) b \|u_{\mu_n}\|_1^4 + 2 \left(\frac{1}{4} - \frac{1}{q} \right) \int_{\Omega} |\nabla u_{\mu_n}|^2 |u_{\mu_n}|^2 dx \\ &\quad + \left(\frac{1}{4} - \frac{1}{q} \right) \int_{\Omega} \phi_{u_{\mu_n}} u_{\mu_n}^2 dx + \frac{2\lambda}{q^2} \int_{\Omega} |u_{\mu_n}|^q dx + \left(\frac{1}{q} - \frac{1}{6} \right) \int_{\Omega} |u_{\mu_n}|^6 dx \\ &\geq \mu \left(\frac{1}{4} - \frac{1}{q} \right) \|u_{\mu_n}\|_W^4 + \left(\frac{1}{2} - \frac{1}{q} \right) \|u_{\mu_n}\|^2. \end{aligned}$$

This means that $\{u_{\mu_n}\}$ is bounded in E . Then up to a subsequence, we may suppose $u_{\mu_n} \rightharpoonup u_0$ in E as $n \rightarrow \infty$.

Claim 2. $\{u_{\mu_n}\}$ is bounded in L^∞ . We will use the Moser iteration to accomplish this. Because $\{u_{\mu_n}\}$ satisfies the equation $\langle (I_{b,\mu_n}^\lambda)'(u_{\mu_n}), \varphi \rangle = 0$, for any $\varphi \in E$, we have

$$\begin{aligned} &\mu_n \int_{\Omega} \left(|\nabla u_{\mu_n}|^2 \nabla u_{\mu_n} \nabla \varphi + |u_{\mu_n}|^2 u_{\mu_n} \varphi \right) dx + \int_{\Omega} (a + |u_{\mu_n}|^2) \nabla u_{\mu_n} \nabla \varphi + |\nabla u_{\mu_n}|^2 u_{\mu_n} \varphi dx \\ &\quad + b \left(\int_{\Omega} |\nabla u_{\mu_n}|^2 dx \right) \left(\int_{\Omega} \nabla u_{\mu_n} \nabla \varphi dx \right) + \int_{\Omega} V(x) u_{\mu_n} \varphi dx + \int_{\Omega} \phi_{u_{\mu_n}} u_{\mu_n} \varphi dx \\ &= \lambda \int_{\Omega} |u_{\mu_n}|^{q-2} u_{\mu_n} \varphi \ln |u_{\mu_n}|^2 dx + \int_{\Omega} |u_{\mu_n}|^4 u_{\mu_n} \varphi dx. \end{aligned} \quad (4.4)$$

Now for any $T > 0$, we take $\varphi = |u_{\mu_n}^T|^{2k} u_{\mu_n}$ as test functions in (4.4) with $k \geq k_0$ for some $k_0 > 0$, where $|u_{\mu_n}^T| = |u_{\mu_n}|$, if $|u_{\mu_n}| \leq T$; $|u_{\mu_n}^T| = T$; if $|u_{\mu_n}| \geq T$. By (4.4) and the Sobolev embedding, for $\varepsilon \in (0, 1)$, we have

$$\begin{aligned} &\int_{\{|T \leq |u_{\mu_n}|\}} (a + |u_{\mu_n}|^2) |\nabla u_{\mu_n}|^2 |u_{\mu_n}^T|^{2k} dx + (2k + 1) \int_{\{|u_{\mu_n}| \leq T\}} (a + |u_{\mu_n}|^2) |\nabla u_{\mu_n}|^2 |u_{\mu_n}^T|^{2k} dx \\ &\quad + \int_{\Omega} |\nabla u_{\mu_n}|^2 |u_{\mu_n}|^2 |u_{\mu_n}^T|^{2k} dx + \int_{\Omega} V(x) |u_{\mu_n}|^2 |u_{\mu_n}^T|^{2k} dx \\ &\leq \int_{\Omega} |u_{\mu_n}|^6 |u_{\mu_n}^T|^{2k} dx + \lambda \int_{\Omega} |u_{\mu_n}|^q |u_{\mu_n} \ln |u_{\mu_n}|^2 |u_{\mu_n}^T|^{2k} dx. \end{aligned}$$

Then there exists $C > 0$ such that

$$\begin{aligned} &\int_{\Omega} |u_{\mu_n}|^6 |u_{\mu_n}^T|^{2k} dx + \lambda \int_{\Omega} |u_{\mu_n}|^q |u_{\mu_n} \ln |u_{\mu_n}|^2 |u_{\mu_n}^T|^{2k} dx \\ &\leq \varepsilon \int_{\Omega} |u_{\mu_n}|^6 |u_{\mu_n}^T|^{2k} dx + C_\varepsilon \int_{\Omega} |u_{\mu_n}|^2 |u_{\mu_n}^T|^{2k} dx. \end{aligned} \quad (4.5)$$

Hence, it follows from the above estimates and the Sobolev imbedding theorem, we can obtain that

$$\begin{aligned} S \left(\int_{\Omega} \left(|u_{\mu_n}^2 |u_{\mu_n}^T|^k \right)^6 dx \right)^{\frac{1}{3}} &\leq \int_{\Omega} \left| \nabla \left(|u_{\mu_n}^2 |u_{\mu_n}^T|^k \right) \right|^2 dx \\ &\leq C(k + 2)^2 \int_{\Omega} |u_{\mu_n}|^2 \left(|u_{\mu_n}^2 |u_{\mu_n}^T|^k \right)^2 dx. \end{aligned} \quad (4.6)$$

Since $|u_{\mu_n}|^2 \in L^s(\Omega)$ for some $s \in (\frac{3}{2}, 3)$, then we have

$$|(u_{\mu_n}^2 |u_{\mu_n}^T|^k)^2|_{L^3} = \left(\int_{\Omega} (u_{\mu_n}^2 |u_{\mu_n}^T|^k)^6 dx \right)^{\frac{1}{3}} \leq C(k+2)^2 |u_{\mu_n}|_{L^2}^2 |(u_{\mu_n}^2 |u_{\mu_n}^T|^k)^2|_{L^{s'}}, \quad (4.7)$$

where $s' = \frac{s}{s-1} < 3$. Assume $u_{\mu_n} \in L^{(4+2k)s'}$. Let $T \rightarrow +\infty$ in (4.7), by the above estimate we have $u_{\mu_n} \in L^{(4+2k)3}$ and

$$|u_{\mu_n}|_{L^{(4+2k)3}} \leq C(k+2)^{\frac{1}{k+2}} |u_{\mu_n}|_{L^{(4+2k)s'}}.$$

Hence, Moser's iteration implies $|u_{\mu_n}|_{\infty} \leq C$.

Claim 3. $u_0 \in E \cap L^{\infty}(\Omega)$ is a solution of problem (1.1). By Claim 1, we may assume $\{u_{\mu_n}\}$ converges to u_0 weakly in E . Taking $\varphi = \psi e^{-u_{\mu_n}}$, where $\psi \in C_0^{\infty}(\Omega)$, $\psi \geq 0$, we have

$$\begin{aligned} & \mu_n \int_{\Omega} (|\nabla u_{\mu_n}|^2 \nabla u_{\mu_n} (\nabla \psi e^{-u_{\mu_n}} - \psi e^{-u_{\mu_n}} \nabla u_{\mu_n}) + |u_{\mu_n}|^2 u_{\mu_n} \psi e^{-u_{\mu_n}}) dx \\ & + b \left(\int_{\Omega} |\nabla u_{\mu_n}|^2 dx \right) \left(\int_{\Omega} \nabla u_{\mu_n} (\nabla \psi e^{-u_{\mu_n}} - \psi e^{-u_{\mu_n}} \nabla u_{\mu_n}) dx \right) \\ & + \int_{\Omega} (a + |u_{\mu_n}|^2) \nabla u_{\mu_n} (\nabla \psi e^{-u_{\mu_n}} - \psi e^{-u_{\mu_n}} \nabla u_{\mu_n}) dx + \int_{\Omega} V(x) u_{\mu_n} \psi e^{-u_{\mu_n}} dx \\ & + \int_{\Omega} |\nabla u_{\mu_n}|^2 u_{\mu_n} \psi e^{-u_{\mu_n}} dx + \int_{\Omega} \phi_{u_{\mu_n}} u_{\mu_n} \psi e^{-u_{\mu_n}} dx \\ & - \lambda \int_{\Omega} |u_{\mu_n}|^{q-2} u_{\mu_n} \psi e^{-u_{\mu_n}} \ln |u_{\mu_n}|^2 dx - \int_{\Omega} |u_{\mu_n}|^4 u_{\mu_n} \psi e^{-u_{\mu_n}} dx \\ \leq & \mu_n \int_{\Omega} (|\nabla u_{\mu_n}|^2 \nabla u_{\mu_n} \nabla \psi + |u_{\mu_n}|^2 u_{\mu_n} \psi) e^{-u_{\mu_n}} dx + \int_{\Omega} (a + |u_{\mu_n}|^2) \nabla u_{\mu_n} \nabla \psi e^{-u_{\mu_n}} dx \\ & + b \left(\int_{\Omega} |\nabla u_{\mu_n}|^2 dx \right) \left(\int_{\Omega} \nabla u_{\mu_n} (\nabla \psi e^{-u_{\mu_n}}) dx \right) + \int_{\Omega} V(x) u_{\mu_n} \psi e^{-u_{\mu_n}} dx \\ & - \int_{\Omega} (a + |u_{\mu_n}|^2 - u_{\mu_n}) |\nabla u_{\mu_n}|^2 \psi e^{-u_{\mu_n}} dx + \int_{\Omega} \phi_{u_{\mu_n}} u_{\mu_n} \psi e^{-u_{\mu_n}} dx \\ & - \lambda \int_{\Omega} |u_{\mu_n}|^{q-2} u_{\mu_n} \psi e^{-u_{\mu_n}} \ln |u_{\mu_n}|^2 dx - \int_{\Omega} |u_{\mu_n}|^4 u_{\mu_n} \psi e^{-u_{\mu_n}} dx. \end{aligned}$$

Since $a > \frac{1}{4}$, then $a + u_{\mu_n}^2 - u_{\mu_n} \geq 0$ and

$$\begin{aligned} & \int_{\Omega} (a + |u_{\mu_n}|^2 - u_{\mu_n}) |\nabla u_{\mu_n}|^2 \psi e^{-u_{\mu_n}} dx \\ & = \int_{\Omega} (a + |u_{\mu_n}|^2 - u_{\mu_n}) \left(|\nabla (u_{\mu_n} - u_0)|^2 + 2 \nabla u_{\mu_n} \nabla u_0 - |\nabla u_0|^2 \right) \psi e^{-u_{\mu_n}} dx \\ & \geq \int_{\Omega} (a + |u_{\mu_n}|^2 - u_{\mu_n}) (2 \nabla u_{\mu_n} \nabla u_0 - |\nabla u_0|^2) \psi e^{-u_{\mu_n}} dx. \end{aligned}$$

Let $v \geq 0$, $v \in C_0^{\infty}(\Omega)$. We choose a sequence of nonnegative functions $\{\psi_n\} \subset C_0^{\infty}(\Omega)$ such that $\psi_n \rightarrow v e^{u_0}$ in E , $\psi_n(x) \rightarrow v e^{u_0}(x)$ a.e. $x \in \Omega$ and $\{\psi_n\}$ is uniformly bounded in $L^{\infty}(\Omega)$. By approximations,

we may obtain for all $v \geq 0$, $v \in C_0^\infty(\Omega)$, we have

$$0 \leq \int_{\Omega} (a \nabla u_0 \nabla v + V(x) u_0 v) dx + b \int_{\Omega} |\nabla u_0|^2 \nabla u_0 \nabla v dx + \int_{\Omega} (|\nabla u_0|^2 u_0 v + |u_0|^2 \nabla u_0 \nabla v) dx \\ + \int_{\Omega} \phi_{u_0} u_0 v dx - \lambda \int_{\Omega} |u_0|^{q-2} v \ln |u_0|^2 dx - \int_{\Omega} |u_0|^4 u_0 v dx.$$

Take $\varphi = \psi e^{u_{\mu_n}}$ in (4.4),

$$0 \geq \int_{\Omega} (a \nabla u_0 \nabla v + V(x) u_0 v) dx + b \int_{\Omega} |\nabla u_0|^2 \nabla u_0 \nabla v dx + \int_{\Omega} (|\nabla u_0|^2 u_0 v + |u_0|^2 \nabla u_0 \nabla v) dx \\ + \int_{\Omega} \phi_{u_0} u_0 v dx - \lambda \int_{\Omega} |u_0|^{q-2} v \ln |u_0|^2 dx - \int_{\Omega} |u_0|^4 u_0 v dx.$$

Therefore, for all $v \in C_0^\infty(\Omega)$, we get

$$\int_{\Omega} (a \nabla u_0 \nabla v + V(x) u_0 v) dx + b \int_{\Omega} |\nabla u_0|^2 \nabla u_0 \nabla v dx \\ + \int_{\Omega} |\nabla u_0|^2 u_0 v + |u_0|^2 \nabla u_0 \nabla v dx + \int_{\Omega} \phi_{u_0} u_0 v dx \\ = \lambda \int_{\Omega} |u_0|^{q-2} v \ln |u_0|^2 dx + \int_{\Omega} |u_0|^4 u_0 v dx.$$

Therefore, u_0 is a critical point of I_b^λ . The proof is completed. \square

Proof of Theorem 1.1. By Lemma 3.4, we choose a sequence $\mu_n \rightarrow 0$, there exists $\{u_{\mu_n}\} \subset E$ satisfies $I_{b,\mu_n}^\lambda(u_{\mu_n}) = c_{b,\mu_n}^\lambda$ and $(I_{b,\mu_n}^\lambda)'(u_{\mu_n}) = 0$.

Claim 1. Problem (1.1) possesses one sign-changing solution u_0 .

Assume $\varphi \in C_0^\infty(\Omega)$ with $\varphi^\pm \neq 0$, there is a pair of positive numbers (α_0, β_0) independent of n such that

$$\langle I_{b,\mu_n}^\lambda(\alpha_0 \varphi^+ + \beta_0 \varphi^-), \alpha_0 \varphi^+ \rangle \leq \langle I_{b,1}^\lambda(\alpha_0 \varphi^+ + \beta_0 \varphi^-), \alpha_0 \varphi^+ \rangle < 0,$$

and

$$\langle I_{b,\mu_n}^\lambda(\alpha_0 \varphi^+ + \beta_0 \varphi^-), \beta_0 \varphi^- \rangle \leq \langle I_{b,1}^\lambda(\alpha_0 \varphi^+ + \beta_0 \varphi^-), \beta_0 \varphi^- \rangle < 0.$$

Let $\varphi_1 = \alpha_0 \varphi^+ + \beta_0 \varphi^-$, according to Lemma 3.1 that there exists a unique pair of positive numbers $(\alpha_n, \beta_n) \subset (0, 1] \times (0, 1]$ such that $\alpha_n \varphi_1^+ + \beta_n \varphi_1^- \in \mathcal{M}_{b,\mu_n}^\lambda$, we have that

$$\begin{aligned}
c_{b,\mu_n}^\lambda &\leq I_{b,\mu_n}^\lambda(\alpha_n\varphi_1^+ + \beta_n\varphi_1^-) - \frac{1}{4} \langle (I_{b,\mu_n}^\lambda)'(\alpha_n\varphi_1^+ + \beta_n\varphi_1^-), \alpha_n\varphi_1^+ + \beta_n\varphi_1^- \rangle \\
&= \frac{1}{4} \left((\alpha_n)^2 \|\varphi_1^+\|^2 + (\beta_n)^2 \|\varphi_1^-\|^2 \right) + \frac{1}{12} \int_\Omega \left((\alpha_n)^6 |\varphi_1^+|^6 + (\beta_n)^6 |\varphi_1^-|^6 \right) dx \\
&\quad + \frac{\lambda}{q^2} \int_\Omega |\alpha_n\varphi_1^+|^q dx + \frac{\lambda}{q^2} \int_\Omega |\beta_n\varphi_1^-|^q dx + \left(\frac{1}{4} - \frac{1}{q} \right) \int_\Omega \left(|\alpha_n\varphi_1^+|^q \ln |\alpha_n\varphi_1^+|^2 \right) dx \\
&\quad + \left(\frac{1}{4} - \frac{1}{q} \right) \int_\Omega \left(|\beta_n\varphi_1^-|^q \ln |\beta_n\varphi_1^-|^2 \right) dx \\
&\leq \frac{1}{4} \left(\|\varphi_1^+\|^2 + \|\varphi_1^-\|^2 \right) + \frac{1}{12} \int_\Omega \left(|\varphi_1^+|^6 + |\varphi_1^-|^6 \right) dx + \frac{\lambda}{q^2} \int_\Omega \left(|\varphi_1^+|^q + |\varphi_1^-|^q \right) dx \\
&\quad + \int_\Omega \left(\frac{1}{4} - \frac{1}{q} \right) |\varphi_1^+|^q \ln |\varphi_1^+|^2 dx + \int_\Omega \left(\frac{1}{4} - \frac{1}{q} \right) |\varphi_1^-|^q \ln |\varphi_1^-|^2 dx \\
&= I_{b,1}^\lambda(\varphi_1) - \frac{1}{4} \langle (I_{b,1}^\lambda)'(\varphi_1), \varphi_1 \rangle.
\end{aligned}$$

Therefore, $\{c_{b,\mu_n}^\lambda\}$ is bounded, according to Lemma 4.2, there exists a critical point u_0 of I_b^λ such that $u_{\mu_n} \rightarrow u_0$ in E .

Now we prove $u_0^\pm \neq 0$. Since $u_{\mu_n} \in \mathcal{M}_{b,\mu_n}^\lambda$, we have that

$$\begin{aligned}
&\mu_n \|u_{\mu_n}^\pm\|_W^4 + \|u_{\mu_n}^\pm\|^2 + b \|u_{\mu_n}^\pm\|_1^4 + 2 \int_\Omega |\nabla u_{\mu_n}^\pm|^2 |u_{\mu_n}^\pm|^2 dx \\
&\quad + \int_\Omega \phi_{u_{\mu_n}^\pm} |u_{\mu_n}^\pm|^2 dx + \int_\Omega \phi_{u_{\mu_n}^\pm} |u_{\mu_n}^\pm|^2 dx \\
&= \int_\Omega |u_{\mu_n}^\pm|^6 dx + \lambda \int_\Omega |u_{\mu_n}^\pm|^q \ln |u_{\mu_n}^\pm|^2 dx.
\end{aligned} \tag{4.8}$$

So, for $\mu_n \rightarrow 0$ and (3.2), we have that

$$\begin{aligned}
\|u_0^\pm\|^2 &\leq \int_\Omega |u_0^\pm|^6 dx + \lambda \int_\Omega |u_0^\pm|^q \ln |u_0^\pm|^2 dx \\
&\leq +\lambda \varepsilon C_1 \|u_0^\pm\|^2 + \lambda C_\varepsilon C_2 \|u_0^\pm\|^r C_3 \|u_0^\pm\|^6.
\end{aligned}$$

Thus, we get

$$(1 - \lambda \varepsilon C_1) \|u_0^\pm\|^2 \leq C_8 \|u_0^\pm\|^6 + \lambda C_\varepsilon C_{10} \|u_0^\pm\|^r.$$

Choosing ε small enough such that $(1 - \mu \varepsilon C_1) > 0$, since $r > 4$, there exists u_0 such that

$$\|u_0^\pm\|^2 \geq \rho > 0.$$

Therefore, $u_0^\pm \neq 0$. Then we obtain that u_0 is a sign-changing solution of (1.1).

Claim 2. u_0 has also exactly two nodal domains.

Since u_0 is a nonzero critical point of I_b^λ , we have that

$$\begin{aligned}
&\int_\Omega (a|\nabla u_0|^2 + V(x)|u_0|^2) dx + b \left(\int_\Omega |\nabla u_0|^2 dx \right)^2 + 2 \int_\Omega |\nabla u_0|^2 |u_0|^2 dx \\
&\quad + \int_\Omega \phi_{u_0} |u_0|^2 dx - \lambda \int_\Omega |u_0|^q \ln |u_0|^2 dx - \int_\Omega |u_0|^6 dx = 0.
\end{aligned} \tag{4.9}$$

On the other hand, $\langle (I_{b,\mu_n}^\lambda)'(u_{\mu_n}), u_{\mu_n} \rangle = 0$ implies that

$$\begin{aligned} & \mu_n \int_{\Omega} (|\nabla u_{\mu_n}|^4 + |u_{\mu_n}|^4) dx + \int_{\Omega} (a|\nabla u_{\mu_n}|^2 + V(x)|u_{\mu_n}|^2) dx + b \left(\int_{\Omega} |\nabla u_{\mu_n}|^2 dx \right)^2 \\ & + 2 \int_{\Omega} |\nabla u_{\mu_n}|^2 |u_{\mu_n}|^2 dx + \int_{\Omega} \phi_{u_{\mu_n}} |u_{\mu_n}|^2 dx - \lambda \int_{\Omega} |u_{\mu_n}|^q \ln |u_{\mu_n}|^2 dx - \int_{\Omega} |u_{\mu_n}|^6 dx = 0. \end{aligned}$$

According to (3.2), we have

$$\lim_{n \rightarrow \infty} \lambda \int_{\Omega} (|u_{\mu_n}|^q) \ln |u_{\mu_n}|^2 dx \rightarrow \lambda \int_{\Omega} (|u_0|^q) \ln |u_0|^2 dx. \quad (4.10)$$

Moreover, according to the proof in Lemma 3.4 that $B_1 = B_2 = 0$, we have

$$\lim_{n \rightarrow \infty} \int_{\Omega} |u_{\mu_n}|^6 dx = \int_{\Omega} |u_0|^6 dx = 0. \quad (4.11)$$

Then, combining (4.8)–(4.11) and using Fatou's lemma and weak semicontinuity of norm, up to a subsequence, we get

$$\begin{aligned} & \int_{\Omega} (a|\nabla u_0|^2 + V(x)|u_0|^2) dx + b \left(\int_{\Omega} |\nabla u_0|^2 dx \right)^2 \\ & \leq \lim_{n \rightarrow \infty} \int_{\Omega} (a|\nabla u_{\mu_n}|^2 + V(x)|u_{\mu_n}|^2) dx + b \left(\int_{\Omega} |\nabla u_{\mu_n}|^2 dx \right)^2 \\ & = \lim_{n \rightarrow \infty} \left(\lambda \int_{\Omega} |u_{\mu_n}|^q \ln |u_{\mu_n}|^2 dx + \int_{\Omega} |u_{\mu_n}|^6 dx - 2 \int_{\Omega} |\nabla u_{\mu_n}|^2 |u_{\mu_n}|^2 dx - \int_{\Omega} \phi_{u_{\mu_n}} |u_{\mu_n}|^2 dx \right) \\ & = \int_{\Omega} (a|\nabla u_0|^2 + V(x)|u_0|^2) dx + b \left(\int_{\Omega} |\nabla u_0|^2 dx \right)^2 + 2 \int_{\Omega} |\nabla u_0|^2 |u_0|^2 dx + \int_{\Omega} \phi_{u_0} |u_0|^2 dx \\ & \quad - \lim_{n \rightarrow \infty} \left(2 \int_{\Omega} |\nabla u_{\mu_n}|^2 |u_{\mu_n}|^2 dx - \int_{\Omega} \phi_{u_{\mu_n}} |u_{\mu_n}|^2 dx \right) \\ & \leq \int_{\Omega} (a|\nabla u_0|^2 + V(x)|u_0|^2) dx + b \left(\int_{\Omega} |\nabla u_0|^2 dx \right)^2. \end{aligned}$$

So $\lim_{n \rightarrow \infty} \|u_{\mu_n}\|_W^2 = \|u_0\|_W^2$. According to Brézis-Lieb lemma that $u_{\mu_n} \rightarrow u_0$ strongly in E as $n \rightarrow \infty$. It means that u_0 has also exactly two nodal domains.

Claim 3. u_0 is a least-energy sign-changing solution.

By Lemma 3.1 it is easy to see that there exists a unique pair $(\alpha_{\mu_n}, \beta_{\mu_n}) \in (0, \infty) \times (0, \infty)$ such that $\alpha_{\mu_n} u_0^+ + \beta_{\mu_n} u_0^- \in \mathcal{M}_{b,\mu_n}^\lambda$. Then we have

$$\begin{aligned} & \mu_n \alpha_{\mu_n}^4 \|u_0^+\|_W^4 + \alpha_{\mu_n}^4 \|u_0^+\|^2 + b \alpha_{\mu_n}^2 \|u_0^+\|_1^4 + b \alpha_{\mu_n}^2 \beta_{\mu_n}^2 \|u_0^+\|_1^2 \|u_0^-\|_1^2 \\ & + 2 \alpha_{\mu_n}^4 \int_{\Omega} |\nabla u_0^+|^2 |u_0^+|^2 dx + \alpha_{\mu_n}^4 \int_{\Omega} \phi_{u_0^+} |u_0^+|^2 dx + \alpha_{\beta_n}^2 \beta_{\mu_n}^2 \int_{\Omega} \phi_{u_0^-} |u_0^+|^2 dx \\ & = \lambda \int_{\Omega} |\alpha_{\mu_n} u_0^+|^q \ln |\alpha_{\mu_n} u_0^+|^2 dx + \alpha_{\mu_n}^6 \int_{\Omega} |u_0^+|^6 dx, \end{aligned} \quad (4.12)$$

and

$$\begin{aligned} & \mu_n \beta_{\mu_n}^4 \|u_0^-\|_W^4 + \beta_{\mu_n}^4 \|u_0^-\|^2 + b \beta_{\mu_n}^2 \|u_0^-\|_1^4 + b \alpha_{\mu_n}^2 \beta_{\mu_n}^2 \|u_0^+\|_1^2 \|u_0^-\|_1^2 \\ & + 2 \beta_{\mu_n}^4 \int_{\Omega} |\nabla u_0^-|^2 |u_0^-|^2 dx + \beta_{\mu_n}^4 \int_{\Omega} \phi_{u_0^-} |u_0^-|^2 dx + \alpha_{\mu_n}^2 \beta_{\mu_n}^2 \int_{\Omega} \phi_{u_0^+} |u_0^-|^2 dx \\ & = \lambda \int_{\Omega} |\beta_{\mu_n} u_0^-|^q \ln |\beta_{\mu_0} u_0^-|^2 dx + \beta_{\mu_n}^6 \int_{\Omega} |u_0^-|^6 dx. \end{aligned} \quad (4.13)$$

According to $\mu_n \rightarrow 0$ as $n \rightarrow \infty$, $\{\alpha_{\mu_n}\}$ and $\{\beta_{\mu_n}\}$ are bounded. Up to a subsequence, suppose that $\alpha_{\mu_n} \rightarrow \alpha_0$ and $\beta_{\mu_n} \rightarrow \beta_0$, then it follows from (4.12) and (4.13) that

$$\begin{aligned} & \alpha_0^2 \|u_0^+\|^2 + b \alpha_0^2 \|u_0^+\|_1^4 + b \alpha_0^2 \beta_0^2 \|u_0^+\|_1^2 \|u_0^-\|_1^2 \\ & + 2 \alpha_0^4 \int_{\Omega} |\nabla u_0^+|^2 |u_0^+|^2 dx + \alpha_0^4 \int_{\Omega} \phi_{u_0^+} |u_0^+|^2 dx + \alpha_0^2 \beta_0^2 \int_{\Omega} \phi_{u_0^-} |u_0^+|^2 dx \\ & = \lambda \int_{\Omega} |\alpha_0 u_0^+|^q \ln |\alpha_0 u_0^+|^2 dx + \alpha_0^6 \int_{\Omega} |u_0^+|^6 dx \end{aligned} \quad (4.14)$$

and

$$\begin{aligned} & \beta_0^2 \|u_0^-\|^2 + b \beta_0^2 \|u_0^-\|_1^4 + b \alpha_0^2 \beta_0^2 \|u_0^+\|_1^2 \|u_0^-\|_1^2 \\ & + 2 \beta_0^4 \int_{\Omega} |\nabla u_0^-|^2 |u_0^-|^2 dx + \beta_0^4 \int_{\Omega} \phi_{u_0^-} |u_0^-|^2 dx + \alpha_0^2 \beta_0^2 \int_{\Omega} \phi_{u_0^+} |u_0^-|^2 dx \\ & = \lambda \int_{\Omega} (\beta_0 u_0^-)^q \ln |\beta_{\mu_0} u_0^-|^2 dx + \beta_0^6 \int_{\Omega} |u_0^-|^6 dx. \end{aligned} \quad (4.15)$$

Because u_0 is a sign-changing solution of problem (1.1), there hold

$$\begin{aligned} & \|u_0^+\|^2 + b \|u_0^+\|_1^4 + b \|u_0^+\|_1^2 \|u_0^-\|_1^2 \\ & + 2 \int_{\Omega} |\nabla u_0^+|^2 |u_0^+|^2 dx + \int_{\Omega} \phi_{u_0^+} |u_0^+|^2 dx + \int_{\Omega} \phi_{u_0^-} |u_0^+|^2 dx \\ & = \lambda \int_{\Omega} (|u_0^+|^q) \ln |u_0^+|^2 dx + \int_{\Omega} |u_0^+|^6 dx \end{aligned} \quad (4.16)$$

and

$$\begin{aligned} & \|u_0^-\|^2 + b \|u_0^-\|_1^4 + b \|u_0^-\|_1^2 \|u_0^+\|_1^2 \\ & + 2 \int_{\Omega} |\nabla u_0^-|^2 |u_0^-|^2 dx + \int_{\Omega} \phi_{u_0^-} |u_0^-|^2 dx + \int_{\Omega} \phi_{u_0^+} |u_0^-|^2 dx \\ & = \lambda \int_{\Omega} (|u_0^-|^q) \ln |u_0^-|^2 dx + \int_{\Omega} |u_0^-|^6 dx. \end{aligned} \quad (4.17)$$

Hence, in view of (4.12)–(4.17), we can easily obtain that $(\alpha_0, \beta_0) = (1, 1)$.

According to Fatou's lemma and weak semicontinuity of norm, we get

$$\begin{aligned}
 & I_b^\lambda(u_0) - \frac{1}{4} \langle (I_b^\lambda)'(u_0), u_0 \rangle \\
 &= \frac{1}{4} \|u_0\|^2 + \frac{1}{12} |u_0|_6^6 + \left(\frac{1}{4} - \frac{1}{q}\right) \lambda \int_{\Omega} |u_0|^q \ln |u_0|^2 dx + \frac{2\lambda}{q^2} \int_{\Omega} |u_0|^q dx \\
 &\leq \frac{1}{4} \|u_{\mu_n}\|^2 + \frac{1}{12} |u_{\mu_n}|_6^6 + \left(\frac{1}{4} - \frac{1}{q}\right) \lambda \int_{\Omega} |u_n|^q \ln |u_n|^2 dx + \frac{2\lambda}{q^2} \int_{\Omega} |u_n|^q dx \\
 &\leq \liminf_{n \rightarrow \infty} \left[I_{b,\mu_n}^\lambda(u_{\mu_n}) - \frac{1}{4} \langle (I_{b,\mu_n}^\lambda)'(u_{\mu_n}), u_{\mu_n} \rangle \right] \\
 &= \liminf_{n \rightarrow \infty} I_{b,\mu_n}^\lambda(u_{\mu_n}) = \lim_{n \rightarrow \infty} c_{b,\mu_n}^\lambda = c_{b,0}^\lambda.
 \end{aligned}$$

Moreover,

$$c_{b,0}^\lambda = \liminf_{n \rightarrow \infty} I_{b,\mu_n}^\lambda(u_{\mu_n}) \leq \liminf_{n \rightarrow \infty} I_{b,\mu_n}^\lambda(\alpha_{\mu_n} u_0^+ + \beta_{\mu_n} u_0^-) = I_b^\lambda(u_0^+ + u_0^-) = I_b^\lambda(u_0).$$

So $I_b^\lambda(u_0) = \liminf_{n \rightarrow \infty} I_{b,\mu_n}^\lambda(u_{\mu_n}) = c_{b,0}^\lambda$. The proof is completed. \square

We get a least-energy sign-changing solution u_0 of issue using Theorem 1.1. Following that, we show that the energy of u_0 is strictly greater than twice the ground state energy.

Proof of Theorem 1.2. Similar to the proof of Lemma 3.4, there exists $\lambda^{**} > 0$ such that for all $\lambda \geq \lambda^{**}$ and when $\mu \rightarrow 0$, there is $v_\mu \in \mathcal{N}_b^\lambda$ such that $I_{b,\mu}^\lambda(v_\mu) = c^* > 0$. The critical points of the functional $I_{b,\mu}^\lambda$ in \mathcal{N}_b^λ are critical points of $I_{b,\mu}^\lambda$ in E are determined using conventional reasoning (see Corollary 2.13 in [11]).

For all $\lambda \geq \lambda^*$, according to Theorem 1.1, for each $\mu \rightarrow 0$, we know that the problem (1.1) has a least-energy sign-changing solution u_0 which changes sign only once. Let $\lambda_2 = \max\{\lambda^*, \lambda^{**}\}$. Suppose that $u_0 = u^+ + u^-$. As the proof of Lemma 3.1, there exist $\alpha_{u^+}, \beta_{u^-} \in (0, 1)$ such that $\alpha_{u^+} u^+ \in \mathcal{N}_b^\lambda, \beta_{u^-} u^- \in \mathcal{N}_b^\lambda$. Therefore, in view of Lemma 3.1, we have that

$$\begin{aligned}
 2c^* &\leq \liminf_{\mu \rightarrow 0} \left[I_{b,\mu}^\lambda(\alpha_{u^+} u^+) + I_{b,\mu}^\lambda(\beta_{u^-} u^-) \right] \\
 &\leq \liminf_{\mu \rightarrow 0} I_{b,\mu}^\lambda(\alpha_{u^+} u^+ + \beta_{u^-} u^-) < \liminf_{\mu \rightarrow 0} I_{b,\mu}^\lambda(u^+ + u^-) = I_b^\lambda(u_0).
 \end{aligned}$$

which shows that $I_b^\lambda(u_0) > 2c^*$ and $c^* > 0$ cannot be achieved by a sign-changing function in E . \square

5. Conclusions

This manuscript has studied the Kirchhoff-Schrödinger-Poisson system with logarithmic and critical nonlinearity. Combining constraint variational method and perturbation method, we prove that the above problem has a least energy sign-changing solution u_0 which has precisely two nodal domains. Moreover, we show that the energy of u_0 is strictly larger than twice the ground state energy.

Acknowledgments

We would like to express our deep gratitude to the anonymous reviewers and editors for their helpful comments and suggestions which led to the improvement of this manuscript.

H. Jian was supported by Jiangxi Provincial Natural Science Foundation (Grant No. 20212BAB211006). L. Wang was supported by the National Natural Science Foundation of China (Grant No. 12161038), and the Science and Technology project of Jiangxi provincial Department of Education (Grant Nos. GJJ212204, GJJ2200635)

Conflict of interest

The authors declare that they have no conflict of interest.

References

1. C. O. Alves, F. J. S. A. Correa, On existence of solutions for a class of problem involving a nonlinear operator, *Comm. Appl. Nonlinear Anal.*, **8** (2001), 43–56.
2. A. Arosio, S. Panizzi, On the well-posedness of the Kirchhoff string, *Trans. Amer. Math. Soc.*, **348** (1996), 305–330. <http://dx.doi.org/10.1090/S0002-9947-96-01532-2>
3. T. Bartsch, T. Weth, Three nodal solutions of singularly perturbed elliptic equations on domains without topology, *Ann. I. H. Poincaré Anal. NonLinéaire*, **22** (2005), 259–281. <http://dx.doi.org/10.1016/j.anihpc.2004.07.005>
4. M. M. Cavalcanti, V. N. D. Cavalcanti, J. A. Soriano, Global existence and uniform decay rates for the Kirchhoff-Carrier equation with nonlinear dissipation, *Adv. Differential Equ.*, **6** (2001), 701–730. <http://dx.doi.org/10.57262/ade/1357140586>
5. J. Chen, X. Tang, Z. Gao, B. Chen, Ground state sign-changing solutions for a class of generalized quasilinear Schrödinger equations with a Kirchhoff-type perturbation, *J. Fixed Point Theory A.*, **19** (2017), 3127–3149. <http://dx.doi.org/10.1007/s11784-017-0475-4>
6. P. D’Ancona, S. Spagnolo, Global solvability for the degenerate Kirchhoff equation with real analytic data, *Invent. Math.*, **108** (1992), 247–262. <http://dx.doi.org/10.1007/BF02100605>
7. Y. Deng, W. Shuai, Sign-changing solutions for non-local elliptic equations involving the fractional Laplacian, *Adv. Differential Equ.*, **23** (2018), 109–134. <http://dx.doi.org/10.57262/ade/1508983363>
8. X. Feng, Y. Zhang, Existence of non-trivial solution for a class of modified Schrödinger-Poisson equations via perturbation method, *J. Math. Anal. Appl.*, **442** (2016), 673–684. <http://dx.doi.org/10.1016/j.jmaa.2016.05.002>
9. G. M. Figueiredo, G. Siciliano, Existence and asymptotic behaviour of solutions for a quasilinear Schrödinger-Poisson system under a critical nonlinearity, *arXiv:1707.05353*, 2017. <https://doi.org/10.48550/arXiv.1707.05353>
10. G. M. Figueiredo, G. Siciliano, Quasilinear Schrödinger-Poisson system under an exponential critical nonlinearity: existence and asymptotic of solutions, *Arch. Math.*, **112** (2019), 313–327. <http://dx.doi.org/10.1007/s00013-018-1287-5>

11. X. He, W. Zou, Ground states for nonlinear Kirchhoff equations with critical growth, *Ann. Mat. Pura. Appl.*, **193** (2014), 473–500. <http://dx.doi.org/10.1007/s10231-012-0286-6>
12. R. Illner, O. Kavian, H. Lange, Stationary solutions of quasi-linear Schrödinger-Poisson systems, *J. Differential Equ.*, **145** (1998), 1–16. <http://dx.doi.org/10.1006/jdeq.1997.3405>
13. G. Kirchhoff, *Mechanik*, Leipzig: Teubner, 1883.
14. Y. Li, D. Wang, J. Zhang, Sign-changing solutions for a class of p -Laplacian Kirchhoff-type problem with logarithmic nonlinearity, *AIMS Math.*, **5** (2020), 2100–2112. <http://dx.doi.org/10.3934/math.2020139>
15. F. Li, X. Zhu, Z. Liang, Multiple solutions to a class of generalized quasilinear Schrödinger equations with a Kirchhoff-type perturbation, *J. Math. Anal. Appl.*, **443** (2016), 11–38. <http://dx.doi.org/10.1016/j.jmaa.2016.05.005>
16. S. Liang, V. D. Rădulescu, Least-energy nodal solutions of critical Kirchhoff problems with logarithmic nonlinearity, *Anal. Math. Phys.*, **10** (2020), 45. <http://dx.doi.org/10.1007/s13324-020-00386-z>
17. J. L. Lions, On some questions in boundary value problems of mathematical physics, *North-Holland Math. Stud.*, **30** (1978), 284–346.
18. X. Liu, J. Liu, Z. Wang, Quasilinear elliptic equations with critical growth via perturbation method, *J. Differential Equ.*, **254** (2013), 102–124. <http://dx.doi.org/10.1016/j.jde.2012.09.006>
19. M. Massar, On a nonlocal Schrödinger-Poisson system with critical exponent, *Appl. Math. E-Notes*, **21** (2021), 44–52.
20. C. Miranda, Unosservazione su un teorema di Brouwer, *Boll. Un. Mat. Ital.*, **3** (1940), 5–7.
21. A. Nakamura, Damping and modification of exciton solitary waves, *J. Phys. Soc. Japan*, **42** (1977), 1824–1835. <http://doi.org/10.1143/JPSJ.42.1824>
22. W. Shuai, Sign-changing solutions for a class of Kirchhoff-type problem in bounded domains, *J. Differential Equ.*, **259** (2015), 1256–1274. <http://dx.doi.org/10.1016/j.jde.2015.02.040>
23. K. Susumu, Large-Amplitude Quasi-Solitons in Superfluid Films, *J. Phys. Soc. Japan*, **50** (1981), 3262–3267. <http://doi.org/10.1143/JPSJ.50.3262>
24. X. Tang, B. Cheng, Ground state sign-changing solutions for Kirchhoff type problems in bounded domains, *J. Differential Equ.*, **261** (2016), 2384–2402. <http://dx.doi.org/10.1016/j.jde.2016.04.032>
25. K. Teng, Existence of ground state solutions for the nonlinear fractional Schrödinger-Poisson system with critical Sobolev exponent, *J. Differential Equ.*, **261** (2016), 3061–3106. <http://dx.doi.org/10.1016/j.jde.2016.05.022>
26. D. Wang, Least energy sign-changing solutions of Kirchhoff-type equation with critical growth, *J. Math. Phys.*, **61** (2020), 011501. <http://dx.doi.org/10.1063/1.5074163>
27. L. Wang, On a quasilinear Schrödinger-Kirchhoff-type equation with radial potentials, *Nonlinear Anal.*, **83** (2013), 58–68. <http://dx.doi.org/10.1016/j.na.2012.12.012>
28. M. Willem, *Minimax theorems*, In: Progress in Nonlinear Differential Equations and their Applications, Boston: Birkhäuser, **24** (1996).



AIMS Press

©2023 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)