

---

*Research article*

## On Hardy-Hilbert-type inequalities with $\alpha$ -fractional derivatives

**Marwa M. Ahmed<sup>1</sup>, Wael S. Hassanein<sup>2</sup>, Marwa Sh. Elsayed<sup>3</sup>, Dumitru Baleanu<sup>4,5,6</sup> and Ahmed A. El-Deeb<sup>7,\*</sup>**

<sup>1</sup> Department of Electrical and Computer Engineering, Faculty of Engineering-Girls Campus, King Abdulaziz University, Jeddah, Saudi Arabia

<sup>2</sup> Department of industrial Engineering Faculty of Engineering King Abdulaziz University, Jeddah, Saudi Arabia

<sup>3</sup> Institute of Culture and Science, 6th October City, Cairo 13759, Egypt

<sup>4</sup> Institute of Space Science, Magurele-Bucharest, Romania

<sup>5</sup> Department of Mathematics, Cankaya University, Ankara 06530, Turkey

<sup>6</sup> Department of Medical Research, China Medical University, 40402, Taiwan

<sup>7</sup> Department of Mathematics, Faculty of Science, Al-Azhar University, Nasr City (11884), Cairo, Egypt

\* **Correspondence:** Email: ahmedeldeeb@zhar.edu.eg; Tel: +201098802022.

**Abstract:** In the current manuscript, new alpha delta dynamic Hardy-Hilbert inequalities on time scales are discussed. These inequalities combine and expand a number of continuous inequalities and their corresponding discrete analogues in the literature. We shall illustrate our results using Hölder's inequality on time scales and a few algebraic inequalities.

**Keywords:** Steffensen's inequality; dynamic inequality; dynamic integral; time scale

**Mathematics Subject Classification:** 26D10, 26D15, 26D20, 34A12, 34A40

---

### 1. Introduction

Hardy-Hilbert's double-series theorem [1] states:

**Theorem 1.1.** If  $\nu, \varpi > 1$  are such that  $\frac{1}{\nu} + \frac{1}{\varpi} \leq 1$  and  $0 < \lambda = 2 - \frac{1}{\nu} - \frac{1}{\varpi} = \frac{1}{\nu'} + \frac{1}{\varpi'} \leq 1$ , such that  $\nu'$  and  $\varpi'$  present the exponents conjugate, then

$$\sum_{j=1}^{\infty} \sum_{i=1}^{\infty} \frac{\vartheta_j \pi_i}{(j+i)^{\lambda}} \leq K \left( \sum_{j=1}^{\infty} \vartheta_j^{\nu} \right)^{\frac{1}{\nu}} \left( \sum_{i=1}^{\infty} \pi_i^{\varpi} \right)^{\frac{1}{\varpi}}, \quad (1.1)$$

where  $K = K(\nu, \varpi)$  depends on  $\nu$  and  $\varpi$  only.

You may find the integral analogue of Theorem 1.1 in [1].

**Theorem 1.2.** Let  $\nu, \varpi, \nu', \varpi'$  and  $\lambda$  be as in Theorem 1.1. If  $\vartheta \in L^\nu(0, \infty)$  and  $\theta \in L^\varpi(0, \infty)$ , then

$$\int_0^\infty \int_0^\infty \frac{\vartheta(\iota)\theta(\varsigma)}{(\iota + \varsigma)^\lambda} d\iota d\varsigma \leq K \left( \int_0^\infty \vartheta^\nu(\iota) d\iota \right)^{\frac{1}{\nu}} \left( \int_0^\infty \theta^\varpi(\varsigma) d\varsigma \right)^{\frac{1}{\varpi}}, \quad (1.2)$$

where  $K = K(\nu, \varpi)$  depends on  $\nu$  and  $\varpi$  only.

In 2000, Pachpatte [2] established different inequalities from inequality (1.1) but, to a certain extent, having a glimpse of inequality (1.1) as follows:

**Theorem 1.3.** Let  $\nu, \varpi, a(\mathfrak{J}), b(\delta), a(0), b(0), \Delta^\alpha a(\mathfrak{J})$  and  $\Delta^\alpha b(\delta)$  be as in [2] then

$$\sum_{\mathfrak{J}=1}^m \sum_{\delta=1}^n \frac{|a(\mathfrak{J})||b(\delta)|}{qs^{\nu-1} + pt^{\varpi-1}} \leq \frac{1}{pq} m^{\frac{\nu-1}{\nu}} n^{\frac{\varpi-1}{\varpi}} \left( \sum_{\mathfrak{J}=1}^m (m - \mathfrak{J} + 1) |\Delta^\alpha a(\mathfrak{J})|^\nu \right)^{\frac{1}{\nu}} \left( \sum_{\delta=1}^n (n - \delta + 1) |\Delta^\alpha b(\delta)|^\varpi \right)^{\frac{1}{\varpi}}. \quad (1.3)$$

In the same paper [2], Pachpatte also established more advanced versions of inequality (1.2) as follows:

**Theorem 1.4.** Let  $\nu, \varpi, \vartheta(\mathfrak{J}), \theta(\delta), \vartheta(0), \theta(0), \vartheta'(\mathfrak{J})$  and  $\theta'(\delta)$  be as in [2], then

$$\int_0^\iota \int_0^s \frac{|\vartheta(\mathfrak{J})||\theta(\delta)|}{qs^{\nu-1} + pt^{\varpi-1}} d\mathfrak{J} d\delta \leq \frac{1}{pq} \iota^{\frac{\nu-1}{\nu}} s^{\frac{\varpi-1}{\varpi}} \left( \int_0^\iota (\iota - \mathfrak{J}) |\vartheta'(\mathfrak{J})|^\nu d\mathfrak{J} \right)^{\frac{1}{\nu}} \left( \int_0^s (\varsigma - \delta) |\theta'(\delta)|^\varpi d\delta \right)^{\frac{1}{\varpi}}. \quad (1.4)$$

In 2011, Zhao et al. [3] proposed a new inequality similar to Theorem 1.2.

**Theorem 1.5.** Let  $h_i \geq 1, \nu_i > 1$  be constants and  $\frac{1}{\nu_i} + \frac{1}{\varpi_i} = 1$ . Let the differentiable fun.  $\vartheta_i(\mathfrak{J}_i)$  on  $[0, \iota_i]$ , where  $\iota_i \in (0, \infty)$  and we use  $\vartheta'_i$  as differentiation of  $\vartheta_i$ . Suppose  $\vartheta_i(0) = 0$  for  $(i = 1, \dots, n)$ . Then

$$\begin{aligned} & \int_0^{\iota_1} \int_0^{\iota_2} \cdots \int_0^{\iota_n} \frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{J}_i)|}{\left( \sum_{i=1}^n \frac{\mathfrak{J}_i}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} d\mathfrak{J}_n d\mathfrak{J}_{n-1} \cdots d\mathfrak{J}_1 \\ & \leq K \prod_{i=1}^n \left( \int_0^{\iota_i} (\iota_i - \mathfrak{J}_i) |\vartheta_i^{h_i-1}(\mathfrak{J}_i) \vartheta'_i(\mathfrak{J}_i)|^{\nu_i} d\mathfrak{J}_i \right)^{\frac{1}{\nu_i}}, \end{aligned}$$

where

$$K = \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n h_i \iota_i^{\frac{1}{\varpi_i}}.$$

Also in 2012, Zhao and Chung [4] proved the following theorem:

**Theorem 1.6.** Let  $\nu_i > 1$ , be constants and  $\frac{1}{\nu_i} + \frac{1}{\varpi_i} = 1$ . Let  $\vartheta_i(\tau_{1i}, \dots, \tau_{ni})$  be real valued nth differentiable functions defined on  $[0, \iota_{1i}] \times \cdots \times [0, \iota_{ni}]$ , where  $0 \leq \iota_{ji} \leq \delta_{ji}, \delta_{ji} \in (0, \infty)$  and  $i, j = 1, \dots, n$ . Suppose

$$\vartheta_i(\iota_{1i}, \dots, \iota_{ni}) = \int_0^{\iota_{1i}} \cdots \int_0^{\iota_{ni}} \frac{\partial^n}{\partial \tau_{1i} \cdots \partial \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) d\tau_{ni} \cdots d\tau_{1i},$$

then

$$\begin{aligned}
 & \int_0^{\delta_{11}} \cdots \int_0^{\delta_{n1}} \int_0^{\delta_{12}} \cdots \int_0^{\delta_{n2}} \cdots \int_0^{\delta_{1n}} \cdots \int_0^{\delta_{nn}} \\
 & \frac{\prod_{i=1}^n \left( \int_0^{\iota_{1i}} \cdots \int_0^{\iota_{ni}} \left| \frac{\partial^n}{\partial \tau_{1i} \cdots \partial \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right|^{\nu_i} d\tau_{ni} \cdots d\tau_{1i} \right)^{\frac{1}{\nu_i}}}{\left( \sum_{i=1}^n \left[ \frac{\iota_{1i} \cdots \iota_{ni}}{\varpi_i} \right] \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \\
 & d\iota_{11} \cdots d\iota_{n1} d\iota_{12} \cdots d\iota_{n2} \cdots d\iota_{1n} \cdots d\iota_{nn} \\
 & \leq N \prod_{i=1}^n \left( \int_0^{\delta_{1i}} \cdots \int_0^{\delta_{ni}} \prod_{j=1}^n (\delta_{ji} - \iota_{ji}) \left| \frac{\partial^n}{\partial \iota_{1i} \cdots \partial \iota_{ni}} \vartheta_i(\iota_{1i}, \dots, \iota_{ni}) \right|^{\nu_i} d\iota_{ni} \cdots d\iota_{1i} \right)^{\frac{1}{\nu_i}},
 \end{aligned} \tag{1.5}$$

where

$$N = \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n [\delta_{1i} \cdots \delta_{ni}]^{\frac{1}{\varpi_i}}.$$

Fractional calculus on fractal sets has been widely applied in science and engineering because it is better able to describe natural events. Many academics have recently used fractal sets and fractional calculus to study various classical inequalities.

All of the aforementioned findings hold true for both continuous and discrete domains. The purpose of the current research is to provide new, more general conclusions to the time-scale-based disparities previously established. Supreme outcomes, from which many other previous and current results may be taken, would be produced in this way. See the publications for various dynamic inequalities, integrals of Hilbert's kind, and other categories of inequalities on time scales [5–8, 11–20].

A very important question here is: Is it possible to prove the time scales version via  $\alpha$ -conformable derivative of the inequalities of Hardy-Hilbert-type presented in [3] due to Zhao et al. and also the inequalities presented in [4] due to Zhao and Chung?

In this manuscript, we intend to address the question above, and establish a few novel delta fractional dynamic inequalities of the Hardy-Hilbert type on time scales, which are studied in [3, 4]. We also extract the discrete counterparts of the continuous Hilbert inequalities that are present in some special situations of our results. The present article is arranged as follows: In Section 2, some basic concepts of the  $\alpha$ -fractional calculus on time scales and useful lemmas are introduced. In Section 3, we state and prove our main results. We end with Section 4 of conclusion.

## 2. Preliminaries on time scales

Now, we present the fundamental results about the fractional time scales calculus. The results are adapted from [9, 10, 21–24]. We assume throughout that  $\mathbb{T}$  has the topology that it inherits from the standard topology on the real numbers  $\mathbb{R}$ . We define the forward jump operator  $\sigma : \mathbb{T} \rightarrow \mathbb{T}$  by

$$\sigma(\zeta) := \inf\{s \in \mathbb{T} : s > \zeta\}, \quad \zeta \in \mathbb{T}, \tag{2.1}$$

and the backward jump operator  $\rho : \mathbb{T} \rightarrow \mathbb{T}$  is defined by

$$\rho(\zeta) := \sup\{s \in \mathbb{T} : s < \zeta\}, \quad \zeta \in \mathbb{T}. \tag{2.2}$$

In the previous two definitions, we set  $\inf \emptyset = \sup \mathbb{T}$  (i.e., if  $\zeta$  is the minimum of  $\mathbb{T}$ , then  $\sigma(\zeta) = \zeta$ ) and  $\sup \emptyset = \inf \mathbb{T}$  (i.e., if  $\zeta$  is the maximum of  $\zeta$ , then  $\rho(\zeta) = \zeta$ ), where  $\emptyset$  is the empty set.

A point  $\zeta \in \mathbb{T}$  with  $\inf \mathbb{T} < \zeta < \sup \mathbb{T}$ , is said to be right-dense if  $\sigma(\zeta) = \zeta$ , left-dense if  $\rho(\zeta) = \zeta$ , right-scattered if  $\sigma(\zeta) > \zeta$ , and left-scattered if  $\rho(\zeta) < \zeta$ . Points that are simultaneously right-dense and left-dense are called dense points, and points that are simultaneously right-scattered and left-scattered are called isolated points. The forward and backward graininess functions  $\mu$  and  $\nu$  for a time scale  $\mathbb{T}$  are defined by  $\mu(\zeta) := \sigma(\zeta) - \zeta$  and  $\nu(\zeta) := \zeta - \rho(\zeta)$ , respectively.

**Definition 2.1.** The number  $T_\alpha^\Delta(f)(\zeta)$  (provided it exists) of the function  $f : \mathbb{T} \rightarrow \mathbb{R}$ , for  $\zeta > 0$  and  $\alpha \in (0, 1]$  is the number which has the property that for any  $\epsilon > 0$ , there exists a neighborhood  $U$  of  $\zeta$  such that

$$|[f(\sigma(\zeta)) - f(s)]\zeta^{1-\alpha} - T_\alpha^\Delta(f(\zeta))[\sigma(\zeta) - s]| \leq \epsilon|\sigma(\zeta) - s|,$$

for all  $s \in U$ .  $T_\alpha^\Delta(f(\zeta))$  is called conformable  $\alpha$ -fractional derivative of function  $f$  of order  $\alpha$  at  $\zeta$ , for conformable fractional derivative on  $\mathbb{T}$  at 0, we define it with  $T_\alpha^\Delta(f(0)) = \lim_{\zeta \rightarrow 0} T_\alpha^\Delta(f(\zeta))$ .

**Remark 2.1.** If  $\alpha = 1$  then we obtain from Definition 2.1 the delta derivative of time scales. The conformable fractional derivative of order zero is defined by the identity operator:  $T_0^\Delta(\eta) = \eta$ .

**Remark 2.2.** Along the work, we also use the notation  $(\eta(\zeta))^{\Delta_\alpha} = T_\alpha^\Delta(\eta)$ .

**Theorem 2.1.** Let  $\alpha \in (0, 1]$  and  $\mathbb{T}$  be a time scale. Assume  $\eta : \mathbb{T} \rightarrow \mathbb{R}$  and  $\zeta \in \mathbb{T}^*$ . The following properties hold.

- (i) If  $\eta$  is conformal fractional differentiable of order  $\alpha$  at  $\zeta > 0$ , then  $\eta$  is continuous at  $\zeta$ .
- (ii) If  $\eta$  is continuous at  $\zeta$  and  $\zeta$  is right-scattered, then  $\eta$  is conformable fractional differentiable of order  $\alpha$  at  $\zeta$  with

$$T_\alpha^\Delta(\eta)(\zeta) = \frac{\eta(\sigma(\zeta)) - \eta(\zeta)}{\mu(\zeta)} \zeta^{1-\alpha}.$$

- (iii) If  $\zeta$  is right-dense, then  $\eta$  is conformable fractional differentiable of order  $\alpha$  at  $\zeta$  if, and only if, the limit  $\lim_{s \rightarrow \zeta} \frac{\eta(\zeta) - \eta(s)}{\zeta - s} \zeta^{1-\alpha}$  exists as a finite number. In this case,

$$T_\alpha^\Delta(\eta)(\zeta) = \lim_{s \rightarrow \zeta} \frac{\eta(\zeta) - \eta(s)}{\zeta - s} \zeta^{1-\alpha}.$$

- (iv) If  $\eta$  is fractional differentiable of order  $\alpha$  at  $\zeta$ , then

$$\eta(\sigma(\zeta)) = \eta(\zeta) + (\mu(\zeta)\zeta^{\alpha-1} T_\alpha^\Delta(\eta)(\zeta)).$$

**Remark 2.3.** In a time scale  $\mathbb{T}$ , due to the inherited topology of the real numbers, a function  $\eta$  is always continuous at any isolated point  $\zeta \in \mathbb{T}$ .

**Example 2.1.** Let  $h > 0$  and  $\mathbb{T} = h\mathbb{Z} = \{hk : k \in \mathbb{Z}\}$ . Then  $\sigma(\zeta) = \zeta + h$  and  $\mu(\zeta) = h$  for all  $\zeta \in \mathbb{T}$ . For function  $\eta : \mathbb{T} \rightarrow \mathbb{R}$ ,  $\eta(\zeta) = \zeta^2$  we have  $T_\alpha^\Delta(\zeta^2) = (2\zeta + h)\zeta^{1-\alpha}$ .

**Example 2.2.** Let  $q > 1$  and  $\mathbb{T} = q\mathbb{Z} := q\mathbb{Z} \cup \{0\}$  with  $q\mathbb{Z} := \{q^k : k \in \mathbb{Z}\}$ . In this time scale

$$\sigma(\zeta) = \begin{cases} q\zeta, & \text{if } \zeta \neq 0; \\ 0, & \text{if } \zeta = 0; \end{cases} \quad \text{and} \quad \mu(\zeta) = \begin{cases} (q-1)\zeta, & \text{if } \zeta \neq 0; \\ 0, & \text{if } \zeta = 0. \end{cases}$$

Here 0 is a right-dense minimum and every other point in  $\mathbb{T}$  is isolated. Now consider the square function of Example 2.1. It follows that

$$T_\alpha^\Delta(\eta)(\zeta) = T_\alpha^\Delta(\zeta^2) = \begin{cases} (q+1)\zeta^{2-\alpha}, & \text{if } \zeta \neq 0; \\ 0, & \text{if } \zeta = 0. \end{cases}$$

The conformable fractional derivative has the following properties

**Theorem 2.2.** Let  $f, g : \mathbb{T} \rightarrow \mathbb{R}$  be conformable fractional derivative of order  $\alpha \in (0, 1]$ , the following properties are hold:

(i) The  $f, g : \mathbb{T} \rightarrow \mathbb{R}$  are  $\alpha$ -conformable fractional derivative and

$$T_\alpha^\Delta(f + g) = T_\alpha^\Delta(f) + T_\alpha^\Delta(g).$$

(ii) For all  $k \in \mathbb{R}$ , then  $kf : \mathbb{T} \rightarrow \mathbb{R}$   $\alpha$ -conformable fractional derivative and

$$T_\alpha^\Delta(kf) = kT_\alpha^\Delta(f).$$

(iii) If  $f$  and  $g$  are  $\alpha$ -conformable fractional differentiable, we have  $fg : \mathbb{T} \rightarrow \mathbb{R}$  is  $\alpha$ -conformable fractional differentiable and

$$T_\alpha^\Delta(fg) = T_\alpha^\Delta(f)g + f^\sigma T_\alpha^\Delta(g).$$

(iv) If  $f$  and  $g$  are  $\alpha$ -conformable fractional differentiable, then  $f/g$  is  $\alpha$ -conformable fractional differentiable with

$$T_\alpha^\Delta\left(\frac{f}{g}\right) = \frac{T_\alpha^\Delta(f)g - fT_\alpha^\Delta(g)}{gg^\sigma}$$

valid  $\forall \zeta \in \mathbb{T}^*$ , where  $gg^\sigma \neq 0$ .

**Theorem 2.3.** Let  $\varsigma : \mathbb{T} \rightarrow \mathbb{R}$  be continuous and  $\alpha$ -fractional differentiable at  $\zeta \in \mathbb{T}$  for  $\alpha \in (0, 1]$  and  $\delta : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous differentiable function. Then there is  $c$  in the interval  $[\zeta, \sigma(\zeta)]$  such that

$$T_\alpha^\Delta(\delta \circ \varsigma) = \delta'(\varsigma(c))T_\alpha^\Delta(\varsigma(\zeta)). \quad (2.3)$$

**Theorem 2.4.** Let  $\delta : \mathbb{R} \rightarrow \mathbb{R}$  be continuously differentiable,  $\alpha \in (0, 1]$  and  $\varsigma : \mathbb{T} \rightarrow \mathbb{R}$  be  $\alpha$ -fractional differentiable function. Then  $(\delta \circ \varsigma) : \mathbb{T} \rightarrow \mathbb{R}$  is  $\alpha$ -fractional differentiable and we have

$$T_\alpha^\Delta(\delta \circ \varsigma)(s) = \left\{ \int_0^1 \delta'(\varsigma(s) + h\mu(s)s^{\alpha-1}T_\alpha^\Delta(\varsigma(s)))dh \right\} T_\alpha^\Delta(\varsigma(s)). \quad (2.4)$$

**Definition 2.2.** Let  $0 < \alpha \leq 1$ , the  $\alpha$ -fractional of  $f$ , is defined as

$$\int f(s)\Delta_\alpha s = \int f(s)s^{\alpha-1}\Delta s.$$

The conformable fractional integral satisfying the next properties

**Theorem 2.5.** Assume  $a, b, c \in \mathbb{T}$ ,  $\lambda \in \mathbb{R}$ . Let  $\delta, \varsigma : \mathbb{T} \rightarrow \mathbb{R}$ . Then

- (i)  $\int_a^b [\varsigma(s) + \delta(s)]\Delta_\alpha s = \int_a^b \varsigma(s)\Delta_\alpha s + \int_a^b \delta(s)\Delta_\alpha s.$
- (ii)  $\int_a^b \lambda \varsigma(s)\Delta_\alpha s = \lambda \int_a^b \varsigma(s)\Delta_\alpha s.$
- (iii)  $\int_a^b \varsigma(s)\Delta_\alpha s = - \int_b^a \varsigma(s)\Delta_\alpha s.$
- (iv)  $\int_a^b \varsigma(s)\Delta_\alpha s = \int_a^c \varsigma(s)\Delta_\alpha s + \int_c^b \varsigma(s)\Delta_\alpha s.$
- (v)  $\int_a^a \varsigma(s)\Delta_\alpha s = 0.$

**Theorem 2.6.** (Dynamic Hölder's Inequality [11]) Let  $u, v \in \mathbb{T}$  with  $u < v$ . If  $\vartheta, \theta \in CC_{rd}^1([u, v]_{\mathbb{T}} \times [u, v]_{\mathbb{T}}, \mathbb{R})$  be integrable functions and  $\frac{1}{v} + \frac{1}{\varpi} = 1$  with  $v > 1$ . Then

$$\begin{aligned} \int_u^v \int_u^v |\vartheta(r, \delta) \theta(r, \delta)| \Delta^\alpha r \Delta^\alpha \delta &\leq \left[ \int_u^v \int_u^v |\vartheta(r, \delta)|^v \Delta^\alpha r \Delta^\alpha \delta \right]^{\frac{1}{v}} \\ &\times \left[ \int_u^v \int_u^v |\theta(r, \delta)|^\varpi \Delta^\alpha r \Delta^\alpha \delta \right]^{\frac{1}{\varpi}}. \end{aligned} \quad (2.5)$$

This inequality is reversed if  $0 < v < 1$  and if  $v < 0$  or  $\varpi < 0$ .

### 3. Main results

**Theorem 3.1.** Let  $\mathbb{T}$  be a time scale with  $\delta_0, \iota_i, \mathfrak{J}_i, \delta_i \in \mathbb{T}$ ,  $(i = 1, \dots, n)$ . Let  $h_i \geq 1$ ,  $\nu_i, \varpi_i > 1$  be constants and  $\frac{1}{\nu_i} + \frac{1}{\varpi_i} = 1$ . Let  $\Delta^\alpha$ -differentiable functions  $\vartheta_i(\mathfrak{J}_i)$  be decreasing on  $[\delta_0, \iota_i]_{\mathbb{T}}$ , where  $\iota_i \in (0, \infty)$ . Suppose  $\vartheta_i(\delta_0) = 0$ . Then

$$\begin{aligned} &\int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\iota_2} \dots \int_{\delta_0}^{\iota_n} \frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{J}_i)|}{\left( \sum_{i=1}^n \frac{(\mathfrak{J}_i - \delta_0)}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \Delta^\alpha \mathfrak{J}_n \Delta^\alpha \mathfrak{J}_{n-1} \dots \Delta^\alpha \mathfrak{J}_1 \\ &\leq K \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} (\sigma(\iota_i) - \sigma(\mathfrak{J}_i)) |\vartheta_i^{h_i-1}(\mathfrak{J}_i) \vartheta_i^{\Delta^\alpha}(\mathfrak{J}_i)|^{\nu_i} \Delta^\alpha \mathfrak{J}_i \right)^{\frac{1}{\nu_i}}, \end{aligned} \quad (3.1)$$

where

$$K = K(\iota_1, \dots, \iota_n) = \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n h_i (\iota_i - \delta_0)^{\frac{1}{\varpi_i}}.$$

*Proof.* From Hölder inequality (2.5), one can see that

$$\begin{aligned} \prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{J}_i)| &\leq \prod_{i=1}^n h_i \int_{\delta_0}^{\mathfrak{J}_i} |\vartheta_i^{h_i-1}(\tau_i) \vartheta_i^{\Delta^\alpha}(\tau_i)| \Delta^\alpha \tau_i \\ &\leq \prod_{i=1}^n h_i (\mathfrak{J}_i - \delta_0)^{\frac{1}{\varpi_i}} \left( \int_{\delta_0}^{\mathfrak{J}_i} |\vartheta_i^{h_i-1}(\tau_i) \vartheta_i^{\Delta^\alpha}(\tau_i)|^{\nu_i} \Delta^\alpha \tau_i \right)^{\frac{1}{\nu_i}}. \end{aligned} \quad (3.2)$$

Using the inequality for the means [25],

$$\left( \prod_{i=1}^n \lambda_i^{\frac{1}{\varpi_i}} \right)^{\frac{1}{\sum_{i=1}^n \frac{1}{\varpi_i}}} \leq \frac{1}{\sum_{i=1}^n \frac{1}{\varpi_i}} \sum_{i=1}^n \frac{\lambda_i}{\varpi_i}, \quad \lambda_i > 0 \quad (i = 1, \dots, n), \quad (3.3)$$

we have

$$\begin{aligned} &\frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{J}_i)|}{\left( \sum_{i=1}^n \frac{(\mathfrak{J}_i - \delta_0)}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \\ &\leq \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n h_i \left( \int_{\delta_0}^{\mathfrak{J}_i} |\vartheta_i^{h_i-1}(\tau_i) \vartheta_i^{\Delta^\alpha}(\tau_i)|^{\nu_i} \Delta^\alpha \tau_i \right)^{\frac{1}{\nu_i}}. \end{aligned} \quad (3.4)$$

Using the integration of (3.4) on  $\mathfrak{J}_i$  from  $\delta_0$  to  $\iota_i$  ( $i = 1, \dots, n$ ) employing the inequality of Hölder's, gets

$$\begin{aligned}
& \int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\iota_2} \dots \int_{\delta_0}^{\iota_n} \frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{J}_i)|}{\left(\sum_{i=1}^n \frac{(\mathfrak{J}_i - \delta_0)}{\varpi_i}\right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \Delta^\alpha \mathfrak{J}_n \Delta^\alpha \mathfrak{J}_{n-1} \dots \Delta^\alpha \mathfrak{J}_1 \\
& \leq \left(n - \sum_{i=1}^n \frac{1}{\nu_i}\right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n h_i \int_{\delta_0}^{\iota_i} \left( \int_{\delta_0}^{\mathfrak{J}_i} |\vartheta_i^{h_i-1}(\tau_i) \vartheta_i^{\Delta^\alpha}(\tau_i)|^{\nu_i} \Delta^\alpha \tau_i \Delta^\alpha \mathfrak{J}_i \right)^{\frac{1}{\nu_i}} \\
& \leq K \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} \int_{\delta_0}^{\mathfrak{J}_i} |\vartheta_i^{h_i-1}(\tau_i) \vartheta_i^{\Delta^\alpha}(\tau_i)|^{\nu_i} \Delta^\alpha \tau_i \Delta^\alpha \mathfrak{J}_i \right)^{\frac{1}{\nu_i}} \\
& = K \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} (\iota_i - \mathfrak{J}_i) |\vartheta_i^{h_i-1}(\mathfrak{J}_i) \vartheta_i^{\Delta^\alpha}(\mathfrak{J}_i)|^{\nu_i} \Delta^\alpha \mathfrak{J}_i \right)^{\frac{1}{\nu_i}}. \tag{3.5}
\end{aligned}$$

By exploiting the fact  $\iota_i \leq \sigma(\iota_i)$ , we get that

$$\begin{aligned}
& \int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\iota_2} \dots \int_{\delta_0}^{\iota_n} \frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{J}_i)|}{\left(\sum_{i=1}^n \frac{(\mathfrak{J}_i - \delta_0)}{\varpi_i}\right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \Delta^\alpha \mathfrak{J}_n \Delta^\alpha \mathfrak{J}_{n-1} \dots \Delta^\alpha \mathfrak{J}_1 \\
& \leq K \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} (\sigma(\iota_i) - \sigma(\mathfrak{J}_i)) |\vartheta_i^{h_i-1}(\mathfrak{J}_i) \vartheta_i^{\Delta^\alpha}(\mathfrak{J}_i)|^{\nu_i} \Delta^\alpha \mathfrak{J}_i \right)^{\frac{1}{\nu_i}}.
\end{aligned}$$

This concludes the evidence.  $\square$

**Remark 3.1.** In Theorem 3.1, taking  $\mathbb{T} = \mathbb{Z}, \alpha = 1, h_i = 1$ , got results thanks to the authors of [3, Theorem 1.1].

**Remark 3.2.** In Theorem 3.1, taking  $\mathbb{T} = \mathbb{R}, \alpha = 1$ , got results thanks to the authors of [3, Theorem 1.3].

**Corollary 3.1.** In Theorem 3.1, taking  $n = 2$ , and  $h_1 = h_2 = 1$ , if  $\nu_1, \nu_2 > 1$  are such that  $\frac{1}{\nu_1} + \frac{1}{\nu_2} \geq 1$  and  $0 < \lambda = 2 - \frac{1}{\nu_1} - \frac{1}{\nu_2} = \frac{1}{\varpi_1} + \frac{1}{\varpi_2} \leq 1$ , inequality (3.1) reduces to

$$\begin{aligned}
& \int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\iota_2} \frac{|\vartheta_1(\mathfrak{J}_1)| |\vartheta_2(\mathfrak{J}_2)|}{\left(\varpi_2(\mathfrak{J}_1 - \delta_0) + \varpi_1(\mathfrak{J}_2 - \delta_0)\right)^\lambda} \Delta^\alpha \mathfrak{J}_2 \Delta^\alpha \mathfrak{J}_1 \leq \frac{1}{(\lambda \varpi_1 \varpi_2)^\lambda} (\iota_1 - \delta_0)^{\frac{1}{\varpi_1}} (\iota_2 - \delta_0)^{\frac{1}{\varpi_2}} \tag{3.6} \\
& \times \left( \int_{\delta_0}^{\iota_1} (\sigma(\iota_1) - \sigma(\mathfrak{J}_1)) |\vartheta_1^{\Delta^\alpha}(\mathfrak{J}_1)|^{\nu_1} \Delta^\alpha \mathfrak{J}_1 \right)^{\frac{1}{\nu_1}} \left( \int_{\delta_0}^{\iota_2} (\sigma(\iota_2) - \sigma(\mathfrak{J}_2)) |\vartheta_2^{\Delta^\alpha}(\mathfrak{J}_2)|^{\nu_2} \Delta^\alpha \mathfrak{J}_2 \right)^{\frac{1}{\nu_2}}.
\end{aligned}$$

**Remark 3.3.** In special case, taking  $\mathbb{T} = \mathbb{R}, \alpha = 1$ , in (3.6), we have that

$$\begin{aligned}
& \int_0^{\iota_1} \int_0^{\iota_2} \frac{|\vartheta_1(\mathfrak{J}_1)| |\vartheta_2(\mathfrak{J}_2)|}{(\varpi_2 \mathfrak{J}_1 + \varpi_1 \mathfrak{J}_2)^\lambda} d\mathfrak{J}_2 d\mathfrak{J}_1 \leq \frac{1}{(\lambda \varpi_1 \varpi_2)^\lambda} (\iota_1)^{\frac{1}{\varpi_1}} (\iota_2)^{\frac{1}{\varpi_2}} \\
& \times \left( \int_0^{\iota_1} (\iota_1 - \mathfrak{J}_1) |\vartheta_1'(\mathfrak{J}_1)|^{\nu_1} d\mathfrak{J}_1 \right)^{\frac{1}{\nu_1}} \left( \int_0^{\iota_2} (\iota_2 - \mathfrak{J}_2) |\vartheta_2'(\mathfrak{J}_2)|^{\nu_2} d\mathfrak{J}_2 \right)^{\frac{1}{\nu_2}}, \tag{3.7}
\end{aligned}$$

which is an interesting variation of inequality (1.2).

**Remark 3.4.** In special case, taking  $\mathbb{T} = \mathbb{Z}, \alpha = 1$ , in (3.6), we have that

$$\sum_{\mathfrak{I}_1=1}^{m_1} \sum_{\mathfrak{I}_2=1}^{m_2} \frac{|a_1(\mathfrak{I}_1)| |a_2(\mathfrak{I}_2)|}{(\varpi_2 \mathfrak{I}_1 + \varpi_1 \mathfrak{I}_2)^\lambda} \leq \frac{1}{(\lambda \varpi_1 \varpi_2)^\lambda} (m_1)^{\frac{1}{\varpi_1}} (m_2)^{\frac{1}{\varpi_2}} \\ \times \left( \sum_{\mathfrak{I}_1=1}^{m_1} (m_1 - \mathfrak{I}_1 + 1) |\Delta^\alpha a_1(\mathfrak{I}_1)|^{\nu_1} \right)^{\frac{1}{\nu_1}} \left( \sum_{\mathfrak{I}_2=1}^{m_2} (m_2 - \mathfrak{I}_2 + 1) |\Delta^\alpha a_2(\mathfrak{I}_2)|^{\nu_2} \right)^{\frac{1}{\nu_2}}, \quad (3.8)$$

which is an interesting variation of inequality (1.1).

**Corollary 3.2.** In Corollary 3.1, if  $\lambda = 1$ , then  $\frac{1}{\nu_1} + \frac{1}{\nu_2} = \frac{1}{\varpi_1} + \frac{1}{\varpi_2} = 1$  and take  $\nu_1 = \varpi_2, \nu_2 = \varpi_1$ . In this case inequality (3.6) reduces to

$$\int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\iota_2} \frac{|\vartheta_1(\mathfrak{I}_1)| |\vartheta_2(\mathfrak{I}_2)|}{\varpi_2(\mathfrak{I}_1 - \delta_0) + \varpi_1(\mathfrak{I}_2 - \delta_0)} \Delta^\alpha \mathfrak{I}_2 \Delta^\alpha \mathfrak{I}_1 \leq \frac{1}{\nu_1 \varpi_1} (\iota_1 - \delta_0)^{\frac{\nu_1-1}{\nu_1}} (\iota_2 - \delta_0)^{\frac{\varpi_1-1}{\varpi_1}} \\ \times \left( \int_{\delta_0}^{\iota_1} (\sigma(\iota_1) - \sigma(\mathfrak{I}_1)) |\vartheta_1^{\Delta^\alpha}(\mathfrak{I}_1)|^{\nu_1} \Delta^\alpha \mathfrak{I}_1 \right)^{\frac{1}{\nu_1}} \left( \int_{\delta_0}^{\iota_2} (\sigma(\iota_2) - \sigma(\mathfrak{I}_2)) |\vartheta_2^{\Delta^\alpha}(\mathfrak{I}_2)|^{\varpi_1} \Delta^\alpha \mathfrak{I}_2 \right)^{\frac{1}{\varpi_1}}. \quad (3.9)$$

**Remark 3.5.** In Corollary 3.2, if  $\mathbb{T} = \mathbb{R}, \alpha = 1$  we obtain an equivalent formulation of the inequality that Pachpatte presented in [2, Theorem 2].

**Remark 3.6.** In Corollary 3.2, if  $\mathbb{T} = \mathbb{Z}, \alpha = 1$  we obtain an equivalent formulation of the inequality that Pachpatte presented in [2, Theorem 1].

**Theorem 3.2.** Let  $\mathbb{T}$  be a time scale with  $\delta_0, \iota_i, \varsigma_i, \mathfrak{I}_i, \delta_i \in \mathbb{T}, (i = 1, \dots, n)$ . Let  $h_i \geq 1, \nu_i, \varpi_i > 1$  be constants and  $\frac{1}{\nu_i} + \frac{1}{\varpi_i} = 1$ . Let the  $\Delta^\alpha$ -differentiable fun.  $\vartheta_i(\mathfrak{I}_i, \delta_i)$  be decreasing funs. on  $[\delta_0, \iota_i]_{\mathbb{T}} \times [\delta_0, \varsigma_i]_{\mathbb{T}}$  and  $\vartheta_i(\delta_0, \delta_i) = \vartheta_i(\mathfrak{I}_i, \delta_0) = 0$ , for  $(i = 1, \dots, n)$ . Partial derivatives of  $\vartheta_i$  are indicated by  $\vartheta_i^{\Delta^\alpha_1}, \vartheta_i^{\Delta^\alpha_2}, \vartheta_i^{\Delta^\alpha_{12}} = \vartheta_i^{\Delta^\alpha_{21}}$ . Let

$$(\vartheta_i^{h_i}(\mathfrak{I}_i, \delta_i))^{\Delta^\alpha_1 \Delta^\alpha_2} \leq (h_i \vartheta_i^{h_i-1}(\mathfrak{I}_i, \delta_i) \cdot \vartheta_i^{\Delta^\alpha_1}(\mathfrak{I}_i, \delta_i))^{\Delta^\alpha_2} = \vartheta_i^{\Delta^\alpha_{12}}(\mathfrak{I}_i, \delta_i).$$

Then

$$\int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\varsigma_1} \cdots \int_{\delta_0}^{\iota_n} \int_{\delta_0}^{\varsigma_n} \frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{I}_i, \delta_i)|}{\left( \sum_{i=1}^n \frac{(\mathfrak{I}_i - \delta_0)(\delta_i - \delta_0)}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \Delta^\alpha \delta_n \Delta^\alpha \mathfrak{I}_n \cdots \Delta^\alpha \delta_1 \Delta^\alpha \mathfrak{I}_1 \\ \leq C \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} \int_{\delta_0}^{\varsigma_i} (\sigma(\iota_i) - \sigma(\mathfrak{I}_i)) (\sigma(\varsigma_i) - \sigma(\delta_i)) |\vartheta_i^{\Delta^\alpha_{12}}(\mathfrak{I}_i, \delta_i)|^{\nu_i} \Delta^\alpha \delta_i \Delta^\alpha \mathfrak{I}_i \right)^{\frac{1}{\nu_i}}, \quad (3.10)$$

where

$$C = C(\iota_1 \varsigma_1, \dots, \iota_n \varsigma_n) = \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n [(\iota_i - \delta_0)(\varsigma_i - \delta_0)]^{\frac{1}{\varpi_i}}.$$

*Proof.* We can type

$$\begin{aligned} \vartheta_i^{h_i}(\mathfrak{I}_i, \delta_i) &= \vartheta_i^{h_i}(\mathfrak{I}_i, \delta_i) - \vartheta_i^{h_i}(\delta_0, \delta_i) - \vartheta_i^{h_i}(\mathfrak{I}_i, \delta_0) + \vartheta_i^{h_i}(\delta_0, \delta_0) \\ &= \int_{\delta_0}^{\mathfrak{I}_i} (\vartheta_i^{h_i}(\xi_i, \delta_i))^{\Delta^\alpha_1} \Delta^\alpha_1 \xi_i - \int_{\delta_0}^{\mathfrak{I}_i} (\vartheta_i^{h_i}(\xi_i, \delta_0))^{\Delta^\alpha_1} \Delta^\alpha_1 \xi_i \end{aligned}$$

$$\begin{aligned}
&= \int_{\delta_0}^{\mathfrak{I}_i} [(\vartheta_i^{h_i}(\xi_i, \delta_i))^{\Delta_1^\alpha} - (\vartheta_i^{h_i}(\xi_i, \delta_0))^{\Delta_1^\alpha}] \Delta^\alpha \xi_i \\
&\leq \int_{\delta_0}^{\mathfrak{I}_i} \int_{\delta_0}^{\delta_i} (h_i \vartheta_i^{h_i-1}(\xi_i, \eta_i) \cdot \vartheta_i^{\Delta_1^\alpha}(\xi_i, \eta_i))^{\Delta_2^\alpha} \Delta^\alpha \eta_i \Delta^\alpha \xi_i \\
&= \int_{\delta_0}^{\mathfrak{I}_i} \int_{\delta_0}^{\delta_i} \vartheta_i^{\Delta_{12}^\alpha}(\xi_i, \eta_i) \Delta^\alpha \eta_i \Delta^\alpha \xi_i.
\end{aligned} \tag{3.11}$$

From (3.11) applying (2.5) and (3.12), gets

$$\begin{aligned}
\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{I}_i, \delta_i)| &\leq \prod_{i=1}^n \int_{\delta_0}^{\mathfrak{I}_i} \int_{\delta_0}^{\delta_i} |\vartheta_i^{\Delta_{12}^\alpha}(\xi_i, \eta_i)| \Delta_1^\alpha \eta_i \Delta_2^\alpha \xi_i \\
&\leq \prod_{i=1}^n [(\mathfrak{I}_i - \delta_0)(\delta_i - \delta_0)]^{\frac{1}{\varpi_i}} \left( \int_{\delta_0}^{\mathfrak{I}_i} \int_{\delta_0}^{\delta_i} |\vartheta_i^{\Delta_{12}^\alpha}(\xi_i, \eta_i)|^{\nu_i} \Delta^\alpha \eta_i \Delta^\alpha \xi_i \right)^{\frac{1}{\nu_i}}.
\end{aligned} \tag{3.12}$$

Using inequality (3.3), we get that

$$\frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{I}_i, \delta_i)|}{\left( \sum_{i=1}^n \frac{(\mathfrak{I}_i - \delta_0)(\delta_i - \delta_0)}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \leq \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n \left( \int_{\delta_0}^{\mathfrak{I}_i} \int_{\delta_0}^{\delta_i} |\vartheta_i^{\Delta_{12}^\alpha}(\xi_i, \eta_i)|^{\nu_i} \Delta^\alpha \eta_i \Delta^\alpha \xi_i \right)^{\frac{1}{\nu_i}}. \tag{3.13}$$

Integrating (3.13) on  $\mathfrak{I}_i$  and  $\delta_i$ , applying (2.5) and Fubini's theorem, yields

$$\begin{aligned}
&\int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\mathfrak{I}_1} \cdots \int_{\delta_0}^{\iota_n} \int_{\delta_0}^{\mathfrak{I}_n} \frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{I}_i, \delta_i)|}{\left( \sum_{i=1}^n \frac{(\mathfrak{I}_i - \delta_0)(\delta_i - \delta_0)}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \Delta^\alpha \delta_n \Delta^\alpha \mathfrak{I}_n \cdots \Delta^\alpha \delta_1 \Delta^\alpha \mathfrak{I}_1 \\
&\leq \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \\
&\quad \times \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} \int_{\delta_0}^{\mathfrak{I}_i} \left( \int_{\delta_0}^{\mathfrak{I}_i} \int_{\delta_0}^{\delta_i} |\vartheta_i^{\Delta_{12}^\alpha}(\xi_i, \eta_i)|^{\nu_i} \Delta^\alpha \eta_i \Delta^\alpha \xi_i \right)^{\frac{1}{\nu_i}} \Delta^\alpha \delta_i \Delta^\alpha \mathfrak{I}_i \right) \\
&\leq \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \\
&\quad \times \prod_{i=1}^n \left[ (\iota_i - \delta_0)(\mathfrak{I}_i - \delta_0) \right]^{\frac{1}{\varpi_i}} \left( \int_{\delta_0}^{\iota_i} \int_{\delta_0}^{\mathfrak{I}_i} \left( \int_{\delta_0}^{\mathfrak{I}_i} \int_{\delta_0}^{\delta_i} |\vartheta_i^{\Delta_{12}^\alpha}(\xi_i, \eta_i)|^{\nu_i} \Delta^\alpha \eta_i \Delta^\alpha \xi_i \right) \Delta^\alpha \delta_i \Delta^\alpha \mathfrak{I}_i \right)^{\frac{1}{\nu_i}} \\
&= C \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} \int_{\delta_0}^{\mathfrak{I}_i} (\iota_i - \mathfrak{I}_i)(\mathfrak{I}_i - \delta_i) |\vartheta_i^{\Delta_{12}^\alpha}(\mathfrak{I}_i, \delta_i)|^{\nu_i} \Delta^\alpha \delta_i \Delta^\alpha \mathfrak{I}_i \right)^{\frac{1}{\nu_i}}.
\end{aligned} \tag{3.14}$$

By exploiting the fact  $\iota_i \leq \sigma(\iota_i)$ , gives

$$\int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\mathfrak{I}_1} \cdots \int_{\delta_0}^{\iota_n} \int_{\delta_0}^{\mathfrak{I}_n} \frac{\prod_{i=1}^n |\vartheta_i^{h_i}(\mathfrak{I}_i, \delta_i)|}{\left( \sum_{i=1}^n \frac{(\mathfrak{I}_i - \delta_0)(\delta_i - \delta_0)}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \Delta^\alpha \delta_n \Delta^\alpha \mathfrak{I}_n \cdots \Delta^\alpha \delta_1 \Delta^\alpha \mathfrak{I}_1$$

$$\leq C \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} \int_{\delta_0}^{\varsigma_i} (\sigma(\iota_i) - \sigma(\mathfrak{J}_i))(\sigma(\varsigma_i) - \sigma(\delta_i)) |\vartheta_i^{\Delta_{12}^\alpha}(\mathfrak{J}_i, \delta_i)|^{\nu_i} \Delta^\alpha \delta_i \Delta^\alpha \mathfrak{J}_i \right)^{\frac{1}{\nu_i}}.$$

This concludes the evidence.  $\square$

**Remark 3.7.** In Theorem 3.2, if we take  $\mathbb{T} = \mathbb{Z}, \alpha = 1, h_i = 1$ , got results thanks to the authors [3, Theorem 1.2].

**Remark 3.8.** In Theorem 3.2, supposing  $\mathbb{T} = \mathbb{R}, \alpha = 1$ , got results thanks the authors [3, Theorem 1.4].

**Corollary 3.3.** Taking  $n = 2$  and  $h_1 = h_2 = 1$  in Theorem 3.2, we have

$$\vartheta_1^{\Delta_{12}^\alpha}(\mathfrak{J}_1, \delta_1) = \vartheta^{\Delta_1^\alpha \Delta_2^\alpha}(\mathfrak{J}_1, \delta_1), \quad \vartheta_2^{\Delta_{12}^\alpha}(\mathfrak{J}_1, \delta_1) = \vartheta^{\Delta_2^\alpha \Delta_1^\alpha}(\mathfrak{J}_2, \delta_2).$$

Moreover, if  $\nu_1, \nu_2 > 1$  satisfy  $\frac{1}{\nu_1} + \frac{1}{\nu_2} \geq 1$  and  $0 < \lambda = 2 - \frac{1}{\nu_1} - \frac{1}{\nu_2} = \frac{1}{\varpi_1} + \frac{1}{\varpi_2} \leq 1$ , inequality (3.10) reduces to

$$\begin{aligned} & \int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\varsigma_1} \left( \int_{\delta_0}^{\iota_2} \int_{\delta_0}^{\varsigma_2} \frac{|\vartheta_1(\mathfrak{J}_1, \delta_1)| |\vartheta_2(\mathfrak{J}_2, \delta_2)|}{(\nu_1(\mathfrak{J}_1 - \delta_0)(\delta_1 - \delta_0) + \varpi_1(\mathfrak{J}_2 - \delta_0)(\delta_2 - \delta_0))^\lambda} \Delta^\alpha \mathfrak{J}_2 \Delta^\alpha \delta_2 \right) \Delta^\alpha \mathfrak{J}_1 \Delta^\alpha \delta_1 \\ & \leq \frac{1}{(\lambda \varpi_1 \varpi_2)^\lambda} \left[ (\iota_1 - \delta_0)(\varsigma_1 - \delta_0) \right]^{\frac{1}{\varpi_1}} \left[ (\iota_2 - \delta_0)(\varsigma_2 - \delta_0) \right]^{\frac{\varpi_1-1}{\varpi_1}} \\ & \quad \times \left( \int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\varsigma_1} (\sigma(\iota_1) - \sigma(\mathfrak{J}_1))(\sigma(\varsigma_1) - \sigma(\delta_1)) |\vartheta_1^{\Delta_1^\alpha \Delta_2^\alpha}(\mathfrak{J}_1, \delta_1)|^{\nu_1} \Delta^\alpha \mathfrak{J}_1 \Delta^\alpha \delta_1 \right)^{\frac{1}{\nu_1}} \\ & \quad \left( \int_{\delta_0}^{\iota_2} \int_{\delta_0}^{\varsigma_2} (\sigma(\iota_2) - \delta_0)(\sigma(\varsigma_2) - \delta_0) |\vartheta_2^{\Delta_2^\alpha \Delta_1^\alpha}(\mathfrak{J}_2, \delta_2)|^{\nu_2} \Delta^\alpha \mathfrak{J}_2 \Delta^\alpha \delta_2 \right)^{\frac{1}{\nu_2}}. \end{aligned} \quad (3.15)$$

**Remark 3.9.** In a unique scenario, if we take  $\mathbb{T} = \mathbb{R}$  in Corollary 3.3, the inequality (3.15) reduces to

$$\begin{aligned} & \int_0^{\iota_1} \int_0^{\varsigma_1} \left( \int_0^{\iota_2} \int_0^{\varsigma_2} \frac{|\vartheta_1(\mathfrak{J}_1, \delta_1)| |\vartheta_2(\mathfrak{J}_2, \delta_2)|}{(\nu_1 \mathfrak{J}_1 \delta_1 + \varpi_1 \mathfrak{J}_2 \delta_2)^\lambda} d\mathfrak{J}_2 d\delta_2 \right) d\mathfrak{J}_1 d\delta_1 \\ & \leq \frac{1}{(\lambda \varpi_1 \varpi_2)^\lambda} [\iota_1 \varsigma_1]^{\frac{1}{\varpi_1}} [\iota_2 \varsigma_2]^{\frac{\varpi_1-1}{\varpi_1}} \\ & \quad \times \left( \int_0^{\iota_1} \int_0^{\varsigma_1} (\iota_1 - \mathfrak{J}_1)(\varsigma_1 - \delta_1) |D_1 D_2 \vartheta_1(\mathfrak{J}_1, \delta_1)|^{\nu_1} d\mathfrak{J}_1 d\delta_1 \right)^{\frac{1}{\nu_1}} \\ & \quad \times \left( \int_0^{\iota_2} \int_0^{\varsigma_2} (\iota_2 - \mathfrak{J}_2)(\varsigma_2 - \delta_2) |D_1 D_2 \vartheta_2(\mathfrak{J}_2, \delta_2)|^{\nu_2} d\mathfrak{J}_2 d\delta_2 \right)^{\frac{1}{\nu_2}}. \end{aligned} \quad (3.16)$$

**Remark 3.10.** In a unique scenario, if we take  $\mathbb{T} = \mathbb{Z}$  in Corollary 3.3, the inequality (3.15) reduces to

$$\begin{aligned} & \sum_{\mathfrak{J}_1=1}^{m_1} \sum_{\delta_1=1}^{n_1} \left( \sum_{\mathfrak{J}_2=1}^{m_2} \sum_{\delta_2=1}^{n_2} \frac{|a_1(\mathfrak{J}_1, \delta_1)| |a_2(\mathfrak{J}_2, \delta_2)|}{(\nu_1 \mathfrak{J}_1 \delta_1 + \varpi_1 \mathfrak{J}_2 \delta_2)^\lambda} \right) \\ & \leq \frac{1}{(\lambda \varpi_1 \varpi_2)^\lambda} [m_1 n_1]^{\frac{1}{\varpi_1}} [m_2 n_2]^{\frac{\varpi_1-1}{\varpi_1}} \end{aligned}$$

$$\begin{aligned} & \times \left( \sum_{\mathfrak{I}_1=1}^{m_1} \sum_{\delta_1=1}^{n_1} (n_1 - \delta_1)(m_1 - \mathfrak{I}_1) |\Delta_1^\alpha \Delta_2^\alpha a_1(\mathfrak{I}_1, \delta_1)|^{\nu_1} \right)^{\frac{1}{\nu_1}} \\ & \times \left( \sum_{\mathfrak{I}_2=1}^{m_2} \sum_{\delta_2=1}^{n_2} (n_2 - \delta_2)(m_2 - \mathfrak{I}_2) |\Delta_1^\alpha \Delta_2^\alpha a_2(\mathfrak{I}_2, \delta_2)|^{\nu_2} \right)^{\frac{1}{\nu_2}}. \end{aligned} \quad (3.17)$$

**Corollary 3.4.** In Corollary 3.3, if  $\lambda = 1$ , then  $\frac{1}{\nu_1} + \frac{1}{\nu_2} = \frac{1}{\varpi_1} + \frac{1}{\varpi_2} = 1$  and take  $\nu_1 = \varpi_2$ ,  $\nu_2 = \varpi_1$ . In this case the inequality (3.15) reduces to

$$\begin{aligned} & \int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\mathfrak{I}_1} \left( \int_{\delta_0}^{\iota_2} \int_{\delta_0}^{\mathfrak{I}_2} \frac{|\vartheta_1(\mathfrak{I}_1, \delta_1)| |\vartheta_2(\mathfrak{I}_2, \delta_2)|}{(\nu_1(\mathfrak{I}_1 - \delta_0)(\delta_1 - \delta_0) + \varpi_1(\mathfrak{I}_2 - \delta_0)(\delta_2 - \delta_0))} \Delta^\alpha \mathfrak{I}_2 \Delta^\alpha \delta_2 \right) \Delta^\alpha \mathfrak{I}_1 \Delta^\alpha \delta_1 \\ & \leq \frac{1}{\nu_1 \varpi_1} \left[ (\iota_1 - \delta_0)(\mathfrak{I}_1 - \delta_0) \right]^{\frac{\nu_1-1}{\nu_1}} \left[ (\iota_2 - \delta_0)(\mathfrak{I}_2 - \delta_0) \right]^{\frac{\varpi_1-1}{\varpi_1}} \\ & \times \left( \int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\mathfrak{I}_1} (\sigma(\iota_1) - \sigma(\mathfrak{I}_1)) (\sigma(\mathfrak{I}_1) - \sigma(\delta_1)) |\vartheta_2^{\Delta_2^\alpha \Delta_1^\alpha}(\mathfrak{I}_1, \delta_1)|^{\nu_1} \Delta^\alpha \mathfrak{I}_1 \Delta^\alpha \delta_1 \right)^{\frac{1}{\nu_1}} \\ & \left( \int_{\delta_0}^{\iota_2} \int_{\delta_0}^{\mathfrak{I}_2} (\sigma(\iota_2) - \delta_0) (\sigma(\mathfrak{I}_2) - \delta_0) |\vartheta_2^{\Delta_2^\alpha \Delta_1^\alpha}(\mathfrak{I}_2, \delta_2)|^{\nu_2} \Delta^\alpha \mathfrak{I}_2 \Delta^\alpha \delta_2 \right)^{\frac{1}{\nu_2}}. \end{aligned} \quad (3.18)$$

**Remark 3.11.** In Corollary 3.4, if  $\mathbb{T} = \mathbb{R}$ ,  $\alpha = 1$  we obtain an equivalent formulation of the inequality that Pachpatte presented in [2, Theorem 4].

**Remark 3.12.** In Corollary 3.4, if  $\mathbb{T} = \mathbb{Z}$ ,  $\alpha = 1$  we obtain an equivalent formulation of the inequality that Pachpatte presented in [2, Theorem 3].

**Theorem 3.3.** Let  $\mathbb{T}$  be a time scale with  $\delta_0, \iota_{ij}, \tau_{ij}, \delta_{ij} \in \mathbb{T}$ ,  $(i, j = 1, \dots, n)$ . Let  $\nu_i, \varpi_i > 1$ , be constants and  $\frac{1}{\nu_i} + \frac{1}{\varpi_i} = 1$ . Let  $\vartheta_i(\tau_{1i}, \dots, \tau_{ni})$  be real valued nth  $\Delta^\alpha$ -differentiable functions also that defined on  $[\delta_0, \iota_{1i}]_{\mathbb{T}} \times \dots \times [\delta_0, \iota_{ni}]_{\mathbb{T}}$ , where  $\delta_0 \leq \iota_{ji} \leq \delta_{ji}$ ,  $\delta_{ji} \in (0, \infty)$  and  $i, j = 1, \dots, n$ . Suppose

$$\vartheta_i(\iota_{1i}, \dots, \iota_{ni}) = \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \frac{\partial^n}{\Delta^\alpha \tau_{1i} \dots \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i},$$

then

$$\begin{aligned} & \int_{\delta_0}^{\delta_{11}} \dots \int_{\delta_0}^{\delta_{n1}} \int_{\delta_0}^{\delta_{12}} \dots \int_{\delta_0}^{\delta_{n2}} \dots \int_{\delta_0}^{\delta_{1n}} \dots \int_{\delta_0}^{\delta_{nn}} \\ & \frac{\prod_{i=1}^n \left( \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \left| \frac{\partial^n}{\Delta^\alpha \tau_{1i} \dots \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right|^{\nu_i} \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i} \right)^{\frac{1}{\nu_i}}}{\left( \sum_{i=1}^n \frac{[(\iota_{1i} - \delta_0) \dots (\iota_{ni} - \delta_0)]}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \\ & \Delta^\alpha \iota_{11} \dots \Delta^\alpha \iota_{n1} \dots \Delta^\alpha \iota_{12} \dots \Delta^\alpha \iota_{n2} \dots \Delta^\alpha \iota_{1n} \dots \Delta^\alpha \iota_{nn} \\ & \leq N \prod_{i=1}^n \left( \int_{\delta_0}^{\delta_{1i}} \dots \int_{\delta_0}^{\delta_{ni}} \prod_{j=1}^n (\sigma(\delta_{ji}) - \iota_{ji}) \left| \frac{\partial^n}{\Delta^\alpha \iota_{1i} \dots \Delta^\alpha \iota_{ni}} \vartheta_i(\iota_{1i}, \dots, \iota_{ni}) \right|^{\nu_i} \Delta^\alpha \iota_{1i} \dots \Delta^\alpha \iota_{ni} \right)^{\frac{1}{\nu_i}}, \end{aligned} \quad (3.19)$$

where

$$N = N(\delta_{1i}, \dots, \delta_{ni}) \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n [(\delta_{1i} - \delta_0) \dots (\delta_{ni} - \delta_0)]^{\frac{1}{\varpi_i}}.$$

*Proof.* From hypothesis of Theorem 3.3, we have

$$|\vartheta_i(\iota_{1i}, \dots, \iota_{ni})| \leq \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \left| \frac{\partial^n}{\Delta^\alpha \tau_{1i} \dots \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right| \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i}. \quad (3.20)$$

On other hand, by using (3.3) and Hölder's dynamic inequality, we obtain

$$\begin{aligned} & \prod_{i=1}^n |\vartheta_i(\iota_{1i}, \dots, \iota_{ni})| \\ & \leq \prod_{i=1}^n \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \left| \frac{\partial^n}{\Delta^\alpha \tau_{1i} \dots \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right| \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i} \\ & \leq \prod_{i=1}^n [(\iota_{1i} - \delta_0) \dots (\iota_{ni} - \delta_0)]^{\frac{1}{\varpi_i}} \\ & \quad \times \left( \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \left| \frac{\partial^n}{\Delta^\alpha \tau_{1i}, \dots, \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right|^{\nu_i} \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i} \right)^{\frac{1}{\nu_i}} \\ & \leq \frac{\left( \sum_{i=1}^n \frac{[(\iota_{1i} - \delta_0) \dots (\iota_{ni} - \delta_0)]}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}}{\left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{n - \sum_{i=1}^n \frac{1}{\nu_i}}} \\ & \quad \times \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \left| \frac{\partial^n}{\Delta^\alpha \tau_{1i} \dots \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right|^{\nu_i} \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i} \right)^{\frac{1}{\nu_i}}. \end{aligned} \quad (3.21)$$

Divide (3.21) by  $\left( \sum_{i=1}^n \frac{[(\iota_{1i} - \delta_0) \dots (\iota_{ni} - \delta_0)]}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}$ , then integrating over  $\iota_{ji}$  from  $\delta_0$  to  $\delta_{ji}$  ( $i, j = 1, \dots, n$ ), respectively, using dynamic Hölder's inequality and using the information  $\sigma(n) \geq n$ , we obtain

$$\begin{aligned} & \int_{\delta_0}^{\delta_{11}} \dots \int_{\delta_0}^{\delta_{n1}} \int_{\delta_0}^{\delta_{12}} \dots \int_{\delta_0}^{\delta_{n2}} \dots \int_{\delta_0}^{\delta_{1n}} \dots \int_{\delta_0}^{\delta_{nn}} \\ & \frac{\prod_{i=1}^n \left( \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \left| \frac{\partial^n}{\Delta^\alpha \tau_{1i} \dots \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right|^{\nu_i} \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i} \right)^{\frac{1}{\nu_i}}}{\left( \sum_{i=1}^n \frac{[(\iota_{1i} - \delta_0) \dots (\iota_{ni} - \delta_0)]}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \\ & \Delta^\alpha \iota_{11} \dots \Delta^\alpha \iota_{n1} \Delta^\alpha \iota_{12} \dots \Delta^\alpha \iota_{n2} \dots \Delta^\alpha \iota_{1n} \dots \Delta^\alpha \iota_{nn} \\ & \leq \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \\ & \quad \times \prod_{i=1}^n \int_{\delta_0}^{\delta_{1i}} \dots \int_{\delta_0}^{\delta_{ni}} \left( \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \left| \frac{\partial^n}{\Delta^\alpha \tau_{1i}, \dots, \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right|^{\nu_i} \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i} \right)^{\frac{1}{\nu_i}} \Delta^\alpha \iota_{ni} \dots \Delta^\alpha \iota_{1i} \\ & \leq \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n [(\delta_{1i} - \delta_0) \dots (\delta_{ni} - \delta_0)]^{\frac{1}{\varpi_i}} \\ & \quad \left( \int_{\delta_0}^{\delta_{1i}} \dots \int_{\delta_0}^{\delta_{ni}} \left( \int_{\delta_0}^{\iota_{1i}} \dots \int_{\delta_0}^{\iota_{ni}} \left| \frac{\partial^n}{\Delta^\alpha \tau_{1i} \dots \Delta^\alpha \tau_{ni}} \vartheta_i(\tau_{1i}, \dots, \tau_{ni}) \right|^{\nu_i} \Delta^\alpha \tau_{ni} \dots \Delta^\alpha \tau_{1i} \right)^{\frac{1}{\nu_i}} \Delta^\alpha \iota_{ni} \dots \Delta^\alpha \iota_{1i} \right)^{\frac{1}{\nu_i}} \end{aligned}$$

$$\begin{aligned}
&= N \prod_{i=1}^n \left( \int_{\delta_0}^{\delta_{1i}} \cdots \int_{\delta_0}^{\delta_{ni}} \prod_{j=1}^n (\delta_{ji} - \iota_{ji}) \left| \frac{\partial^n}{\Delta^\alpha \iota_{1i} \cdots \Delta^\alpha \iota_{ni}} \vartheta_i(\iota_{1i}, \dots, \iota_{ni}) \right|^{\nu_i} \Delta^\alpha \iota_{ni} \cdots \Delta^\alpha \iota_{1i} \right)^{\frac{1}{\nu_i}} \\
&\leq N \prod_{i=1}^n \left( \int_{\delta_0}^{\delta_{1i}} \cdots \int_{\delta_0}^{\delta_{ni}} \prod_{j=1}^n (\sigma(\delta_{ji}) - \iota_{ji}) \left| \frac{\partial^n}{\Delta^\alpha \iota_{1i} \cdots \Delta^\alpha \iota_{ni}} \vartheta_i(\iota_{1i}, \dots, \iota_{ni}) \right|^{\nu_i} \Delta^\alpha \iota_{ni} \cdots \Delta^\alpha \iota_{1i} \right)^{\frac{1}{\nu_i}}.
\end{aligned}$$

This concludes the evidence.  $\square$

**Remark 3.13.** In Theorem 3.3, supposing  $\mathbb{Z} = \mathbb{T}$ , and with  $\alpha = 1$ , obtains [4, Theorem 3.1].

**Remark 3.14.** In Theorem 3.3, supposing  $\mathbb{R} = \mathbb{T}$ , and with  $\alpha = 1$ , obtains [4, Theorem 3.2].

**Corollary 3.5.** Let  $\vartheta_i(\iota_{1i}, \dots, \iota_{ni})$  change to  $\vartheta_i(\mathfrak{I}_i)$  in Theorem 3.3 and in view of  $\vartheta_i(\delta_0) = 0$ , ( $i = 1, \dots, n$ ), then

$$\begin{aligned}
&\int_{\delta_0}^{\iota_1} \int_{\delta_0}^{\iota_2} \cdots \int_{\delta_0}^{\iota_n} \frac{\prod_{i=1}^n |\vartheta_i(\mathfrak{I}_i)|}{\left( \sum_{i=1}^n \frac{(\mathfrak{I}_i - \delta_0)}{\varpi_i} \right)^{\sum_{i=1}^n \frac{1}{\varpi_i}}} \Delta^\alpha \mathfrak{I}_n \Delta^\alpha \mathfrak{I}_{n-1} \cdots \Delta^\alpha \mathfrak{I}_1 \\
&\leq R \prod_{i=1}^n \left( \int_{\delta_0}^{\iota_i} (\sigma(\iota_i) - \sigma(\mathfrak{I}_i)) \left| \vartheta_i^{\Delta^\alpha}(\mathfrak{I}_i) \right|^{\nu_i} \Delta^\alpha \tau_i \Delta^\alpha \mathfrak{I}_i \right)^{\frac{1}{\nu_i}},
\end{aligned} \tag{3.22}$$

where

$$R = \left( n - \sum_{i=1}^n \frac{1}{\nu_i} \right)^{\sum_{i=1}^n \frac{1}{\nu_i} - n} \prod_{i=1}^n (\iota_i - \delta_0)^{\frac{1}{\varpi_i}}.$$

**Remark 3.15.** Taking  $n = 2$ , in Corollary 3.5, if  $\nu_1, \nu_2 > 1$  are such that  $\frac{1}{\nu_1} + \frac{1}{\nu_2} \geq 1$  and  $0 < \lambda = 2 - \frac{1}{\nu_1} - \frac{1}{\nu_2} = \frac{1}{\varpi_1} + \frac{1}{\varpi_2} \leq 1$ , inequality (3.22) reduces to inequality (3.6).

#### 4. Conclusions

In this work, we used Holder's inequality to prove a number of Hilbert's inequalities on the time scale. Some integer and discrete inequalities were obtained as special cases of the results. This work builds on the multiple inequalities reported by Pachpatte in 1998 and his 2000 and by Handley et al. and by Zhao et al. in 2013. In the future, one can generalize the results proved here by using diamond alpha calculus and also by trying to get the inverse inequalities.

#### Use of AI tools declaration

The authors declare they have not used Artificial Intelligence (AI) tools in the creation of this article.

#### Acknowledgments

This research work was funded by Institutional Fund Project under grant no. (IFPIP: 827-144-1443). The authors gratefully acknowledge technical and financial support provided by the Ministry of Education and King Abdulaziz University, DSR, Jeddah, Saudi Arabia.

---

## Conflict of interest

The authors declare that there is no competing interest.

## References

1. G. H. Hardy, J. E. Littlewood, G. Polya, *Inequalities*, Cambridge Univ. Press, Cambridge, 1952.
2. B. G. Pachpatte. Inequalities similar to certain extensions of Hilbert's inequality, *J. Math. Anal. Appl.*, **243** (2000), 217–227. <https://doi.org/10.1006/jmaa.1999.6646>
3. C. J. Zhao, L. Y. Chen, W. S. Cheung, On some new Hilbert-type inequalities, *Math. Slovaca*, **61** (2011), 15–28. <https://doi.org/10.2478/s12175-010-0056-0>
4. C. J. Zhao, W. S. Cheung, On Hilbert type inequalities, *J. Inequal. Appl.*, **2012** (2012), 145. <https://doi.org/10.1186/1029-242X-2012-145>
5. B. C. Yang, S. H. Wu, Q. Chen, A new extension of Hardy-Hilbert's inequality containing kernel of double power functions, *Mathematics*, **8** (2020), 894. <https://doi.org/10.3390/math8060894>
6. B. C. Yang, S. H. Wu, A. Z. Wang, A new Hilbert-type inequality with positive homogeneous kernel and its equivalent forms, *Symmetry*, **12** (2020), 342. <https://doi.org/10.3390/sym12030342>
7. Y. G. Sun, T. Hassan, Some nonlinear dynamic integral inequalities on time scales, *Appl. Math. Comput.*, **220** (2013), 221–225. <https://doi.org/10.1016/j.amc.2013.06.036>
8. J. Q. Liao, S. H. Wu, B. C. Yang, On a new half-discrete Hilbert-type inequality involving the variable upper limit integral and partial sums, *Mathematics*, **8** (2020), 229. <https://doi.org/10.3390/math8020229>
9. M. Bohner, A. Peterson, *Dynamic equations on time scales: An introduction with applications*, Birkhäuser Boston, Inc., Boston, MA, 2001.
10. S. Hilger, Analysis on measure chains—a unified approach to continuous and discrete calculus, *Results Math.*, **18** (1990), 18–56. <https://doi.org/10.1007/BF03323153>
11. R. Agarwal, D. O'Regan, S. Saker, *Dynamic inequalities on time scales*, Springer, Cham, 2014.
12. E. Akin-Bohner, M. Bohner, F. Akin, Pachpatte inequalities on time scales, *J. Inequal. Pure Appl. Math.*, **6** (2005).
13. M. Bohner, T. Matthews, The Grüss inequality on time scales, *Commun. Math. Anal.*, **3** (2007), 1–8.
14. M. Bohner, T. Matthews, Ostrowski inequalities on time scales, *J. Inequal. Pure Appl. Math.*, **9** (2008).
15. C. Dinu, Hermite-Hadamard inequality on time scales, *J. Inequal. Appl.*, **2008** (2008). <https://doi.org/10.1155/2008/287947>
16. R. Hilscher, A time scales version of a Wirtinger-type inequality and applications, *J. Comput. Appl. Math.*, **141** (2002), 219–226. [https://doi.org/10.1016/S0377-0427\(01\)00447-2](https://doi.org/10.1016/S0377-0427(01)00447-2)
17. W. N. Li, Some delay integral inequalities on time scales, *Comput. Math. Appl.*, **59** (2010), 1929–1936. <https://doi.org/10.1016/j.camwa.2009.11.006>

---

18. P. Rehak, Hardy inequality on time scales and its application to half-linear dynamic equations, *J. Inequal. Appl.*, **2005** (2005), 495–507. <https://doi.org/10.1155/JIA.2005.495>

19. R. P. Agarwal, V. Lakshmikantham, *Uniqueness and nonuniqueness criteria for ordinary differential equations*, World Scientific Publishing, Singapore, vol. 6 of Series in Real Analysis, 1993.

20. J. D. Li, Opial-type integral inequalities involving several higher order derivatives, *J. Math. Anal. Appl.*, **167** (1992), 98–110.

21. M. Bohner, A. C. Peterson, *Advances in dynamic equations on time scales*, Springer Science and Business Media, 2002.

22. M. Bohner, A. Peterson, *Dynamic equations on time scales: An introduction with applications*, Birkhauser, Boston, Mass, USA, 2001.

23. N. Benkhettou, S. Hassani, D. F. M. Torres, A conformable fractional calculus on arbitrary time scales, *J. King Saud Univ.-Sci.*, **28** (2016), 93–98. <https://doi.org/10.1016/j.jksus.2015.05.003>

24. E. R. Nwaeze, D. F. M. Torres, Chain rules and inequalities for the BHT fractional calculus on arbitrary timescales, *Arab. J. Math.*, **6** (2017), 13–20. <https://doi.org/10.1007/s40065-016-0160-2>

25. M. J. Cloud, B. C. Drachman, L. Lebedev, *Inequalities*, Springer, 1998.



AIMS Press

© 2023 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)