



---

*Research article*

## Solvability and controllability of second-order non-autonomous impulsive neutral evolution hemivariational inequalities

Yong-Ki Ma<sup>1</sup>, N. Valliammal<sup>2</sup>, K. Jothimani<sup>3</sup> and V. Vijayakumar<sup>4,\*</sup>

<sup>1</sup> Department of Applied Mathematics, Kongju National University, Chungcheongnam-do 32588, Korea

<sup>2</sup> Department of Mathematics, Sri Eshwar College of Engineering, Coimbatore 641202, Tamil Nadu, India

<sup>3</sup> Department of Mathematics, Sri G.V.G. Visalakshi College for Women, Udumalpet 642128, Tamil Nadu, India

<sup>4</sup> Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology, Vellore 632014, Tamil Nadu, India

\* **Correspondence:** Email: vijaysarovel@gmail.com.

**Abstract:** The primary aim of this article was to explore the approximate controllability of second-order impulsive hemivariational inequalities with initial conditions in Hilbert space. The mild solution was initially derived using the properties of the cosine and sine family of operators, Clarke's subdifferential, and the fact that the related linear equation has an evolution operator. The results of the approximate controllability of the considered systems were then taken into account using the fixed-point theorem method. An application was provided to support our theoretical findings.

**Keywords:** existence; approximate controllability; generalized Clarke's subdifferential; hemivariational inequalities

**Mathematics Subject Classification:** 34K30, 35R10, 35R12, 35R70, 93B05

---

### 1. Introduction

Clarke's subdifferential operator is associated with a type of nonlinear inclusion known as hemivariational inequalities. These inequalities have applications in structural analysis and non-convex optimization. The two main types of inequality problems are variational inequalities and hemivariational inequalities. Hemivariational inequalities handle nonsmooth, nonconvex energy functions, whereas variational inequalities primarily deal with convex energy functions. In 1981, Panagiotopoulos introduced the concept of hemivariational inequality as a method to represent

mechanical obstacles. Hemivariational inequalities or subdifferential inclusions can be used to model various nonsmooth contact mechanics issues involving multivalued and nonmonotone constitutive laws with boundary conditions (see [1–3]). Since then, numerous researchers have made significant contributions to the field of hemivariational inequalities, as seen in [4–9]. Moreover, neutral differential systems with impulsive effects have become a prominent area of research, modeling real-world processes that undergo sudden changes at specific points. This field has far-reaching applications in areas like finance, economics, mechanics, neural networks, electronics, and telecommunications. Notably, the authors of [10] investigated the controllability of nonlocal neutral differential inclusions with impulse effects. Researchers looked at the existence of impulsive multivalued neutral functional differential inclusions [11, 12].

Research on controllability in hemivariational-type systems has gained significant attention from the scientific community in recent times. Control problems have significant implications for various fields, including engineering, physics, and finance [13]. Despite the progress made, many intriguing questions and concepts remain unexplored. Notably, the authors of [14, 15] established the existence of optimal control in hemivariational inequalities, while [16, 17] investigated optimal control in parabolic hemivariational inequalities. Furthermore, [18, 19] demonstrated the existence of optimal control in hyperbolic systems, contributing to the advancement of the field. Due to their wide application to numerous pragmatist mathematics fields, neutral systems have attracted attention recently. Neutral systems have numerous applications in various fields, including thermal expansion of materials, biological advancements, surface waves, and stretchability, which benefit from neutral systems either directly or indirectly. Additionally, researchers have extensively studied hemivariational inequalities with a neutral type in [20, 21]. For further information regarding the system with hemivariational inequalities, refer to [22–26]. Recently, there has been a surge of interest in Clarke’s subdifferential evolution inclusion problems, particularly in the context of nonsmooth analysis and optimization. Frictional contact analysis can effectively characterize the interaction between a piezoelectric body and an electrically conducting foundation, and the frictional contact between a piezoelectric cylinder and a foundation exhibits anti-plane shear deformations. Moreover, the authors of [27] have established approximate controllability results for Sobolev-type Hilfer fractional neutral evolution problems with Clarke’s subdifferential-type problems. Our proposed problem presents a model that combines the key elements of hemivariational inequality and fractional impulsive differential equations into a unified framework. This study establishes approximate controllability results for a neutral differential system with impulsive effects, formulated as a hemivariational inequality. Next, we will define the specific system under consideration, which will be the focus of our investigation:

$$\left\{ \begin{array}{l} \langle -\frac{d}{d\varphi}[u'(\varphi) - g_1(\varphi, u(\varphi))] + \mathcal{A}(\varphi)u(\varphi) + \mathcal{B}v(\varphi), \omega \rangle_{\mathcal{M}} \\ \quad + H^\circ(\varphi, u(\varphi); \omega) \geq 0, \varphi_i \neq \varphi \in \mathfrak{L} = [0, a], \forall \omega \in \mathcal{M}, \\ u(0) = u_0, u'(0) = u_1, \\ \Delta u(\varphi_i) = \mathcal{L}_i(u(\varphi_i)), \\ \Delta u'(\varphi_i) = \mathcal{K}_i(u(\varphi_i)), i = 1, 2, 3, \dots, m, 0 < \varphi_1 < \varphi_2 < \dots < \varphi_m < a. \end{array} \right. \quad (1.1)$$

Here  $\langle \cdot, \cdot \rangle_{\mathcal{M}}$  stands for the inner product of the separable Hilbert space  $\mathcal{M}$  and  $\mathcal{A} : D(\mathcal{A}) \subseteq \mathcal{M} \rightarrow \mathcal{M}$  is a closed, linear, and densely defined operator on  $\mathcal{M}$ .  $H^\circ(\varphi, \cdot; \cdot)$  denotes the generalized Clarke’s directional derivative [6] of a locally Lipschitz function  $H(\varphi, \cdot) : \mathcal{M} \rightarrow \mathbb{R}$ . The function  $g_1 : \mathfrak{L} \times \mathcal{M} \rightarrow$

$\mathcal{M}$ . The control function  $v \in V$  takes values from  $L^2(\mathcal{Q}, V)$  and  $V$  is the control set which is also a Hilbert space. Let  $\mathcal{B} : V \rightarrow \mathcal{M}$  be the bounded linear operator. Let  $0 < \varphi_1 < \varphi_2 < \dots < \varphi_m < \varphi_{m+1} = a$  be pre-fixed points and the symbol  $\Delta u(\varphi_i)$  represents the jump in the state  $u$  at time  $t$  which is defined by  $\Delta u(\varphi_i) = u(\varphi_i^+) - u(\varphi_i^-)$ , and  $u(\varphi_i^-)$  and  $u(\varphi_i^+)$  denote the left and right limit of  $u$  at  $\varphi_i$ .  $\mathcal{L}_i : \mathcal{M} \rightarrow \mathcal{M}$  and  $\mathcal{K}_i : \mathcal{M} \rightarrow \mathcal{M}, i = 1, 2, 3, \dots, m$ , are impulsive functions.

The structure of the article is presented in the following way:

- We begin by reviewing key definitions, fundamental theorems, and initial findings from a previous section.
- The core focus then shifts to exploring the existence of mild solutions for the system represented by Eq (1.1).
- Next, we investigate the approximate controllability of the system (1.1).
- Finally, we solidify our theoretical results by presenting a concrete example.

## 2. Foundations and background

Let  $\mathcal{Z}$  and  $\mathcal{Z}^*$  be a Banach space and its dual, respectively with  $\|\cdot\|_{\mathcal{Z}}$  and  $\langle \cdot, \cdot \rangle$  is the duality pairing of  $\mathcal{Z}$  and  $\mathcal{Z}^*$ . The Banach space  $\mathcal{PC}(\mathcal{Q}, \mathcal{Z})$  is the set of all piecewise continuous functions from  $\mathcal{Q} = [0, a]$  into  $\mathcal{Z}$  together with  $\|u\|_{\mathcal{PC}} = \sup_{\varphi \in \mathcal{Q}} \|u(\varphi)\|_{\mathcal{Z}}$ . Also,

$$\mathcal{M}_{(w)k(e)}(\mathcal{Z}) := \{\bar{U} \subseteq \mathcal{Z} : \bar{U} \neq \emptyset \text{ is (weakly) compact (convex)}\};$$

$$\mathcal{M}_f(e)(\mathcal{Z}) := \{\bar{U} \subseteq \mathcal{Z} : \bar{U} \neq \emptyset \text{ is closed (convex)}\}.$$

Consider the non-autonomous second-order initial value problem:

$$z''(\varphi) = \mathcal{A}(\varphi)z(\varphi) + f(\varphi), \quad 0 \leq z, \varphi \leq a, \quad (2.1)$$

$$z(z) = z_0, \quad z'(z) = z_1, \quad (2.2)$$

where  $\mathcal{A}(\varphi) : D(\mathcal{A}(\varphi)) \subseteq \mathcal{M} \rightarrow \mathcal{M}, \varphi \in [0, a]$  is a closed densely defined operator and  $f : [0, a] \rightarrow \mathcal{M}$  is an appropriate function. One can refer to [15, 28–30] and the references therein. A significant number of articles relate the existence of solutions to the (2.1)-(2.2) problem to the existence of the evolution operator  $Q(\varphi, z)$  for the homogeneous equation

$$z''(\varphi) = \mathcal{A}(\varphi)z(\varphi), \quad 0 \leq \varphi \leq a. \quad (2.3)$$

Assume that the domain of  $\mathcal{A}(\varphi)$  is a subspace  $D$  that is dense in  $\mathcal{M}$  and independent of  $\varphi$ , and that the function  $\varphi \mapsto \mathcal{A}(\varphi)z$  is continuous for each  $z \in D(\mathcal{A}(\varphi))$ .

In view of Kozak's work [31], we shall apply the following evolution operator notion in this study.

**Definition 2.1.** A family  $Q$  of bounded linear operators  $Q(\varphi, z) : [0, a] \times [0, a] \rightarrow \mathcal{L}(\mathcal{M})$  is called an evolution operator for (2.3) if the following conditions are satisfied:

(Z1) For each  $z \in \mathcal{M}$ , the mapping  $[0, a] \times [0, a] \ni (\varphi, z) \rightarrow Q(\varphi, z)z \in \mathcal{M}$  is of class  $C^1$  and

(i) for each  $\varphi \in [0, a], Q(\varphi, \varphi) = 0$ ,

(ii) for all  $\varphi, z \in [0, a]$ , and for each  $\mathfrak{z} \in \mathcal{M}$ ,

$$\frac{\partial}{\partial \varphi} Q(\varphi, z)\mathfrak{z}\Big|_{\varphi=z} = \mathfrak{z}, \quad \frac{\partial}{\partial z} Q(\varphi, z)\mathfrak{z}\Big|_{\varphi=z} = -\mathfrak{z}.$$

(Z2) For all  $\varphi, z \in [0, a]$ , if  $\mathfrak{z} \in D(\mathcal{A})$ , then  $Q(\varphi, z)\mathfrak{z} \in D(\mathcal{A})$ , the mapping  $[0, a] \times [0, a] \ni (\varphi, z) \rightarrow Q(\varphi, z)\mathfrak{z} \in \mathcal{M}$  is of class  $C^2$  and

- (i)  $\frac{\partial^2}{\partial \varphi^2} Q(\varphi, z)\mathfrak{z} = \mathcal{A}(\varphi)Q(\varphi, z)\mathfrak{z}$ ,
- (ii)  $\frac{\partial^2}{\partial z^2} Q(\varphi, z)\mathfrak{z} = Q(\varphi, z)B(z)\mathfrak{z}$ ,
- (iii)  $\frac{\partial}{\partial z} \frac{\partial}{\partial \varphi} Q(\varphi, z)\mathfrak{z}\Big|_{\varphi=z} = 0$ .

(Z3) For all  $\varphi, z \in [0, a]$ , if  $\mathfrak{z} \in D(\mathcal{A})$ , and then  $\frac{\partial}{\partial z} Q(\varphi, z)\mathfrak{z} \in D(\mathcal{A})$ , then  $\frac{\partial^2}{\partial \varphi^2} \frac{\partial}{\partial z} Q(\varphi, z)\mathfrak{z}$ ,  $\frac{\partial^2}{\partial z^2} \frac{\partial}{\partial \varphi} Q(\varphi, z)\mathfrak{z}$ , and

- (i)  $\frac{\partial^2}{\partial \varphi^2} \frac{\partial}{\partial z} Q(\varphi, z)\mathfrak{z} = \mathcal{A}(\varphi) \frac{\partial}{\partial z} Q(\varphi, z)\mathfrak{z}$ ,
- (ii)  $\frac{\partial^2}{\partial z^2} \frac{\partial}{\partial \varphi} Q(\varphi, z)\mathfrak{z} = \frac{\partial}{\partial \varphi} Q(\varphi, z)\mathcal{A}(z)\mathfrak{z}$ ,

and the mapping  $[0, a] \times [0, a] \ni (\varphi, z) \rightarrow \mathcal{A}(\varphi) \frac{\partial}{\partial z} Q(\varphi, z)\mathfrak{z}$  is continuous.

Let us take

$$C(\varphi, z) = -\frac{\partial Q(\varphi, z)}{\partial z}.$$

Also, for some positive constants  $\mathcal{N}_1$  and  $\mathcal{N}_2$ , we set  $\sup_{0 \leq \varphi, z \leq a} \|Q(\varphi, z)\| \leq \mathcal{N}_2$  and  $\sup_{0 \leq \varphi, z \leq a} \|C(\varphi, z)\| \leq \mathcal{N}_1$  and

$$\|Q(\varphi + g, z) - Q(\varphi, z)\| \leq \mathcal{N}_1 |g|, \quad (2.4)$$

for all  $z, \varphi, \varphi + g \in [0, a]$ . The mild solution  $\mathfrak{z} : [0, a] \rightarrow \mathcal{M}$  of (2.1)-(2.2) is:

$$\mathfrak{z}(\varphi) = C(\varphi, z)\mathfrak{z}_0 + Q(\varphi, z)\mathfrak{z}_1 + \int_0^\varphi Q(\varphi, \eta)f(\eta)d\eta.$$

Let us start by discussing the necessary definitions and findings from the multivalued analysis. The following sources [7, 32, 33] are recommended for readers in addition to multivalued maps.

**Definition 2.2.** Given a Banach space  $\mathcal{Z}$  and a multivalued map  $H : \mathcal{Z} \rightarrow 2^{\mathcal{Z}} \setminus \{\emptyset\} = \mathcal{N}(\mathcal{Z})$ , we say

- (i)  $H$  is convex and closed valued, only when  $H(u)$  is convex and closed valued for all  $u \in \mathcal{Z}$ .
- (ii)  $H$  is said to be upper semicontinuous on  $\mathcal{Z}$ , if for all  $y \in \mathcal{Z}$ ,  $H(y) \neq \emptyset$  is closed in  $\mathcal{Z}$  and if for each open set  $\mathbb{J}_1$  of  $\mathcal{Z}$  which contains  $H(y)$ , then there is an open neighborhood  $\mathbb{E}$  of  $y$  such that  $H(\mathbb{E}) \subseteq \mathbb{J}_1$ .
- (iii)  $H$  is bounded on bounded sets if  $H(B) = \cup_{u \in B} H(u)$  is bounded in  $\mathcal{Z}$  for all  $B \in \mathcal{M}_b(\mathcal{Z})$  (i.e.,  $\sup_{u \in B} \{\sup\{\|k\| : k \in H(u)\}\} < \infty$ ).
- (iv)  $H$  is supposed to be completely continuous provided that  $H(\mathbb{J}_1)$  is relatively compact, for all bounded subset  $\mathbb{J}_1 \subseteq \mathcal{Z}$ .
- (v)  $H$  has a fixed point if there is a  $u \in \mathcal{Z}$  such that  $u \in H(u)$ .

For a locally Lipschitzian functional  $H : \mathcal{Z} \rightarrow \mathbb{R}$ , we denote  $H^0(q; p)$ , the Clarke's generalized directional derivative of  $H$  at point  $q$  in the direction of  $p$ , that is,

$$H^0(q; p) := \limsup_{\gamma \rightarrow 0^+} \sup_{\vartheta \rightarrow q} \frac{H(\vartheta + \gamma p) - H(\vartheta)}{\gamma}.$$

Also,  $\partial H(q) := \{q^* \in \mathcal{Z}^* : H^0(q; p) \geq \langle q^*, p \rangle, \text{ for every } p \in \mathcal{Z}\}$  denotes the generalized Clarke's subdifferential.

The subsequent features and outcomes will facilitate our goal achievement:

**Lemma 2.3.** [9] *If the function  $H : \mathcal{U} \rightarrow \mathbb{R}$  is a locally Lipschitz on an open set  $\mathcal{U}$  of  $\mathcal{Z}$ , then*

- (i) *For all  $p \in \mathcal{Z}$ , it has  $H^0(q; p) = \max\{\langle q^*, p \rangle : \text{for all } q^* \in \partial H(q)\}$ ;*
- (ii) *For all  $q \in \mathcal{U}$ , the gradient  $\partial H(q)$  is a nonempty, convex, weak\*-compact subset of  $\mathcal{Z}^*$  and  $\|q^*\|_{\mathcal{Z}^*} \leq M$ , for all  $q^* \in \partial H(q)$ ;*
- (iii) *The graph of  $\partial H$  is closed in  $\mathcal{U} \times \mathcal{Z}^*$ . That is, if  $\{q_n\} \subset \mathcal{U}$ ,  $\{q_n^*\} \subset \mathcal{Z}^*$  are sequences as  $q_n^* \in \partial H(q_n)$  and  $q_n \rightarrow q \in \mathcal{Z}$ ,  $q_n^* \rightarrow q^*$  weakly\* in  $\mathcal{Z}^*$ , then  $q^* \in \partial H(q)$  (where the Banach space  $\mathcal{Z}^*$  furnished with the  $\ell^*$ -topology is denoted by  $\mathcal{Z}^*$ );*
- (iv) *The multifunction  $\mathcal{U} \ni q \rightarrow \partial H(q) \subseteq \mathcal{Z}^*$  is upper semicontinuous which maps  $\mathcal{U}$  into  $\mathcal{Z}^*$ .*

**Lemma 2.4.** [9] *Let  $\mathcal{Z}$  be the separable reflexive Banach space,  $0 < a < \infty$  and  $H : (0, a) \times \mathcal{Z} \rightarrow \mathbb{R}$ , such that  $H(\cdot, w)$  is measurable for each  $w \in \mathcal{Z}$  and  $H(\varphi, \cdot)$  is locally Lipschitz on  $\mathcal{Z}$  for all  $\varphi \in (0, a)$ . Then the multifunction  $(0, a) \times \mathcal{Z} \ni (\varphi, w) \mapsto \partial H(\varphi, w) \subset \mathcal{Z}^*$  is measurable.*

As discussed in [9], we can investigate the existence of mild solutions and approximate controllability for the following semilinear inclusions:

$$\begin{cases} \left[ \frac{d}{d\varphi} \left[ z'(\varphi) - g_1(\varphi, u(\varphi)) \right] \right] \in \mathcal{A}(\varphi)u(\varphi) + \mathcal{B}v(\varphi) + \partial H(\varphi, z(\varphi)), \varphi \in \mathcal{Q} = [0, a], \\ u(0) = u_0, u'(0) = u_1, \\ \Delta u(\varphi_i) = \mathcal{L}_i(u(\varphi_i)), \Delta u'(\varphi_i) = \mathcal{K}_i(u(\varphi_i)), i = 1, 2, 3, \dots, m. \end{cases} \quad (2.5)$$

Now, we can explore the implication that every solution to Eq (2.5) is also a solution to Eq (1.1). Hence, if  $u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M})$  is a solution of (2.5), there exists  $h(\varphi) \in \partial H(\varphi, u(\varphi))$  provided  $h \in L^2(\mathcal{Q}, \mathcal{M})$  and

$$\begin{cases} \left[ \frac{d}{d\varphi} \left[ u'(\varphi) - g_1(\varphi, u(\varphi)) \right] \right] = \mathcal{A}(\varphi)u(\varphi) + \mathcal{B}v(\varphi) + h(\varphi), \varphi \in \mathcal{Q} = [0, a], \\ u(0) = u_0, u'(0) = u_1, \\ \Delta z(\varphi_i) = \mathcal{L}_i(u(\varphi_i)), \Delta u'(\varphi_i) = \mathcal{K}_i(u(\varphi_i)), i = 1, 2, 3, \dots, m, \end{cases}$$

which implies

$$\begin{cases} \left\langle -\frac{d}{d\varphi} \left[ u'(\varphi) - g_1(\varphi, u(\varphi)) \right] + \mathcal{A}(\varphi)u(\varphi) + \mathcal{B}v(\varphi), \omega \right\rangle_{\mathcal{M}} \\ \quad + \langle h(\varphi), \omega \rangle_{\mathcal{M}} = 0, \varphi \in \mathcal{Q} = [0, a], \text{ for all } \omega \in \mathcal{M}, \\ u(0) = u_0, u'(0) = u_1, \\ \Delta z(\varphi_i) = \mathcal{L}_i(u(\varphi_i)), \Delta u'(\varphi_i) = \mathcal{K}_i(u(\varphi_i)), i = 1, 2, 3, \dots, m. \end{cases}$$

Since  $\langle h(\varphi), \omega \rangle_{\mathcal{M}} \leq H^0(\varphi, u(\varphi); \omega)$  and  $h(\varphi) \in \partial H(\varphi, u(\varphi))$ ,

$$\begin{cases} \left\langle -\frac{d}{d\varphi} \left[ u'(\varphi) - g_1(\varphi, u(\varphi)) \right] + \mathcal{A}(\varphi)u(\varphi) + \mathcal{B}v(\varphi), \omega \right\rangle_{\mathcal{M}} \\ \quad + H^0(\varphi, u(\varphi); \omega), \quad \varphi \in \mathcal{Q} = [0, a], \text{ for all } \omega \in \mathcal{M}, \\ u(0) = u_0, \quad u'(0) = u_1, \\ \Delta u(\varphi_i) = \mathcal{L}_i(u(\varphi_i)), \quad \Delta u'(\varphi_i) = \mathcal{K}_i(u(\varphi_i)), \quad i = 1, 2, 3, \dots, m. \end{cases}$$

Therefore, our initial focus will be on examining the semilinear inclusion (2.5), which proceeds our investigation of the hemivariational inequality (1.1). According to established literature [33, 34], the mild solution for problem (2.5) is defined as below.

**Definition 2.5.** For all  $v \in L^2(\mathcal{Q}, V)$ , a function  $u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M})$  is a mild solution for (2.5) if there exists  $h \in L^2(\mathcal{Q}, \mathcal{M})$  as  $h(\varphi) \in \partial H(\varphi, u(\varphi))$  almost everywhere on  $\varphi \in \mathcal{Q}$ ,

$$\begin{aligned} u(\varphi) = & C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] + \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz + \int_0^\varphi Q(\varphi, z)h(z)dz \\ & + \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz + \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)), \quad \varphi \in \mathcal{Q}. \end{aligned}$$

Let us take the following assumptions:

(A1) The function  $H : \mathcal{Q} \times \mathcal{M} \rightarrow \mathbb{R}$  satisfies:

- (i)  $\forall u \in \mathcal{M}, \varphi \mapsto H(\varphi, z)$  is measurable;
- (ii)  $u \mapsto H(\varphi, u)$  is locally Lipschitz continuous for a.e.  $\varphi \in \mathcal{Q}$ ;
- (iii) There is a function  $b(\varphi) \in L^2(\mathcal{Q}, \mathbb{R}^+)$  and  $e > 0$  such that

$$\begin{aligned} \|\partial H(\varphi, u)\| &= \sup\{\|h\| : h(\varphi) \in \partial H(\varphi, u)\} \leq b(\varphi) + e\|u\|, \\ &\text{for a.e. } \varphi \in \mathcal{Q} \text{ and for each } u \in \mathcal{M}. \end{aligned}$$

(A2) In the function  $g_1 : \mathcal{Q} \times \mathcal{M} \rightarrow \mathcal{M}$ , there exists some constants  $C_1, C_2 > 0$ , and for every  $u \in \mathcal{M}$ ,  $\varphi_1, \varphi_2 \in \mathcal{Q}$ , we have

$$\begin{aligned} \|g_1(\varphi_1, u(\varphi_1)) - g_1(\varphi_2, u(\varphi_2))\| &\leq C_1\|u(\varphi_1) - u(\varphi_2)\|, \\ \|g_1(\varphi_1, u(\varphi_1))\| &\leq C_2(1 + \|u(\varphi_1)\|). \end{aligned}$$

(A3) For some constants  $c_i, l_i > 0$ , the maps  $\mathcal{L}_i, \mathcal{K}_i : \mathcal{M} \rightarrow \mathcal{M}$  are continuous, and

$$\|\mathcal{L}_i(\varphi)\| \leq c_i, \quad \|\mathcal{K}_i(\varphi)\| \leq l_i, \quad i = 1, 2, 3, \dots, m, \text{ for all } u \in \mathcal{M}.$$

Let  $\mathbb{M} : L^2(\mathcal{Q}, \mathcal{M}) \rightarrow 2^{L^2(\mathcal{Q}, \mathcal{M})}$  be defined as below:

$$\mathbb{M}(u) = \{k \in L^2(\mathcal{Q}, \mathcal{M}) : k(\varphi) \in \partial H(\varphi, u(\varphi)) \text{ almost everywhere, } \varphi \in \mathcal{Q}\}, \text{ for all } u \in L^2(\mathcal{Q}, \mathcal{M}).$$

**Lemma 2.6.** [35] Let (A1) and  $\mathbb{M}$  be true. If  $u_n \rightarrow u$  in  $L^2(\mathcal{Q}, \mathcal{M})$ ,  $w_n \rightarrow w$  weakly in  $L^2(\mathcal{Q}, \mathcal{M})$  and  $u_n \in \mathbb{M}(u_n)$ , and then  $u \in \mathbb{M}(u)$ .

**Lemma 2.7.** [9] Let all hypotheses and **(A1)** hold. Then for every  $u \in L^2(\mathcal{Q}, \mathcal{M})$ , the set  $\mathbb{M}(u)$  has nonempty, weakly compact and convex values.

**Theorem 2.8.** [9] Consider the Banach space  $\mathcal{Z}$  which is locally convex and  $\Upsilon : \mathcal{Z} \rightarrow 2^{\mathcal{Z}}$  is a compact convex valued, upper semicontinuous multivalued map such that there exists a closed neighborhood  $\mathbb{J}$  of 0 for which  $\Upsilon(\mathbb{J})$  is relatively compact provided

$$\bar{U} = \{x \in \mathcal{Z} : \gamma x \in \Upsilon(u), \text{ for some } \gamma > 1\}$$

is bounded. Then  $\Upsilon$  has a fixed point.

### 3. Existence results

**Theorem 3.1.** For all  $v \in L^2(\mathcal{Q}, V)$ , provided that **(A1)–(A3)** are fulfilled, then (2.5) has a mild solution on  $\mathcal{Q}$  such that  $a(\mathcal{N}_1 C_2 + \mathcal{N}_2 e) < 1$ , where  $\mathcal{N}_1 := \sup_{\varphi, z \in [0, a]} \|C(\varphi, z)\|$ ,  $\mathcal{N}_2 := \sup_{\varphi, z \in [0, a]} \|Q(\varphi, z)\|$ .

*Proof.* Initially, choose any  $u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M}) \subset L^2(\mathcal{Q}, \mathcal{M})$ , by Lemma 2.7. Now, define  $\Upsilon : \mathcal{PC}(\mathcal{Q}, \mathcal{M}) \rightarrow 2^{\mathcal{PC}(\mathcal{Q}, \mathcal{M})}$  as:

$$\begin{aligned} \Upsilon(u) = & \left\{ f \in \mathcal{PC}(\mathcal{Q}, \mathcal{M}) : f(\varphi) = C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \right. \\ & + \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz + \int_0^\varphi Q(\varphi, z)h(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \\ & \left. + \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)), h \in \mathbb{M}(u) \right\}, u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M}). \end{aligned}$$

It is clear that we can determine a fixed point of the multivalued map  $\Upsilon$  that satisfies Theorem 2.8. First, note that the set-valued map  $\Upsilon(u)$  is convex due to the properties of  $\mathbb{M}(u)$ . Now, let us proceed with the proof of the theorem as follows:

**Step 1.**  $\Upsilon(\mathcal{B}_r) \subseteq (\mathcal{B}_r)$ ,  $r > 0$ , is bounded in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$ , where  $\mathcal{B}_r = \{u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M}) : \|u\|_{\mathcal{PC}} \leq r\}$ . Here, it suffices to demonstrate the existence of a positive constant  $\ell$  such that for each  $\sigma \in \Upsilon(u)$ ,  $u \in \mathcal{B}_r$ ,  $\|\sigma\|_{\mathcal{PC}} \leq \ell$ . If  $\sigma \in \Upsilon(u)$ , then there is a  $h \in \mathbb{M}(u)$  provided

$$\begin{aligned} \sigma(\varphi) = & C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \\ & + \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz + \int_0^\varphi Q(\varphi, z)h(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \\ & + \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)), \quad \varphi \in \mathcal{Q}. \end{aligned}$$

By Hölder's inequality,

$$\begin{aligned} \|\sigma(\varphi)\| \leq & \|C(\varphi, 0)u_0\| + \|Q(\varphi, 0)[u_1 - g_1(0, u(0))]\| + \left\| \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz \right\| \\ & + \left\| \int_0^\varphi Q(\varphi, z)h(z)dz \right\| + \left\| \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \right\| \end{aligned}$$

$$\begin{aligned}
& + \sum_{0 < \varphi_i < \varphi} \|C(\varphi, \varphi_i) \mathcal{L}_i(u(\varphi_i))\| + \sum_{0 < \varphi_i < \varphi} \|C(\varphi, \varphi_i) \mathcal{K}_i(u(\varphi_i))\| \\
& \leq \mathcal{N}_1 \|u_0\| + \mathcal{N}_2 \| [u_1 - g_1(0, u(0))] \| + \mathcal{N}_1 \int_0^\varphi C_2(1+r) dz \\
& \quad + \mathcal{N}_2 \int_0^\varphi [b(z) + er + \|\mathcal{B}\| \|v(z)\|] dz \\
& \quad + \mathcal{N}_1 \sum_{0 < \varphi_i < \varphi} \|\mathcal{L}_i(u(\varphi_i))\| + \mathcal{N}_2 \sum_{0 < \varphi_i < \varphi} \|\mathcal{K}_i(u(\varphi_i))\| \\
& \leq \mathcal{N}_1 \|u_0\| + \mathcal{N}_2 \varphi_2 \| [u_1 - g_1(0, u(0))] \| + \mathcal{N}_1 a C_2(1+r) + \mathcal{N}_2 [\sqrt{a} \|b\|_{L^2(\mathcal{Q}, \mathbb{R}^+)} + era \\
& \quad + \sqrt{a} \|\mathcal{B}\| \|v\|_{L^2(\mathcal{Q}, V)}] + \mathcal{N}_1 \sum_{i=1}^m c_i + \mathcal{N}_2 \sum_{i=1}^m l_i := \ell.
\end{aligned}$$

Thus,  $\Upsilon(\mathfrak{B}_r)$  is bounded.

**Step 2.**  $\{\Upsilon(u) : u \in \mathfrak{B}_r\}$  is completely continuous.

Let us note that for any  $u \in \mathfrak{B}_r$ ,  $\sigma \in \Upsilon(u)$ , there exists  $h \in \mathbb{M}(u)$  such that for all  $\varphi \in \mathcal{Q}$ ,

$$\begin{aligned}
\|\sigma(\varphi)\| & \leq \|C(\varphi, 0)u_0\| + \|Q(\varphi, 0)[u_1 - g_1(0, u(0))]\| + \left\| \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz \right\| \\
& \quad + \left\| \int_0^\varphi Q(\varphi, z)h(z)dz \right\| + \left\| \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \right\| \\
& \quad + \sum_{0 < \varphi_i < \varphi} \|C(\varphi, \varphi_i) \mathcal{L}_i(u(\varphi_i))\| + \sum_{0 < \varphi_i < \varphi} \|C(\varphi, \varphi_i) \mathcal{K}_i(u(\varphi_i))\|.
\end{aligned}$$

For  $0 < \xi_1 < \xi_2 \leq a$  and  $\mathfrak{t} > 0$  very small,

$$\begin{aligned}
\|\sigma(\xi_2) - \sigma(\xi_1)\|_{\mathcal{M}} & \leq \|C(\xi_2, 0) - C(\xi_1, 0)\| \|u_0\| + \|Q(\xi_2, 0) - Q(\xi_1, 0)\| \| [u_1 - g_1(0, u(0))] \| \\
& \quad + \int_0^{\xi_1 - \mathfrak{t}} \|C(\xi_2, z) - C(\xi_1, z)\| \|g_1(z, u(z))\| dz \\
& \quad + \int_{\xi_1 - \mathfrak{t}}^{\xi_1} \|C(\xi_2, z) - C(\xi_1, z)\| \|g_1(z, u(z))\| dz \\
& \quad + \int_{\xi_1}^{\xi_2} \|C(\xi_2, z)\| \|g_1(z, u(z))\| dz \\
& \quad + \int_0^{\xi_1 - \mathfrak{t}} \|Q(\xi_2, z) - Q(\xi_1, z)\| \|h(z) + \mathcal{B}v(z)\| dz \\
& \quad + \int_{\xi_1 - \mathfrak{t}}^{\xi_1} \|Q(\xi_2, z) - Q(\xi_1, z)\| \|h(z) + \mathcal{B}v(z)\| dz \\
& \quad + \int_{\xi_1}^{\xi_2} \|Q(\xi_2, z)\| \|h(z) + \mathcal{B}v(z)\| dz \\
& \quad + \sum_{0 < \varphi_i < a} \|C(\xi_2, \varphi_i) - C(\xi_1, \varphi_i)\| \|\mathcal{L}_i(u(\varphi_i))\| \\
& \quad + \sum_{0 < \varphi_i < a} \|Q(\xi_2, \varphi_i) - Q(\xi_1, \varphi_i)\| \|\mathcal{K}_i(u(\varphi_i))\|
\end{aligned}$$



$$\begin{aligned}
&\leq \|C(\xi_2, 0) - C(\xi_1, 0)\| \|u_0\| + \|Q(\xi_2, 0) - Q(\xi_1, 0)\| \|u_1 - g_1(0, u(0))\| \\
&\quad + \int_0^{\xi_1 - \mathfrak{k}} \|C(\xi_2, z) - C(\xi_1, z)\| C_2(1+r) dz \\
&\quad + \int_{\xi_1 - \mathfrak{k}}^{\xi_1} \|C(\xi_2, z) - C(\xi_1, z)\| C_2(1+r) dz + \mathcal{N}_1 C_2(1+r)(\xi_2 - \xi_1) \\
&\quad + \int_0^{\xi_1 - \mathfrak{k}} \|Q(\xi_2, \varphi_i) - Q(\xi_1, \varphi_i)\| [b(z) + er + \|\mathcal{B}\| \|v(z)\|] dz \\
&\quad + \int_{\xi_1 - \mathfrak{k}}^{\xi_1} \|Q(\xi_2, z) - Q(\xi_1, z)\| [b(z) + er + \|\mathcal{B}\| \|v(z)\|] dz \\
&\quad + \mathcal{N}_2 \int_{\xi_1}^{\xi_2} [b(z) + er + \|\mathcal{B}\| \|v(z)\|] dz + \sum_{0 < \varphi_i < a} \|C(\xi_2, \varphi_i) - C(\xi_1, \varphi_i)\| \|\mathcal{L}_i(u(\varphi_i))\| \\
&\quad + \sum_{0 < \varphi_i < a} \|Q(\xi_2, \varphi_i) - Q(\xi_1, \varphi_i)\| \|\mathcal{K}_i(u(\varphi_i))\|. \tag{3.1}
\end{aligned}$$

From the uniform operator topology [33, Lemma 6.2], it is easily understood that (3.1) tends to zero of  $u \in \mathfrak{B}_r$  as  $\xi_2 \rightarrow \xi_1$  and  $\mathfrak{k} \rightarrow 0$ .

Equivalently, for  $\xi_1 = 0$  and  $0 < \xi_2 \leq a$ , we can show that  $\|\sigma(\xi_2) - u_0\|_{\mathcal{M}}$  tends to zero independently of  $u \in \mathfrak{B}_r$  as  $\xi_2 \rightarrow 0$ . Hence, we can conclude that  $\{\Upsilon(u) : u \in \mathfrak{B}_r\}$  is equicontinuous of  $\mathcal{P}\mathcal{C}(\mathcal{Q}, \mathcal{M})$ .

Finally, from the assumptions (A1) and (A3) and by the definition of a relatively compact set, it is not difficult to check that  $\{\sigma(t) : \sigma \in \Upsilon(\mathfrak{B}_r)\}$  is relatively compact in  $\mathcal{M}$ . Thus, by the generalized Arzelà-Ascoli theorem, we get that  $\Upsilon$  is a multivalued compact map.

Therefore, based on the above arguments,  $\Upsilon$  is completely continuous.

**Step 3.** Assume  $u_n \rightarrow u_*$  in  $\mathcal{P}\mathcal{C}(\mathcal{Q}, \mathcal{M})$ ,  $\sigma_n \in \Upsilon(u_n)$  and  $\sigma_n \rightarrow \sigma_*$  in  $\mathcal{P}\mathcal{C}(\mathcal{Q}, \mathcal{M})$ . Let us check  $\sigma_* \in \Upsilon(u_*)$ . It is obvious that  $\sigma_n \in \Upsilon(u_n)$  exist only when  $h_n \in \mathbb{M}(u_n)$  such that

$$\begin{aligned}
\sigma_n(\varphi) &= C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \\
&\quad + \int_0^\varphi C(\varphi, z)g_1(z, u_n(z))dz + \int_0^\varphi Q(\varphi, z)h_n(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \\
&\quad + \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)). \tag{3.2}
\end{aligned}$$

Here  $\{h_n\}_{n \geq 1} \subseteq L^2(\mathcal{Q}, \mathcal{M})$  is bounded from the hypothesis (A2). Hence we may assume that

$$h_n \rightarrow h_* \text{ weakly in } L^2(\mathcal{Q}, \mathcal{M}). \tag{3.3}$$

From (3.2) and (3.3), we have

$$\begin{aligned}
\sigma_n(\varphi) &\rightarrow C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \\
&\quad + \int_0^\varphi C(\varphi, z)g_1(z, u_*(z))dz + \int_0^\varphi Q(\varphi, z)h_*(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \\
&\quad + \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)). \tag{3.4}
\end{aligned}$$

We can see that  $\sigma_n \rightarrow \sigma_*$  in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$  and  $h_n \in \mathbb{M}(u)$ . According to Lemma 2.6 and (3.4),  $h_* \in \mathbb{M}(u_*)$ . Then,  $\sigma_* \in \Upsilon(u_*)$ , and this shows that  $\Upsilon$  has a closed graph. Hence using Proposition 3.12 of [36] implies that it is upper semicontinuous.

**Step 4.** A priori estimate.

Based on the results from Steps 1–3, we have established that the multivalued map  $\Upsilon$  satisfies the following properties: upper semicontinuity, convex-valuedness, compactness, and relative compactness of  $\Upsilon(\mathcal{B}_r)$ . Therefore,  $\Upsilon$  meets the conditions of Theorem 2.8, which implies that

$$\mathcal{U} = \{u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M}) : \gamma u \in \Upsilon(u), \gamma > 1\},$$

is bounded. To prove  $\Upsilon$  has a fixed point, let  $u \in \mathcal{U}$ , and  $h \in \mathbb{M}(u)$  such that

$$\begin{aligned} u(\varphi) &= \gamma^{-1}C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] + \gamma^{-1} \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz \\ &+ \gamma^{-1} \int_0^\varphi Q(\varphi, z)h(z)dz + \gamma^{-1} \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \\ &+ \gamma^{-1} \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \gamma^{-1} \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)). \end{aligned}$$

From our assumptions,

$$\begin{aligned} \|u(\varphi)\|_{\mathcal{M}} &\leq \|C(\varphi, 0)u_0\| + \|Q(\varphi, 0)[u_1 - g_1(0, u(0))]\| + \left\| \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz \right\| \\ &+ \left\| \int_0^\varphi Q(\varphi, z)h(z)dz \right\| + \left\| \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \right\| \\ &+ \sum_{0 < \varphi_i < \varphi} \|C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i))\| + \sum_{0 < \varphi_i < \varphi} \|Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i))\| \\ &\leq \mathcal{N}_1\|u_0\| + \mathcal{N}_2\|u_1 - g_1(0, u(0))\| + \mathcal{N}_1 \int_0^\varphi C_2(1 + \|u(z)\|)dz \\ &+ \mathcal{N}_2 \int_0^\varphi [a(z) + e\|u(z)\| + \|\mathcal{B}\|\|v(z)\|]dz + \mathcal{N}_1 \sum_{0 < \varphi_i < \varphi} \|\mathcal{L}_i(u(\varphi_i))\| \\ &+ \mathcal{N}_2 \sum_{0 < \varphi_i < \varphi} \|\mathcal{K}_i(u(\varphi_i))\| \\ &\leq \rho + \mathcal{K}_1\|u\|, \end{aligned} \tag{3.5}$$

where

$$\begin{aligned} \rho &= \mathcal{N}_1[u_0] + \mathcal{N}_2[u_1 - g_1(0, u(0))] + \mathcal{N}_1C_2b + \mathcal{N}_2(\|b\|_{L^2(\mathcal{Q}, \mathbb{R}^+)}) \\ &+ \|\mathcal{B}\|\|v\|_{L^2(\mathcal{Q}, V)}\sqrt{a} + \sum_{i=1}^m l_i + \mathcal{N}_1 \sum_{i=1}^m c_i, \\ \mathcal{K}_1 &= a(\mathcal{N}_1C_2 + \mathcal{N}_2e). \end{aligned}$$

Hence, by the assumption  $\mathcal{K}_1 < 1$  and (3.5), we can see that

$$\|u\| = \sup_{\varphi \in \mathcal{Q}} \|u(\varphi)\| \leq \rho + \mathcal{K}_1\|u\|, \text{ thus } \|u\| \leq \frac{\rho}{1 - \mathcal{K}_1} =: \ell_2.$$

Hence,  $\Upsilon$  has a fixed point.  $\square$

#### 4. Approximate controllability results

Consider that the mild solution for the Eq (2.5) is  $u(\cdot; v)$ , the control variable  $v$  has values in  $L^2(\mathfrak{Q}, V)$ , and the initial value is  $u_0, u_1 \in \mathcal{M}$ . At the terminal time  $a$ , the accessible set of the system (2.5) is defined as  $\mathbb{R}(a, u_0, u_1) = \{u(a; u_0, u_1) : v \in L^2(\mathfrak{Q}, V)\}$ .

**Definition 4.1.** *The Eq (2.5) is approximately controllable on  $\mathfrak{Q}$ , if for any initial value  $u_0, u_1 \in \mathcal{M}$ , then  $\overline{\mathbb{R}(a, u_0, u_1)} = \mathcal{M}$ .*

Consider the linear differential system:

$$\begin{cases} u''(\varphi) = \mathcal{A}(\varphi)u(\varphi) + \mathcal{B}v(\varphi), & \varphi \in \mathfrak{Q} = [0, a], \\ u(0) = u_0, & u'(0) = u_1. \end{cases} \quad (4.1)$$

Now, define the operators for the system (4.1) as:

$$\Upsilon_0^a = \int_0^a Q(a, z)\mathcal{B}\mathcal{B}^*Q^*(a, z)dz \quad \text{and} \quad \mathcal{R}(\beta, \Upsilon_0^a) = (\beta I + \Upsilon_0^a)^{-1}, \quad \beta > 0,$$

where  $\mathcal{B}^*$  and  $Q^*(\varphi)$  are adjoint of  $\mathcal{B}$  and  $Q(\varphi)$ , respectively.

**Lemma 4.2.** [9] *The system (4.1) is approximately controllable on  $\mathfrak{Q}$  iff  $\beta \mathcal{R}(\beta, \Upsilon_0^a) \rightarrow 0$  as  $\beta \rightarrow 0^+$  in the strong operator topology.*

Choose any  $\beta > 0$ ,  $u \in \mathcal{PC}(\mathfrak{Q}, \mathcal{M}) \subset L^2(\mathfrak{Q}, \mathcal{M})$  and  $u_a \in \mathcal{M}$ , as stated in Lemma 2.7, and it is possible to define a multivalued map  $\Upsilon_\beta : \mathcal{PC}(\mathfrak{Q}, \mathcal{M}) \rightarrow 2^{\mathcal{PC}(\mathfrak{Q}, \mathcal{M})}$  given by

$$\begin{aligned} \Upsilon_\beta(u) = \{f \in \mathcal{PC}(\mathfrak{Q}, \mathcal{M}) : f(\varphi) = & C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \\ & + \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz + \int_0^\varphi Q(\varphi, z)h(z)dz \\ & + \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz + \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) \\ & + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)), \quad h \in \mathbb{M}(u)\}, \end{aligned}$$

and

$$\begin{aligned} v_\beta(\varphi) = & \mathcal{B}^*Q^*(a, z)\mathcal{R}(\beta, \Upsilon_0^a)\left(u_a - C(a, 0)u_0 - Q(a, 0)[u_1 - g_1(0, u(0))]\right. \\ & - \int_0^a C(a, z)g_1(z, u(z))dz - \int_0^a Q(a, z)h(z)dz \\ & \left. - \sum_{0 < \varphi_i < a} C(a, \varphi_i)\mathcal{L}_i(u(\varphi_i)) - \sum_{0 < \varphi_i < a} Q(a, \varphi_i)\mathcal{K}_i(u(\varphi_i))\right). \end{aligned}$$

**Theorem 4.3.** *Let (A1)–(A3) be true.  $\Upsilon_\beta$ , for all  $\beta > 0$ , has a fixed point on  $\mathfrak{Q} = [0, a]$  if*

$$a(\mathcal{N}_1\mathcal{C}_2 + \mathcal{N}_2e)\left(1 + \frac{\mathcal{N}_2^2\|\mathcal{B}\|^2}{\beta}\right) < 1,$$

where  $\mathcal{N}_1 := \sup_{\varphi, z \in [0, a]} \|C(\varphi, z)\|$ ,  $\mathcal{N}_2 := \sup_{\varphi, z \in [0, a]} \|Q(\varphi, z)\|$ .

*Proof.* For every  $u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M})$ , by the nature of  $\mathbb{M}(u)$ , we can say  $\Upsilon_\beta$  is convex.

**Step 1.** For every  $p > 0$ ,  $\Upsilon_\beta(\mathfrak{B}_p)$  is bounded in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$ ,

$$\mathfrak{B}_p = \{u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M}) : \|u\|_{\mathcal{PC}} \leq p\}.$$

Here, it is sufficient to prove that there exists a positive constant  $l_\beta$  and for all  $\sigma \in \Upsilon_\beta(u)$ ,  $u \in \mathfrak{B}_p$ ,  $\|\sigma\|_{\mathcal{PC}} \leq l_\beta$ . If  $\sigma \in \Upsilon_\beta(u)$ , there is  $h \in \mathbb{M}(u)$  such that

$$\begin{aligned} \sigma(\varphi) &= C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \\ &+ \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz + \int_0^\varphi Q(\varphi, z)h(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \\ &+ \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)), \quad \varphi \in \mathcal{Q}. \end{aligned}$$

Notice that

$$\begin{aligned} \|v_\beta(\varphi)\| &= \left\| \mathcal{B}^* Q^*(a, z)\mathcal{R}(\beta, \Upsilon_0^a)(u_a - C(a, 0)u_0 - Q(a, 0)[u_1 - g_1(0, u(0))]) \right. \\ &\quad \left. - \int_0^a C(a, z)g_1(z, u(z))dz - \int_0^a Q(a, z)h(z)dz \right. \\ &\quad \left. - \sum_{0 < \varphi_i < a} C(a, \varphi_i)\mathcal{L}_i(u(\varphi_i)) - \sum_{0 < \varphi_i < a} Q(a, \varphi_i)\mathcal{K}_i(u(\varphi_i)) \right\| \\ &\leq \frac{\mathcal{N}_2\|\mathcal{B}\|}{\beta} [\|u_a\| + \mathcal{N}_1\|u_0\| + \mathcal{N}_2\|u_1 - g_1(0, u(0))\| + a\mathcal{N}_1\mathcal{C}_2(1 + p) \\ &\quad + \mathcal{N}_2[\|b\|_{L^2(\mathcal{Q}, \mathbb{R}^+)} \sqrt{a} + epa] + \mathcal{N}_1 \sum_{i=1}^m c_i + \mathcal{N}_2 \sum_{i=1}^m l_i] := \Psi. \end{aligned} \quad (4.2)$$

From (4.2),

$$\begin{aligned} \|\sigma(\varphi)\|_{\mathcal{M}} &\leq \|C(\varphi, 0)u_0\|_{\mathcal{M}} + \|Q(\varphi, 0)[u_1 - g_1(0, u(0))]\|_{\mathcal{M}} + \left\| \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz \right\| \\ &\quad + \left\| \int_0^\varphi Q(\varphi, z)h(z)dz \right\| + \left\| \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \right\| \\ &\quad + \sum_{0 < \varphi_i < \varphi} \|C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i))\| + \sum_{0 < \varphi_i < \varphi} \|C(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i))\| \\ &\leq \mathcal{N}_1\|u_0\| + \mathcal{N}_2\|u_1 - g_1(0, u(0))\| + \mathcal{N}_1a\mathcal{C}_2(1 + p) + \mathcal{N}_2[\sqrt{a}\|b\|_{L^2(\mathcal{Q}, \mathbb{R}^+)} + epa \\ &\quad + \|\mathcal{B}\|\Psi a] + \mathcal{N}_1 \sum_{i=1}^m c_i + \mathcal{N}_2 \sum_{i=1}^m l_i := l_\beta. \end{aligned}$$

Thus  $\Upsilon_\beta(\mathfrak{B}_p)$  is bounded in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$ .

**Step 2.** Consider any  $u \in \mathfrak{B}_p$ ,  $\sigma \in \Upsilon_\beta(u)$ . There exists  $h \in \mathbb{M}(u)$ , for every  $\varphi \in \mathcal{Q}$ ,

$$\begin{aligned}\sigma(\varphi) &= C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \\ &+ \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz + \int_0^\varphi Q(\varphi, z)h(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}v(z)dz \\ &+ \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)).\end{aligned}$$

Using  $\|v_\beta(t)\|$  as (4.2) and also from Theorem 3.1, Step 2, one can obtain that  $\{\Upsilon_\beta(u) : u \in \mathfrak{B}_p\}$  is completely continuous.

**Step 3.** Consider  $u_n \rightarrow u_*$  in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$ ,  $\sigma_n \in \Upsilon_\beta(u_n)$  and  $\sigma_n \rightarrow \sigma_*$  in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$ . We investigate  $\sigma_* \in \Upsilon_\beta(u_*)$ . Indeed,  $\sigma_n \in \Upsilon_\beta(u_n)$  exists only when  $h_n \in \mathbb{M}(u_n)$  such that

$$\begin{aligned}\sigma_n(\varphi) &= C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \\ &+ \int_0^\varphi C(\varphi, z)g_1(z, u_n(z))dz + \int_0^\varphi Q(\varphi, z)h_n(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}\mathcal{B}^*Q^*(a, z)\mathcal{R}(\beta, \Upsilon_0^a) \\ &(\times)(u_a - C(a, 0)u_0 - Q(a, 0)[u_1 - g_1(0, u(0))]) - \int_0^a C(a, \eta)g_1(\eta, u_n(\eta))d\eta \\ &- \int_0^a Q(a, \eta)h_n(\eta)d\eta - \sum_{0 < \varphi_i < a} C(a, \varphi_i)\mathcal{L}_i(u(\varphi_i)) - \sum_{0 < \varphi_i < a} Q(a, \varphi_i)\mathcal{K}_i(u(\varphi_i))dz \\ &+ \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)).\end{aligned}\quad (4.3)$$

From (A1), we will prove  $\{h_n\}_{n \geq 1} \subseteq L^2(\mathcal{Q}, \mathcal{M})$  is bounded. Hence,

$$h_n \rightarrow h_* \text{ weakly in } L^2(\mathcal{Q}, \mathcal{M}).\quad (4.4)$$

From (4.3) and (4.4),

$$\begin{aligned}\sigma_n(\varphi) &\rightarrow C(\varphi, 0)u_0 + Q(\varphi, 0)[z_1 - g_1(0, z(0))] \\ &+ \int_0^\varphi C(\varphi, z)g_1(z, u_*(z))dz + \int_0^\varphi Q(\varphi, z)h_*(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}\mathcal{B}^*Q^*(a, z)\mathcal{R}(\beta, \Upsilon_0^a) \\ &(\times)(u_a - C(a, 0)u_0 - Q(a, 0)[u_1 - g_1(0, u(0))]) - \int_0^a C(a, \eta)g_1(\eta, u_*(\eta))d\eta \\ &- \int_0^a Q(a, \eta)h_*(\eta)d\eta - \sum_{0 < \varphi_i < a} C(a, \varphi_i)\mathcal{L}_i(u(\varphi_i)) - \sum_{0 < \varphi_i < a} Q(a, \varphi_i)\mathcal{K}_i(u(\varphi_i))dz \\ &+ \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)).\end{aligned}\quad (4.5)$$

Clearly,  $\sigma_n \rightarrow \sigma_*$  in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$  and  $h_n \in \mathbb{M}(u_n)$ . According to Lemma 2.6 and (4.5),  $h_* \in \mathbb{M}(u_*)$ . Then,  $\sigma_* \in \Upsilon(u_*)$ , and this shows that  $\Upsilon$  has a closed graph. Hence, by using Proposition 3.12 of [36] implies that it is upper semicontinuous.

**Step 4.** A priori estimate.

By Steps 1–3,  $\Upsilon_\beta$  is convex valued, compact upper semicontinuous,  $\Upsilon_\beta(\mathfrak{B}_p)$  is a relatively compact set and meets Theorem 2.8, and

$$\mathbb{J} = \{u \in \mathcal{PC}(\mathcal{Q}, \mathcal{M}) : \gamma u \in \Upsilon_\beta(u), \gamma > 1\},$$

is bounded.

Consider  $u \in \mathbb{J}$ . Then there exists  $h \in \mathcal{M}(u)$  such that

$$\begin{aligned} u(\varphi) &= \gamma^{-1}C(\varphi, 0)u_0 + \gamma^{-1}Q(\varphi, 0)[u_1 - g_1(0, u(0))] + \gamma^{-1} \int_0^\varphi C(\varphi, z)g_1(\varphi, u(\varphi))dz \\ &\quad + \gamma^{-1} \int_0^\varphi Q(\varphi, z)h(z)dz + \gamma^{-1} \int_0^\varphi Q(\varphi, z)\mathcal{B}v_\beta(z)dz \\ &\quad + \gamma^{-1} \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \gamma^{-1} \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)), \end{aligned}$$

and

$$\begin{aligned} v_\beta(\varphi) &= \mathcal{B}^*Q^*(a, z)\mathcal{R}(\beta, \Upsilon_0^a)(u_a - C(a, 0)u_0 - Q(a, 0)[u_1 - g_1(0, u(0))]) \\ &\quad - \int_0^a C(a, z)g_1(z, u(z))dz - \int_0^a Q(a, z)h(z)dz \\ &\quad - \sum_{0 < \varphi_i < a} C(a, \varphi_i)\mathcal{L}_i(u(\varphi_i)) - \sum_{0 < \varphi_i < a} Q(a, \varphi_i)\mathcal{K}_i(u(\varphi_i)). \end{aligned}$$

Then from our assumptions,

$$\begin{aligned} \|u(\varphi)\|_{\mathcal{M}} &\leq \|C(\varphi, 0)u_0\| + \|Q(\varphi, 0)[u_1 - g_1(0, u(0))]\| + \left\| \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz \right\| \\ &\quad + \left\| \int_0^\varphi Q(\varphi, z)h(z)dz \right\| + \left\| \int_0^\varphi Q(\varphi, z)\mathcal{B}v_\beta(z)dz \right\| \\ &\quad + \sum_{0 < \varphi_i < \varphi} \|C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i))\| + \sum_{0 < \varphi_i < \varphi} \|Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i))\| \\ &\leq \mathcal{N}_1\|z_0\| + \mathcal{N}_2\|[u_1 - g_1(0, u(0))]\| + \mathcal{N}_1 \int_0^\varphi C_2(1 + \|u(\varphi)\|)dz + \mathcal{N}_2 \int_0^\varphi [b(z) + e\|u(z)\| \\ &\quad + \|\mathcal{B}\| \left( \frac{\mathcal{N}_2\|\mathcal{B}\|}{\beta} [\|u_a\| + \mathcal{N}_1\|u_0\| + \mathcal{N}_2\|u_1 - g_1(0, z(0))\| + a\mathcal{N}_1C_2(1 + p) \right. \\ &\quad \left. + \mathcal{N}_2[\|b\|_{L^2(\mathcal{Q}, \mathbb{R}^+)} \sqrt{a} + epa] + \mathcal{N}_1 \sum_{i=1}^m c_i + \mathcal{N}_2 \sum_{i=1}^m l_i \right)] dz \\ &\quad + \mathcal{N}_1 \sum_{0 < \varphi_i < \varphi} \|\mathcal{L}_i(u(\varphi_i))\| + \mathcal{N}_2 \sum_{0 < \varphi_i < \varphi} \|\mathcal{K}_i(u(\varphi_i))\| \\ &\leq \rho + \mathcal{K}_2\|u\|, \end{aligned} \tag{4.6}$$

where

$$\begin{aligned} \rho &= \mathcal{N}_1 \|u_0\| + \mathcal{N}_2 \| [u_1 - g_1(0, u(0))] \| + \mathcal{N}_1 C_2 a + \mathcal{N}_2 [ \|b\|_{L^2(\mathcal{Q}, \mathbb{R}^+)} ] \sqrt{a} \\ &\quad + \frac{\mathcal{N}_2^2 \| \mathcal{B} \|^2}{\beta} \left( \|u_a\| + \mathcal{N}_1 \|u_0\| + \mathcal{N}_2 \| [u_1 - g_1(0, u(0))] \| \right. \\ &\quad \left. + \mathcal{N}_1 C_2 a + \mathcal{N}_2 [ \|b\|_{L^2(\mathcal{Q}, \mathbb{R}^+)} ] \sqrt{a} \right) + \mathcal{N}_1 \sum_{i=1}^m c_i + \mathcal{N}_2 \sum_{i=1}^m l_i. \\ \mathcal{K}_2 &= a(\mathcal{N}_1 C_2 + \mathcal{N}_2 e) \left( 1 + \frac{\mathcal{N}_2^2 \| \mathcal{B} \|^2}{\beta} \right). \end{aligned}$$

According to  $\mathcal{K}_2 < 1$  and (4.6), we conclude,

$$\|u\| = \sup_{\varphi \in \mathcal{Q}} \|u(\varphi)\| \leq \rho + \mathcal{K}_2 \|u\|, \text{ and thus } \|u\| \leq \frac{\rho}{1 - \mathcal{K}_2} =: \ell_3.$$

Therefore  $\mathbb{J}$  is bounded which leads to the conclusion that  $\Upsilon_\beta$  has a fixed point.  $\square$

**Theorem 4.4.** *Suppose the conditions of the above theorem are satisfied. Then, if system (4.1) is approximately controllable on the set  $\mathcal{Q}$ , it follows that system (2.5) is also approximately controllable on  $\mathcal{Q}$ .*

*Proof.* By Theorem 4.3,  $\Upsilon_\beta$ , for all  $\beta > 0$ , has a fixed point in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$ . Let  $u^\beta$  be a fixed point of  $\Upsilon_\beta$  in  $\mathcal{PC}(\mathcal{Q}, \mathcal{M})$ . Clearly,  $\Upsilon_\beta$  is a mild solution of (2.5). Then, there exists  $h^\beta \in \mathbb{M}(u^\beta)$  such that for each  $\varphi \in \mathcal{Q}$ ,

$$\begin{aligned} u^\beta(\varphi) &= C(\varphi, 0)u_0 + Q(\varphi, 0)[u_1 - g_1(0, u(0))] \\ &\quad + \int_0^\varphi C(\varphi, z)g_1(z, u(z))dz + \int_0^\varphi Q(\varphi, z)h^\beta(z)dz + \int_0^\varphi Q(\varphi, z)\mathcal{B}\mathcal{B}^*Q^*(a, z)\mathcal{R}(\beta, \Upsilon_0^a) \\ &\quad (\times) \left( u_a - C(a, 0)u_0 - Q(a, 0)[u_1 - g_1(0, u(0))] - \int_0^a C(a, \eta)g_1(\eta, u(\eta))d\eta \right. \\ &\quad \left. - \int_0^a Q(a, \eta)h^\beta(\eta)d\eta - \sum_{0 < \varphi_i < a} C(a, \varphi_i)\mathcal{L}_i(u(\varphi_i)) - \sum_{0 < \varphi_i < a} Q(a, \varphi_i)\mathcal{K}_i(u(\varphi_i)) \right) du \\ &\quad + \sum_{0 < \varphi_i < \varphi} C(\varphi, \varphi_i)\mathcal{L}_i(u(\varphi_i)) + \sum_{0 < \varphi_i < \varphi} Q(\varphi, \varphi_i)\mathcal{K}_i(u(\varphi_i)). \end{aligned}$$

Since  $I - \Upsilon_0^a \mathcal{R}(\beta, \Upsilon_0^a) = \beta \mathcal{R}(\beta, \Upsilon_0^a)$ , we have  $u^\beta(a) = u_a - \beta \mathcal{R}(\beta, \Upsilon_0^a) \mathbb{E}(h^\beta)$ . From the above,

$$\begin{aligned} \mathbb{E}(h^\beta) &= u_a - C(a, 0)u_0 - Q(a, 0)[u_1 - g_1(0, u(0))] - \int_0^a C(a, \eta)g_1(\eta, u(\eta))d\eta \\ &\quad - \int_0^a Q(a, \eta)h^\beta(\eta)d\eta - \sum_{0 < \varphi_i < a} C(a, \varphi_i)\mathcal{L}_i(u(\varphi_i)) - \sum_{0 < \varphi_i < a} Q(a, \varphi_i)\mathcal{K}_i(u(\varphi_i)). \end{aligned}$$

From the hypothesis **(A1)** and from Theorem 4.3,  $\|\partial H(\varphi, u)\| \leq b(\varphi) + e\|u(\varphi)\| \leq b(\varphi) + e\rho := \nu(\varphi)$ . Then,

$$\int_0^a \|h^\beta(z)\| dz \leq \|\nu\|_{L^2(\mathcal{Q}, \mathbb{R}^+)} \sqrt{a}.$$

Consequently  $\{h^\beta\}$  is a bounded sequence in  $L^2(\mathcal{Q}, \mathcal{M})$ . So, there exists a subsequence,  $\{h^\beta\}$ , which will converge weakly to  $h$  in  $L^2(\mathcal{Q}, \mathcal{M})$ . It is expressed as

$$g = u_a - C(a, 0)u_0 - Q(a, 0)[u_1 - g_1(0, u(0))] - \int_0^a C(a, \eta)g_1(\eta, u(\eta))d\eta \\ - \int_0^a Q(a, \eta)h(\eta)d\eta - \sum_{0 < \varphi_i < a} C(a, \varphi_i)\mathcal{L}_i(u(\varphi_i)) - \sum_{0 < \varphi_i < a} Q(a, \varphi_i)\mathcal{K}_i(u(\varphi_i)).$$

Now,

$$\|\mathbb{E}(h^\beta) - g\| = \left\| \int_0^a C(a, \eta)[g_1(\eta, y^\beta(\eta)) - g_1(\eta, y(\eta))]d\eta \right\| + \left\| \int_0^a \mathcal{S}(a, \eta)[h^\beta(\eta) - h(\eta)]d\eta \right\| \\ \leq \mathcal{M}_1 \sup_{0 \leq \eta \leq a} [g_1(\eta, y^\beta(\eta)) - g_1(\eta, y(\eta))] + \mathcal{M}_2 \sup_{0 \leq \eta \leq a} [h^\beta(\eta) - h(\eta)]. \quad (4.7)$$

From Step 2 in Theorem 4.3 and by the Arzelà-Ascoli theorem, we get that the compactness of the right-hand side of (4.7) tends to zero as  $\beta \rightarrow 0^+$ , which gives

$$\|u^\beta(a) - u_1\| = \|\beta \mathcal{R}(\beta, \Upsilon_0^a)\mathbb{E}(h^\beta)\| \\ \leq \|\beta \mathcal{R}(\beta, \Upsilon_0^a)(g)\| + \|\mathbb{E}(h^\beta) - g\| \rightarrow 0, \text{ as } \beta \rightarrow 0^+.$$

Hence, (2.5) is approximately controllable on  $\mathcal{Q}$ . □

## 5. Applications

We utilize our theoretical findings on a concrete partial differential equation. We need to provide the required technological resources to accomplish our goals.

Now, let us take

$$\mathcal{A}(\varphi) = \mathcal{A} + \tilde{\mathcal{A}}(\varphi),$$

where  $\mathcal{A}$  is the infinitesimal generator of a cosine function  $C(\varphi)$  with associated sine function  $Q(t)$ , and  $\tilde{\mathcal{A}}(\varphi) : D(\tilde{\mathcal{A}}(\varphi)) \rightarrow \mathcal{M}$  is a closed linear operator with  $D \subset D(\tilde{\mathcal{A}}(\varphi))$ , for all  $\varphi \in \mathcal{Q}$ . We take the space  $\mathcal{M} = L^2(\mathbb{T}, \mathbb{C})$ , where the group  $\mathbb{T}$  is defined as the quotient  $\mathbb{R}/2\pi\mathbb{Z}$ , and we denote by  $L^2(\mathbb{T}, \mathbb{C})$  the space of  $2\pi$  periodic 2-integrable functions from  $\mathbb{R}$  to  $\mathbb{C}$ . Also, we use the identification between functions on  $\mathbb{T}$  and  $2\pi$  periodic functions on  $\mathbb{R}$ . Furthermore,  $H^2(\mathbb{T}, \mathbb{C})$  denotes the Sobolev space of  $2\pi$  periodic from  $\mathbb{R}$  to  $\mathbb{C}$  such that  $u'' \in L^2(\mathbb{T}, \mathbb{C})$ .

We define  $\mathcal{A}u(\varphi) = u''(\varphi)$  with domain  $D(\mathcal{A}) = H^2(\mathbb{T}, \mathbb{C})$ . Then,  $\mathcal{A}$  can be written as

$$\mathcal{A}u = \sum_{n=1}^{\infty} -n^2 \langle u, x_n \rangle x_n, u \in D(A),$$

where  $x_n(\varphi) = \frac{1}{\sqrt{2\pi}} e^{in\varphi}$  ( $n \in \mathbb{Z}$ ) is an orthonormal basis of  $\mathcal{M}$ . It is well known that  $\mathcal{A}$  is the infinitesimal generator of a strongly continuous cosine function  $C(\varphi)$  on  $\mathcal{M}$ . The cosine function  $C(\varphi)$  is given by

$$C(\varphi)u = \sum_{n=1}^{\infty} \cos nt \langle u, x_n \rangle x_n, u \in \mathcal{M}, \varphi \in \mathbb{R}.$$



The connected sine operator  $(Q(\varphi))_{\varphi \in \mathbb{R}}$  is

$$Q(t)u = \sum_{n=1}^{\infty} \frac{\sin nt}{n} \langle u, x_n \rangle x_n, \quad u \in \mathcal{M}, \quad \varphi \in \mathbb{R}.$$

It is clear that  $\|C(\varphi)\| \leq 1$  for all  $\varphi \in \mathbb{R}$ , so it is uniformly bounded on  $\mathbb{R}$ .

Assume the following second-order non-autonomous neutral differential system of the form:

$$\begin{aligned} \frac{\partial}{\partial \varphi} \left[ \frac{\partial}{\partial \varphi} u(\varphi, \wp) - \hat{g}_1(\varphi, y(\varphi, \wp)) \right] &= \frac{\partial^2}{\partial \varphi^2} u(\varphi, \wp) + \mathbb{B}(\varphi) \frac{\partial}{\partial \varphi} u(\varphi, \wp) \\ &\quad + \mathcal{B}\tilde{v}(\varphi, \wp) + \phi(\varphi, u(\varphi, \wp)), \quad 0 < \varphi < a, \\ u(\varphi, 0) &= u(\varphi, \pi) = 0, \quad 0 < \varphi < a, \\ u(0, \wp) &= u_0(\wp), \quad \wp \in (0, \pi), \\ \frac{\partial}{\partial \varphi} y(0, \wp) &= u_1(\wp), \\ \Delta u(\varphi_k)(\wp) &= \int_0^{\varphi_k} b_k(\varphi_k - \varphi) u(\varphi, \wp) d\varphi, \\ \Delta \frac{\partial}{\partial \varphi} u(\varphi_k)(\wp) &= \int_0^{\varphi_k} b_k(\varphi_k - \varphi) u_1(\varphi, \wp) d\varphi, \end{aligned} \tag{5.1}$$

where  $\mathbb{B} : \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function such that  $\sup_{\varphi \in [0, a]} \|\mathbb{B}(\varphi)\| = c_0$ , and  $u(\varphi, \wp)$  represents the temperature at  $\varphi \in (0, \pi)$  and  $\varphi \in (0, a)$ . Let  $\phi(\varphi, u(\varphi, \wp)) = \phi_1(\xi_1, \varphi(\varphi, \wp)) + \phi_2(\varphi, u(\varphi, \wp))$  and  $\phi_2(\varphi, u(\varphi, \wp))$  is the temperature function of the form  $-\phi_2(\varphi, u(\varphi, \wp)) \in \partial H(\varphi, \wp, u(\varphi, \wp))$ ,  $(\varphi, \wp) \in (0, a) \times (0, \pi)$ . Here, the nonsmooth and nonconvex function  $H = H(\varphi, \wp, k)$  is defined as a locally Lipschitz energy function.  $\partial H$  is the generalized Clarke's gradient in the third variable  $k$  [6]. Assume that  $H$  fulfills the assumptions **(A1)**,  $H(k) = \min\{h_1(v), h_2(v)\}$ , and  $h_i = \mathbb{R} \rightarrow \mathbb{R}$  ( $i = 1, 2$ ) are convex quadratic functions [16].

Now we take  $\tilde{\mathcal{A}}(\varphi)u(\varphi) = \mathbb{B}(\varphi)u(\varphi)$  defined on  $H^1(\mathbb{T}, \mathbb{C})$ . It is easy to see that  $\mathcal{A}(\varphi) = \mathcal{A} + \tilde{\mathcal{A}}(\varphi)$  is a closed linear operator. Initially, we will show that  $\mathcal{A} + \tilde{\mathcal{A}}(\varphi)$  generates an evolution operator. It is well known that the solution of the scalar initial value problem

$$\begin{aligned} p''(\varphi) &= -n^2 p(\varphi) + q(\varphi), \\ p(s) &= 0, \quad p'(s) = p_1, \end{aligned}$$

is given by

$$p(t) = \frac{p_1}{n} \sin n(\varphi - z) + \frac{1}{n} \int_z^\varphi \sin n(\varphi - \iota) q(\iota) d\iota.$$

Therefore, the solution of the scalar initial value problem

$$p''(\varphi) = -n^2 p(\varphi) + in\mathbb{B}(\varphi)p(\varphi), \tag{5.2}$$

$$p(z) = 0, \quad p'(z) = p_1, \tag{5.3}$$

satisfies the following equation:

$$p(\varphi) = \frac{p_1}{n} \sin n(\varphi - z) + i \int_z^\varphi \sin n(\varphi - t) \mathbb{B}(t) q(t) dt.$$

By the Gronwall-Bellman lemma, we obtain

$$|p(t)| \leq \frac{p_1}{n} e^{c(\varphi-z)} \quad (5.4)$$

for  $z \leq \varphi$  and  $c$  is a constant. We denote by  $p_n(\varphi, z)$  the solution of (5.2)-(5.3). We define

$$Q(\varphi, z)u = \sum_{n=1}^{\infty} p_n(\varphi, z) \langle u, x_n \rangle x_n, \quad u \in \mathcal{M}, \quad \varphi \in \mathbb{R}.$$

It follows from the estimate (5.4) that  $Q(\varphi, z) : \mathcal{M} \rightarrow \mathcal{M}$  is well defined and satisfies the condition of Definition 2.1. We set  $u(t) = u(\varphi, \cdot)$ , that is,  $u(\varphi)(\varphi) = u(\varphi, \varphi)$ ,  $\varphi \in \mathcal{Q}$ ,  $\varphi \in [0, \pi]$ . Then, we assume the infinite dimensional Hilbert space  $\mathcal{V}$ , and we have

$$\mathcal{V} = \left\{ v : v \in \sum_{j=1}^{\infty} v_j x_j \text{ with } \sum_{j=2}^{\infty} v_j^2 < \infty \right\},$$

with  $\mathcal{V}$  as

$$\|v\|_{\mathcal{V}} = \left( \sum_{j=2}^{\infty} v_j^2 \right)^{\frac{1}{2}}.$$

Let us define  $\mathcal{B} \in \mathcal{L}(\mathcal{V}, \mathcal{M})$  as below:

$$\mathcal{B}v = 2v_2 x_1 + \sum_{j=2}^{\infty} v_j x_j < \infty, \text{ for all } v = \sum_{j=2}^{\infty} v_j x_j \in \mathcal{V}.$$

It continuous that

$$\mathcal{B}^*v = (2v + v_2)x_2 + \sum_{j=3}^{\infty} v_j x_j, \text{ for all } v = \sum_{j=2}^{\infty} v_j x_j \in \mathcal{M}.$$

Assume these functions satisfy the requirements of the hypotheses. From the above choices of the functions and evolution operator  $\mathcal{A}(\varphi)$  with  $\mathcal{B}$ , system (5.1) can be formulated as system (2.5) in  $\mathcal{M}$ . Since all hypotheses of Theorem 4.4 are satisfied, the approximate controllability of system (5.1) on  $\mathcal{Q}$  follows from Theorem 4.4.

## 6. Conclusions

The principles of approximate controllability of second-order differential impulsive systems with the impact of hemivariational inequalities are the main focus of this article. The generalized Clarke's subdifferential technique and multivalued maps were used to suggest and demonstrate the necessary requirements for existence and approximate controllability. In the future, we will extend the results with finite delay and stochastic systems.

## Author contributions

Yong-Ki Ma: Conceptualization, Methodology, Validation, Visualization, Writing–original draft. N. Valliammal: Conceptualization, Formal analysis, Methodology, Validation, Visualization, Writing–original draft. K. Jothimani: Conceptualization, Formal analysis, Investigation, Resources, Supervision, Writing–original draft. V. Vijayakumar: Conceptualization, Formal analysis, Resources, Supervision, Writing–original draft, Writing–Review & Editing. All authors have read and approved the final version of the manuscript for publication.

## Acknowledgments

This work was supported by the research grant of Kongju National University in 2024. The authors are immensely grateful to the anonymous referees for their careful reading of this paper and helpful comments, which have been very useful for improving the quality of this paper.

## Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

1. Y. Huang, Z. Liu, B. Zeng, Optimal control of feedback control systems governed by hemivariational inequalities, *Comput. Math. Appl.*, **70** (2015), 2125–2136. <https://doi.org/10.1016/j.camwa.2015.08.029>
2. P. D. Panagiotopoulos, *Hemivariational inequalities: applications in mechanics and engineering*, Berlin: Springer, 1993. <https://doi.org/10.1007/978-3-642-51677-1>
3. P. D. Panagiotopoulos, Non-convex superpotentials in the sense of F.H. Clarke and applications, *Mech. Res. Commun.*, **8** (1981), 335–340. [https://doi.org/10.1016/0093-6413\(81\)90064-1](https://doi.org/10.1016/0093-6413(81)90064-1)
4. S. Carl, D. Motreanu, Extremal solutions of quasilinear parabolic inclusions with generalized Clarke's gradient, *J. Differ. Equations*, **191** (2003), 206–233. [https://doi.org/10.1016/S0022-0396\(03\)00022-6](https://doi.org/10.1016/S0022-0396(03)00022-6)
5. S. Carl, Existence of extremal solutions of boundary hemivariational inequalities, *J. Differ. Equations*, **171** (2001), 370–396. <https://doi.org/10.1006/jdeq.2000.3845>
6. F. H. Clarke, *Optimization and nonsmooth analysis*, Wiley, 1990. <https://doi.org/10.1137/1.9781611971309>
7. Z. Denkowski, S. Migórski, N. S. Papageorgiou, *An introduction to non-linear analysis: theory*, New York: Springer, 2003. <https://doi.org/10.1007/978-1-4419-9158-4>
8. Z. Liu, N. S. Papageorgiou, Double phase Dirichlet problems with unilateral constraints, *J. Differ. Equations*, **316** (2022), 249–269. <https://doi.org/10.1016/j.jde.2022.01.040>
9. Z. H. Liu, X. Li, Approximate controllability for a class of hemivariational inequalities, *Nonlinear Anal.*, **22** (2015), 581–591. <https://doi.org/10.1016/j.nonrwa.2014.08.010>

10. F. S. Acharya, N. C. Dimplekumar, Controllability of neutral impulsive differential inclusions with nonlocal conditions, *Appl. Math.*, **2** (2011), 1486–1496. <https://doi.org/10.4236/am.2011.212211>
11. M. Benchohra, J. Henderson, S. K. Ntouyas, Existence results for impulsive multivalued semilinear neutral functional differential inclusions in Banach spaces, *J. Math. Anal. Appl.*, **263** (2001), 763–780. <https://doi.org/10.1006/jmaa.2001.7663>
12. X. Hao, L. Liu, Y. Wu, Mild solutions of impulsive semilinear neutral evolution equations in Banach spaces, *J. Nonlinear Sci. Appl.*, **9** (2016), 6183–6194. <https://doi.org/10.22436/jnsa.009.12.23>
13. N. Valliammal, K. Jothimani, S. K. Panda, V. Vijayakumar, An investigation on the existence and approximate controllability of neutral stochastic hemivariational inequalities, *Rend. Circ. Mat. Palermo, II. Ser.*, **73** (2024), 941–958. <https://doi.org/10.1007/s12215-023-00967-4>
14. J. Haslinger, P. D. Panagiotopoulos, Optimal control of systems governed by hemivariational inequalities. Existence and approximation results, *Nonlinear Anal.*, **24** (1995), 105–119. [https://doi.org/10.1016/0362-546X\(93\)E0022-U](https://doi.org/10.1016/0362-546X(93)E0022-U)
15. Y. Liu, Z. Liu, N. S. Papageorgiou, Sensitivity analysis of optimal control problems driven by dynamic history-dependent variational-hemivariational inequalities, *J. Differ. Equations*, **342** (2023), 559–595. <https://doi.org/10.1016/j.jde.2022.10.009>
16. S. Migórski, On existence of solutions for parabolic hemivariational inequalities, *J. Comput. Appl. Math.*, **129** (2001), 77–87. [https://doi.org/10.1016/S0377-0427\(00\)00543-4](https://doi.org/10.1016/S0377-0427(00)00543-4)
17. J. Zhao, L. Liu, E. Vilches, C. Wen, J. C. Yao, Optimal control of an evolution hemivariational inequality involving history-dependent operators, *Commun. Nonlinear Sci. Numer. Simul.*, **103** (2021), 105992. <https://doi.org/10.1016/j.cnsns.2021.105992>
18. J. Y. Park, S. H. Park, Optimal control problems for anti-periodic quasi-linear hemivariational inequalities, *Optim. Control Appl. Met.*, **28** (2007), 275–287. <https://doi.org/10.1002/oca.803>
19. J. Y. Park, S. H. Park, Existence of solutions and optimal control problems for hyperbolic hemivariational inequalities, *ANZIAM J.*, **47** (2005), 51–63. <https://doi.org/10.1017/S1446181100009767>
20. X. Li, Z. Liu, N. S. Papageorgiou, Solvability and pullback attractor for a class of differential hemivariational inequalities with its applications, *Nonlinearity*, **36** (2023), 1323–1348. <https://doi.org/10.1088/1361-6544/acb191>
21. Y. K. Ma, J. Pradeesh, A. Shukla, V. Vijayakumar, K. Jothimani, An analysis on the approximate controllability of neutral impulsive stochastic integrodifferential inclusions via resolvent operators, *Heliyon*, **9** (2023), e20837. <https://doi.org/10.1016/j.heliyon.2023.e20837>
22. J. Chen, Z. Liu, F. E. Lomovtsev, V. Obukhovskii, Optimal feedback control for a class of second-order evolution differential inclusions with Clarke’s subdifferential, *J. Nonlinear Var. Anal.*, **6** (2022), 551–565. <https://doi.org/10.23952/jnva.6.2022.5.08>
23. Y. Liu, Z. Liu, S. Peng, C. Wen, Optimal feedback control for a class of fractional evolution equations with history-dependent operators, *Fract. Calc. Appl. Anal.*, **25** (2022), 1108–1130. <https://doi.org/10.1007/s13540-022-00054-y>

24. J. Pradeesh, V. Vijayakumar, Investigating the existence results for Hilfer fractional stochastic evolution inclusions of order  $1 < \mu < 2$ , *Qual. Theory Dyn. Syst.*, **23** (2024), 46. <https://doi.org/10.1007/s12346-023-00899-5>
25. K. Rykaczewski, Approximate controllability of differential inclusions in Hilbert spaces, *Nonlinear Anal.*, **75** (2012), 2701–2712. <https://doi.org/10.1016/j.na.2011.10.049>
26. V. Vijayakumar, Approximate controllability for a class of second-order stochastic evolution inclusions of Clarke's subdifferential type, *Results Math.*, **73** (2018), 1–17. <https://doi.org/10.1007/s00025-018-0807-8>
27. J. Pradeesh, V. Vijayakumar, A new approach on the approximate controllability results for Hilfer fractional stochastic hemivariational inequalities of order  $1 < \mu < 2$ , *Qual. Theory Dyn. Syst.*, **23** (2024), 158. <https://doi.org/10.1007/s12346-024-01012-0>
28. C. J. K. Batty, R. Chill, S. Srivastava, Maximal regularity for second order non-autonomous Cauchy problems, *Studia Math.*, **189** (2008), 205–223.
29. F. Faraci, A. Iannizzotto, A multiplicity theorem for a perturbed second-order non-autonomous system, *Proc. Edinburgh Math. Soc.*, **49** (2006), 267–275. <https://doi.org/10.1017/S001309150400149X>
30. E. Obrecht, Evolution operators for higher order abstract parabolic equations, *Czech. Math. J.*, **36** (1986), 210–222. <https://doi.org/10.21136/CMJ.1986.102085>
31. M. Kozak, A fundamental solution of a second order differential equation in a Banach space, *Univ. Lagel. Acta Math.*, **32** (1995), 275–289.
32. S. Hu, N. S. Papageorgiou, *Handbook of multivalued analysis, volume I: theory*, New York: Springer, 1997.
33. A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, New York: Springer-Verlag, 1983. <https://doi.org/10.1007/978-1-4612-5561-1>
34. T. W. Ma, *Topological degrees of set-valued compact fields in locally convex spaces*, Warszawa: Instytut Matematyczny Polskiej Akademi Nauk, 1972.
35. S. Migórski, A. Ochal, Quasi-static hemivariational inequality via vanishing acceleration approach, *SIAM J. Control Optim.*, **41** (2009), 1415–1435. <https://doi.org/10.1137/080733231>
36. S. Migórski, A. Ochal, M. Sofonea, *Nonlinear inclusions and hemivariational inequalities: models and analysis of contact problems*, Advances in Mechanics and Mathematics, Vol. 26, New York: Springer, 2013. <https://doi.org/10.1007/978-1-4614-4232-5>



AIMS Press

©2024 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>)