



Research article

Global stability of a tridiagonal competition model with seasonal succession

Xizhuang Xie* and **Meixiang Chen**

School of Mathematical Sciences, Huaqiao University, Quanzhou 362021, China

* **Correspondence:** Email: xzx@hqu.edu.cn.

Abstract: In this paper, we investigate a tridiagonal three-species competition model with seasonal succession. The Floquet multipliers of all nonnegative periodic solutions of such a time-periodic system are estimated via the stability analysis of equilibria. Together with the Brouwer degree theory, sufficient conditions for existence and uniqueness of the positive periodic solution are given. We further obtain the global dynamics of coexistence and extinction for three competing species in this periodically forced environment. Finally, some numerical examples are presented to illustrate the effectiveness of our theoretical results.

Keywords: tridiagonal competition model; seasonal succession; carrying simplex; Poincaré-map; global dynamics

1. Introduction

The far-reaching consequences of ecological interactions in the dynamics of biological communities remain an intriguing subject. The interaction between two species can basically be of three different kinds: competition, mutualism and predator-prey. Competition and predator-prey are, perhaps, the most well-known types of interactions. They embody the natural dispute for resources in ecology. Competition is characterized by a decrease of the growth rate as the density of the other species increases, which has been extensively studied (see [1–4]). One special competition example occurs in the water column of an ocean or on a steep mountain side or on island groups, where each species dominates a species zone (depth, altitude or different island) but is obliged to interact with other species in the narrow overlap of their zones of dominance. Taking three species as an example, one can think of a hierarchy of species x_1 , x_2 and x_3 , where x_i is the density or biomass of the i -th species. In this hierarchy, x_1 only competes with x_2 , x_3 only with x_2 , and x_2 competes with x_1 and x_3 .

The concrete form is as follows,

$$\begin{cases} \frac{dx_1}{dt} = x_1(r_1 - a_{11}x_1 - a_{12}x_2), \\ \frac{dx_2}{dt} = x_2(r_2 - a_{21}x_1 - a_{22}x_2 - a_{23}x_3), \\ \frac{dx_3}{dt} = x_3(r_3 - a_{32}x_2 - a_{33}x_3), \\ (x_1(0), x_2(0), x_3(0)) = x^0 \in \mathbb{R}_+^3, \end{cases} \quad (1.1)$$

where r_i and a_{ij} are all positive real numbers. Such system is called tridiagonal competitive system and there have also been a number of theoretical works (see [5–8]).

As we know, interactive species often live in a fluctuating environment. Due to seasonal or daily variations, species experience a periodic dynamical environment such as temperature, rainfall, humidity, wind, and other resources. A typical example here is the phytoplankton and zooplankton of the temperate lakes, where the species grow during the warmer months and die off or form resting stages in winter. The phenomenon is described as seasonal succession. It has been a fascinating subject for ecologists and mathematicians to explore the dynamics of periodic models by means of seasonal succession numerically and analytically. Recently, Klausmeier [9] proposed a novel approach, called successional state dynamics (SSD), to modeling seasonally forced food webs. The SSD approach treats succession as a series of state transitions driven by both the internal dynamics of species interactions and external forcing, and can uncover unexpected phenomena such as multiple stable annual trajectories and year-to-year irregularity in successional trajectories (chaos). Steiner et al. [10] later validated the utility of the SSD approach as a framework for predicting the effects of altered seasonality on the structure and dynamics of multitrophic communities by using controlled laboratory experiments.

However, there are few analytic results on these ecological models with seasonal succession. One of the major reasons is that the vector fields of these models are discontinuous and periodic in time t . Hsu and Zhao [11] first studied the global dynamics of a Lotka-Volterra two-species competition model with seasonal succession and obtained a complete classification for the global dynamics and the effects of season succession on the competition outcomes via the theory of monotone dynamical systems. Recently, Niu et al. [12] were concerned with an n -dimensional Lotka-Volterra competition model with seasonal succession and obtained the existence of a carrying simplex. Based on this, they reconsidered the two-dimensional model proposed by Hsu and Zhao [11] and simplified their approach to acquire the complete classification of global dynamics. In [13], Xie and Niu analyzed a three-dimensional Lotka-Volterra cooperation model with seasonal succession and derived a completed dynamics of global coexistence and extinction, which extends previous results with respect to three-dimensional cooperation models. There are other works on seasonal succession, such as [14–18] and references therein.

Yet, to our knowledge, there are few works on the global stability for high dimensional competitive systems with seasonal succession due to such obstacles as the estimates for the Floquet multipliers, the existence and uniqueness of the positive periodic solutions, etc. Therefore, it is interesting for us to introduce the seasonal succession into the three-species tridiagonal competition model and study the global stability for such a periodically forced system. Motivated by the modelling methods in Klausmeier [9], we propose a tridiagonal three-species competition model with seasonal succession as

follows:

$$\begin{cases} \frac{dx_i}{dt} = -\lambda_i x_i, & m\omega \leq t \leq m\omega + (1 - \varphi)\omega, \quad i = 1, 2, 3, \\ \frac{dx_1}{dt} = x_1(r_1 - a_{11}x_1 - a_{12}x_2), & m\omega + (1 - \varphi)\omega \leq t \leq (m + 1)\omega, \\ \frac{dx_2}{dt} = x_2(r_2 - a_{21}x_1 - a_{22}x_2 - a_{23}x_3), & m\omega + (1 - \varphi)\omega \leq t \leq (m + 1)\omega, \\ \frac{dx_3}{dt} = x_3(r_3 - a_{32}x_2 - a_{33}x_3), & m\omega + (1 - \varphi)\omega \leq t \leq (m + 1)\omega, \\ (x_1(0), x_2(0), x_3(0)) = x^0 \in \mathbb{R}_+^3, & m = 0, 1, 2, \dots, \end{cases} \quad (1.2)$$

where $m \in \mathbb{Z}_+$, $\varphi \in [0, 1]$ and $\omega, \lambda_i, r_i, a_{ij}$ ($i, j = 1, 2, 3$) are all positive constants.

In particular, if $\varphi = 0$, then the system (1.2) become the linear system

$$\begin{cases} \frac{dx_i}{dt} = -\lambda_i x_i, & i = 1, 2, 3, \\ (x_1(0), x_2(0), x_3(0)) = x^0 \in \mathbb{R}_+^3. \end{cases} \quad (1.3)$$

While, if $\varphi = 1$, system (1.2) turns out to be the tridiagonal competitive system (1.1).

Obviously, system (1.2) is a time periodic system in a season alternate environment. Overall period is ω , and φ stands for the switching proportion of a period between two subsystems (1.1) and (1.3). Biologically, φ is used to describe the proportion of the period in the good season in which the species follow system (1.1), while $(1 - \varphi)$ represents the proportion of the period in the bad season in which the species die exponentially according to system (1.3).

In addition, when we write

$$r_i(t) = \begin{cases} -\lambda_i, & [m\omega, m\omega + (1 - \varphi)\omega), \\ b_i, & [m\omega + (1 - \varphi)\omega, (m + 1)\omega], \end{cases} \quad a_{ij}(t) = \begin{cases} 0, & [m\omega, m\omega + (1 - \varphi)\omega), \\ a_{ij}, & [m\omega + (1 - \varphi)\omega, (m + 1)\omega], \end{cases}$$

where $i, j = 1, 2, 3$, system (1.2) can be expressed as a three-dimensional time ω -periodic tridiagonal competitive system with discontinuous ω -periodic coefficients,

$$\begin{cases} \frac{dx_1(t)}{dt} = x_1(t)(r_1(t) - a_{11}(t)x_1(t) - a_{12}(t)x_2(t)), \\ \frac{dx_2(t)}{dt} = x_2(t)(r_2(t) - a_{21}(t)x_1(t) - a_{22}(t)x_2(t) - a_{23}(t)x_3(t)), \\ \frac{dx_3(t)}{dt} = x_3(t)(r_3(t) - a_{32}(t)x_2(t) - a_{33}(t)x_3(t)), \\ (x_1(0), x_2(0), x_3(0)) = x^0 \in \mathbb{R}_+^3. \end{cases} \quad (1.4)$$

From system (1.2), it can easily be seen that system (1.2) admits a unique nonnegative global solution $x(t, x^0)$ on $[0, +\infty)$ for any $x^0 \in \mathbb{R}_+^3$. Since the system is ω -periodic, it suffices to consider the Poincaré map S on \mathbb{R}_+^3 , that is,

$$S(x^0) = x(\omega, x^0), \quad \forall x^0 \in \mathbb{R}_+^3.$$

Let us first define a linear map L by

$$L(x_1, x_2, x_3) = (e^{-\lambda_1(1-\varphi)\omega} x_1, e^{-\lambda_2(1-\varphi)\omega} x_2, e^{-\lambda_3(1-\varphi)\omega} x_3), \quad \forall (x_1, x_2, x_3) \in \mathbb{R}_+^3. \quad (*)$$

We also let $\{Q_t\}_{t \geq 0}$ represent the solution flow associated with the tridiagonal competitive system (1.1). Then, $Q_t(x^0)$ is the unique global solution of the system (1.1) on $[0, +\infty)$. Thus, we have

$$S(x^0) = Q_{\varphi\omega}(Lx^0), \quad \forall x^0 \in \mathbb{R}_+^3, \quad \text{i.e.,} \quad S = Q_{\varphi\omega} \circ L.$$

We only need to focus on the dynamics of the discrete-time system $\{S^n\}_{n \geq 0}$.

The purpose of this paper is to investigate the global stability for system (1.2). Firstly, the Floquet multipliers of all non-negative periodic solutions, including trivial periodic solution, semi-trivial periodic solutions and the positive periodic solutions, are estimated via the stability analysis of equilibria (see Lemmas 3.1–3.8). To obtain the existence and uniqueness for the positive periodic solution, we provide the index theory of the fixed points for the Poincaré map S (see Lemma 4.1). Compared to the proof of the existence and uniqueness for the positive periodic solution in Xie and Niu [13] and [Lemmas 4.1 and 4.2], our approach is much more general which avoids Brouwer fixed point theorem and the connecting orbits theorem and can be applied to more mappings, such as the the Poincaré maps associated with competition and cooperation models with seasonal succession. Together with the existence of carrying simplex, we obtain that system (1.2) has a unique positive periodic solution under appropriate conditions, and moreover the positive periodic solution is globally asymptotically stable in $\text{Int}\mathbb{R}_+^3$ (see Theorem 4.4). In addition, sufficient conditions for two-species coexistence, two-species extinction and global extinction are given (see Theorems 4.5–4.9). Some numerical examples are provided to illustrate our theoretical results (see Figures 1–5).

The paper is organized as follows. In Section 2, we introduce some notations, relevant definitions and preliminaries. Section 3 is devoted to analyzing the local dynamics of all nonnegative periodic solutions of system (1.2). In Section 4, the global dynamics of coexistence and extinction for three competing species in terms of system (1.2) are obtained. In Section 5, we present some numerical simulations to illustrate our analytic results. The paper ends with a discussion in Section 6.

2. Notations and preliminaries

In this section, we first introduce some definitions and describe some results which are essential tools for the later sections.

Define \mathbb{Z}_+ to be the set of nonnegative integers. Let $\mathbb{R}_+^3 = \{x \in \mathbb{R}^3 : x_i \geq 0, \forall i \in \Lambda\}$ and $\text{Int}\mathbb{R}_+^3 = \{x \in \mathbb{R}^3 : x_i > 0, \forall i \in \Lambda\}$, where $\Lambda := \{1, 2, 3\}$. Let $\emptyset \neq L \subset \Lambda$ and $\bar{L} = \Lambda \setminus L$ be its complementary set in Λ . We define the sets $H_L = \{x \in \mathbb{R}^3 : x_j = 0 \text{ for } j \in \bar{L}\}$, $H_L^+ = \mathbb{R}_+^3 \cap H_L$, and $\text{Int}H_L^+ = \{x \in H_L^+ : x_i > 0 \text{ for all } i \in L\}$. For two vectors $x, y \in \mathbb{R}^3$, we write $x \leq y$ if $x_i \leq y_i$ for all $i \in \Lambda$, and $x \ll y$ if $x_i < y_i$ for all $i \in \Lambda$. If $x \leq y$ but $x \neq y$, we write $x < y$. If $x, y \in \mathbb{R}^3$ and $x \leq y$, let $[x, y] = \{z \in \mathbb{R}^3 : x \leq z \leq y\}$ be a closed order interval, and if $x \ll y$, let $[[x, y]] = \{z \in \mathbb{R}^3 : x \ll z \ll y\}$ be an open order interval. For an $n \times n$ matrix A , we write $A \geq 0$ iff A is a nonnegative matrix (i.e., all the entries are nonnegative) and $A \gg 0$ iff A is a positive matrix (i.e., all the entries are positive).

Let $X \subset \mathbb{R}_+^3$ and $S : X \rightarrow X$ be the Poincaré (period) map. The orbit of a state x for S is $\gamma(x) = \{S^n(x), n \in \mathbb{Z}_+\}$. A fixed point x of S is a point $x \in X$ such that $S(x) = x$. A point $y \in X$ is called a k -periodic point of S if there exists some positive integer $k > 1$, such that $S^k(y) = y$ and $S^m(y) \neq y$ for every positive integer $m < k$. The k -periodic orbit of the k -periodic point y , $\gamma(y) = \{y, S(y), S^2(y), \dots, S^{k-1}(y)\}$, is often called a periodic orbit for short. The ω -limit set of x is defined by $\omega(x) = \{y \in \mathbb{R}_+^3 : S^{n_k}x \rightarrow y (k \rightarrow \infty) \text{ for some sequence } n_k \rightarrow +\infty \text{ in } \mathbb{Z}_+\}$. A set $V \subset X$ is positively

invariant under S if $SV \subset V$, and invariant if $SV = V$. Note that if the orbit $\gamma(x)$ of x has compact closure, $\omega(x)$ is nonempty, compact and invariant. If S is a differentiable map, we write $DS(x)$ as the Jacobian matrix of S at the point x , and $r(DS(x))$ stands for the spectral radius of $DS(x)$.

A continuous map $S : \mathbb{R}_+^3 \rightarrow \mathbb{R}_+^3$ is said to be monotone if $S(x) \leq S(y)$ whenever $x \leq y$ with $x, y \in \mathbb{R}_+^3$; S is called strictly monotone if $x < y$, then $S(x) < S(y)$; strongly monotone if $x < y$, then $S(x) \ll S(y)$.

A *carrying simplex* for the periodic map S is a subset $\Sigma \subset \mathbb{R}_+^n \setminus \{0\}$ with the following properties [26]:

(P1) Σ is compact, invariant and unordered;

(P2) Σ is homeomorphic via radial projection to the $(n - 1)$ -dim standard probability simplex $\Delta^{n-1} := \{x \in \mathbb{R}_+^n \mid \sum_i x_i = 1\}$;

(P3) $\forall x \in \mathbb{R}_+^n \setminus \{0\}$, there exists some $y \in \Sigma$ such that $\lim_{n \rightarrow \infty} |S^n x - S^n y| = 0$.

Lemma 2.1. ([11, Lemma 2.1]) *Let $x(t, x_0)$ be the unique solution of the following system*

$$\begin{cases} \frac{dx}{dt} = -\lambda x, & m\omega \leq t \leq m\omega + (1 - \varphi)\omega, \quad m = 0, 1, 2, \dots, \\ \frac{dx}{dt} = x(r - ax), & m\omega + (1 - \varphi)\omega \leq t \leq (m + 1)\omega, \\ x(0) = x^0 \in \mathbb{R}_+, \end{cases} \quad (2.1)$$

where λ, r, a are all positive constants. Then the following two statements are valid:

(i) If $r\varphi - \lambda(1 - \varphi) \leq 0$, then $\lim_{t \rightarrow \infty} x(t, x_0) = 0$ for all $x_0 \in \mathbb{R}_+$;

(ii) If $r\varphi - \lambda(1 - \varphi) > 0$, then system (2.1) admits a unique positive ω -periodic solution $x^*(t)$, and $\lim_{t \rightarrow \infty} (x(t, x_0) - x^*(t)) = 0$ for all $x_0 \in \mathbb{R}_+ \setminus \{0\}$.

Lemma 2.2. (Boundness) *The Poincaré map S associated with system (1.2) is bounded in \mathbb{R}_+^3 . Moreover, every forward orbit of S is precompact in \mathbb{R}_+^3 .*

Proof. Firstly, we rewrite system (1.1) as

$$\begin{cases} \dot{x}_i(t) = F_i(x), & i = 1, 2, 3, \\ (x_1(0), x_2(0), x_3(0)) = x^0 \in \mathbb{R}_+^3, \end{cases}$$

where

$$\begin{cases} F_1(x) = x_1(r_1 - a_{11}x_1 - a_{12}x_2), \\ F_2(x) = x_2(r_2 - a_{21}x_1 - a_{22}x_2 - a_{23}x_3), \\ F_3(x) = x_3(r_3 - a_{32}x_2 - a_{33}x_3). \end{cases}$$

Let $b = \max\{\frac{r_1}{a_{11}} + 1, \frac{r_2}{a_{22}} + 1, \frac{r_3}{a_{33}} + 1\}$, we have $B := (b, b, b) \gg 0$. For any positive number $l \geq 1$, whenever $x := (x_1, x_2, x_3) \in [0, lB]$ satisfies $x_1 = lb$, then

$$\dot{x}_1(t) = F_1(x) = lb(r_1 - a_{11}lb - a_{12}x_2) \leq lb(r_1 - a_{11}lb) < 0.$$

Similarly, for any $x \in [0, lB]$ satisfies $x_2 = lb$, then

$$\dot{x}_2(t) = F_2(x) = lb(r_2 - a_{22}lb - a_{21}x_1 - a_{23}x_3) \leq lb(r_2 - a_{22}lb) < 0.$$

while, for any $x \in [0, lB]$ satisfies $x_3 = lb$, we also have

$$\dot{x}_3(t) = F_3(x) \leq lb(r_3 - a_{33}lb - a_{32}x_2) \leq lb(r_3 - a_{33}lb) < 0.$$

Thus, $[0, lB]$ is positive invariant for system (1.1).

For any $x^0 \in \mathbb{R}_+^3$, there exists some positive number $l_0 \geq 1$ such that $0 \leq x^0 \leq l_0B$. The positive invariance of $[0, l_0B]$ implies that $0 \leq x(t, x^0) \leq l_0B$ for all $t \geq 0$ where $x(t, x^0) = (x_1(t), x_2(t), x_3(t))$ stands for the solution of system (1.1) with initial value x^0 in \mathbb{R}_+^3 . Note that $\{Q_t\}_{t \geq 0}$ is the solution flow of system (1.1), it follows that $0 \leq Q_t(x^0) \leq l_0B$ for any $t \geq 0$. By the expression (*) of $L(x_1, x_2, x_3)$, we have $Lx^0 \leq x^0$, and then, $0 \leq Q_t(Lx^0) \leq l_0B$ for any $t \geq 0$. This implies that $0 \leq Q_{\varphi\omega}(Lx^0) = S(x^0) \leq l_0B$, that is, S is bounded. Based on this, we easily see that every forward orbit of S is precompact. The proof is completed. \square

Lemma 2.3. (Monotonicity) *If $x, y \in \mathbb{R}_+^3$ and $S(x) < S(y)$, then $x < y$. In particular, if $x, y \in \text{Int}\mathbb{R}_+^3$ and $S(x) < S(y)$, then $x \ll y$.*

Proof. For $x, y \in \mathbb{R}_+^3$, if $S(x) < S(y)$, then $Q_{\varphi\omega}(Lx) < Q_{\varphi\omega}(Ly)$. Define $u(t) := Q_{\varphi\omega-t}(Lx)$ and $v(t) := Q_{\varphi\omega-t}(Ly)$. It is clear that $u(t)$ and $v(t)$ are two solutions of a three dimensional cooperative system for $t \in [0, \varphi\omega]$. Since $u(0) = Q_{\varphi\omega}(Lx) < Q_{\varphi\omega}(Ly) = v(0)$, it follows from the comparison theorem of cooperative systems that $Lx = u(\varphi\omega) < v(\varphi\omega) = Ly$. By the expression of the linear map L , we have $x < y$. For the strong monotonicity of S , see Smith [24, Chapter 3]. \square

Remark 2.1. *By the expression of system (1.4), H_L, H_L^+ and $\text{Int}H_L^+$ are positively invariant under the map S . So we can rewrite the Poincaré map S as:*

$$S(x_1, x_2, x_3) = (x_1G_1(x), x_2G_2(x), x_3G_3(x)), \quad x \in \mathbb{R}_+^3$$

where

$$G_i(x) := \begin{cases} \frac{S_i(x)}{x_i} & \text{if } x_i \neq 0, \\ \frac{\partial S_i}{\partial x_i}(x) & \text{if } x_i = 0. \end{cases}$$

Moreover, $G_i(x)$ are continuous functions with $G_i(x) \geq 0$ for $x \in \mathbb{R}_+^3, i = 1, 2, 3$.

3. Local dynamics

In this section, we will analyze the local stabilities of all nonnegative fixed points of S in \mathbb{R}_+^3 . Clearly, $O := (0, 0, 0)$ is a trivial fixed point of S . Let $h_i := (r_i\varphi - \lambda_i(1 - \varphi))\omega, i = 1, 2, 3$. By Lemma 2.1, if $h_i > 0$ ($i = 1, 2, 3$), then system (1.2) has three axial fixed points, say $R_1 := (x_1^*, 0, 0)$, $R_2 := (0, x_2^*, 0)$ and $R_3 := (0, 0, x_3^*)$. By the second and fourth equations of system (1.2), it is not difficult to see that there admits a unique interior fixed point in coordinate planar $\{x_2 = 0\}$, say $E_2 = (\hat{x}_1, 0, \hat{x}_3)$. Noticing that Hsu and Zhao [11, Theorems 2.3 and 2.4], system (1.2) has interior fixed points in coordinate planar $\{x_1 = 0\}$ and $\{x_3 = 0\}$ under certain conditions. If exist, we write them as $E_3 = (\bar{x}_1, \bar{x}_2, 0)$ and $E_1 = (0, \check{x}_2, \check{x}_3)$, respectively.

Lemma 3.1. (Stability of the fixed point O)

(i) *If $h_i < 0$ ($i = 1, 2, 3$), then O is an asymptotically stable fixed point of S .*

(ii) If one of h_1, h_2 and h_3 is greater than 0, then O is an unstable fixed point of S . In particular, if $h_i > 0$ ($i = 1, 2, 3$), then O is a hyperbolic repeller.

Proof. Let $F(x) = (F_1, F_2, F_3)^T$, where

$$\begin{cases} F_1 = r_1 x_1 - a_{11} x_1^2 - a_{12} x_1 x_2, \\ F_2 = r_2 x_2 - a_{21} x_1 x_2 - a_{22} x_2^2 - a_{23} x_2 x_3, \\ F_3 = r_3 x_3 - a_{32} x_2 x_3 - a_{33} x_3^2. \end{cases}$$

Then, $DF(x) =$

$$\begin{pmatrix} r_1 - 2a_{11}x_1 - a_{12}x_2 & -a_{12}x_1 & 0 \\ -a_{21}x_2 & r_2 - a_{21}x_1 - 2a_{22}x_2 - a_{23}x_3 & -a_{23}x_2 \\ 0 & -a_{32}x_3 & r_3 - a_{32}x_2 - 2a_{33}x_3 \end{pmatrix}.$$

For simplicity, we denote $u(t, x) := Q_t(x)$ and $V(t, x) := D_x u(t, x) = D_x Q_t(x)$. Then $S(x) = Q_{\varphi\omega}(Lx) = u(\varphi\omega, Lx)$. Thus,

$$DS(x) = D(Q_{\varphi\omega}(Lx)) \cdot D(Lx) = V(\varphi\omega, Lx) \cdot D(Lx) \quad (3.1)$$

$$= V(\varphi\omega, Lx) \cdot \text{diag}\{e^{-\lambda_1(1-\varphi)\omega}, e^{-\lambda_2(1-\varphi)\omega}, e^{-\lambda_3(1-\varphi)\omega}\}. \quad (3.2)$$

Note that $V(t, x)$ satisfies

$$\frac{dV(t)}{dt} = DF(u(t, x))V(t), \quad V(0) = I. \quad (3.3)$$

Taking $x = O$, we have $u(t, LO) = (0, 0, 0)$, which implies that $DF(O) = \text{diag}\{r_1, r_2, r_3\}$. Then,

$$V(\varphi\omega, LO) = \text{diag}\{e^{\int_0^{\varphi\omega} r_1 dt}, e^{\int_0^{\varphi\omega} r_2 dt}, e^{\int_0^{\varphi\omega} r_3 dt}\},$$

and hence, one has

$$DS(O) = \text{diag}\{e^{(r_1\varphi - \lambda_1(1-\varphi))\omega}, e^{(r_2\varphi - \lambda_2(1-\varphi))\omega}, e^{(r_3\varphi - \lambda_3(1-\varphi))\omega}\} = \text{diag}\{e^{h_1}, e^{h_2}, e^{h_3}\}.$$

Consequently, the matrix $DS(O)$ has three positive eigenvalues μ_1, μ_2 and μ_3 given by

$$\mu_1 = e^{h_1}, \quad \mu_2 = e^{h_2} \quad \text{and} \quad \mu_3 = e^{h_3}.$$

Then, the conclusion is immediate. □

By Lemma 3.1, we assume that $h_i > 0$ ($i = 1, 2, 3$) in following Lemmas 3.2–3.8.

Lemma 3.2. (Stability of the axial fixed point R_1)

(i) If $\frac{h_2}{a_{21}} > \frac{h_1}{a_{11}}$, then R_1 is a saddle point with one-dimensional stable manifold.

(ii) If $\frac{h_2}{a_{21}} < \frac{h_1}{a_{11}}$, then R_1 is a saddle point with two-dimensional stable manifold.

Proof. Let $u(t, LR_1) := (u_1^*(t), 0, 0)$. By the proof of Lemma 3.1, it is not difficult to see that

$$DF(u(t, LR_1)) = \begin{pmatrix} r_1 - 2a_{11}u_1^*(t) & -a_{12}u_1^*(t) & 0 \\ 0 & r_2 - a_{21}u_1^*(t) & 0 \\ 0 & 0 & r_3 \end{pmatrix}.$$

Then,

$$V(\varphi\omega, LR_1) = \begin{pmatrix} e^{\int_0^{\varphi\omega} r_1 - 2a_{11}u_1^*(t)dt} & * & * \\ 0 & e^{\int_0^{\varphi\omega} r_2 - a_{21}u_1^*(t)dt} & * \\ 0 & 0 & e^{\int_0^{\varphi\omega} r_3 dt} \end{pmatrix}.$$

Note that $u_1^*(t)$ satisfies the equation $\frac{du_1^*(t)}{dt} = u_1^*(t)(r_1 - a_{11}u_1^*(t))$, it follows that

$$\int_0^{\varphi\omega} u_1^*(t)dt = \frac{h_1}{a_{11}},$$

and then,

$$DS(R_1) = \begin{pmatrix} e^{-h_1} & * & * \\ 0 & e^{(h_2 - \frac{a_{21}}{a_{11}})h_1} & * \\ 0 & 0 & e^{h_3} \end{pmatrix},$$

where * stands for unknown algebraic expression. Therefore, the matrix $DS(R_1)$ has three positive eigenvalues μ_1, μ_2 and μ_3 given by

$$\mu_1 = e^{-h_1} < 1, \quad \mu_2 = e^{(h_2 - \frac{a_{21}}{a_{11}})h_1} \quad \text{and} \quad \mu_3 = e^{h_3} > 1.$$

Therefore, the conclusion is immediate. \square

Similarly, we have the Lemmas 3.3 and 3.4.

Lemma 3.3. (Stability of the axial fixed point R_2)

- (i) If $\frac{h_2}{a_{22}} > \max\{\frac{h_1}{a_{12}}, \frac{h_3}{a_{32}}\}$, then R_2 is an asymptotically stable fixed point of S .
- (ii) If $\frac{h_2}{a_{22}} < \min\{\frac{h_1}{a_{12}}, \frac{h_3}{a_{32}}\}$, then R_2 is a saddle point with one-dimensional stable manifold.
- (iii) If $\frac{h_1}{a_{12}} < \frac{h_2}{a_{22}} < \frac{h_3}{a_{32}}$ or $\frac{h_1}{a_{12}} > \frac{h_2}{a_{22}} > \frac{h_3}{a_{32}}$, then R_2 is a saddle point with two-dimensional stable manifold.

Lemma 3.4. (Stability of the axial fixed point R_3)

- (i) If $\frac{h_2}{a_{23}} > \frac{h_3}{a_{33}}$, then R_3 is a saddle point with one-dimensional stable manifold.
- (ii) If $\frac{h_2}{a_{23}} < \frac{h_3}{a_{33}}$, then R_3 is a saddle point with two-dimensional stable manifold.

Lemma 3.5. (Stability of the planar fixed point E_3) If $\frac{h_2}{a_{21}} > (<) \frac{h_1}{a_{11}}, \frac{h_1}{a_{12}} > (<) \frac{h_2}{a_{22}}$, then there admits a unique interior fixed point, say $E_3 := (\bar{x}_1, \bar{x}_2, 0)$, for S in the coordinate plane $\{x_3 = 0\}$. Moreover,

- (i) If $\frac{h_2}{a_{21}} > \frac{h_1}{a_{11}}, \frac{h_1}{a_{12}} > \frac{h_2}{a_{22}}$ and $\frac{h_3}{a_{32}} < \frac{h_1 a_{21} - h_2 a_{11}}{a_{12} a_{21} - a_{11} a_{22}}$, then E_3 is an asymptotically stable fixed point of S .

(ii) If $\frac{h_2}{a_{21}} < \frac{h_1}{a_{11}}$, $\frac{h_1}{a_{12}} < \frac{h_2}{a_{22}}$ and $\frac{h_3}{a_{32}} < \frac{h_1 a_{21} - h_2 a_{11}}{a_{12} a_{21} - a_{11} a_{22}}$, then E_3 is a saddle point with two-dimensional stable manifold.

(iii) If $\frac{h_2}{a_{21}} < \frac{h_1}{a_{11}}$, $\frac{h_1}{a_{12}} < \frac{h_2}{a_{22}}$ and $\frac{h_3}{a_{32}} > \frac{h_1 a_{21} - h_2 a_{11}}{a_{12} a_{21} - a_{11} a_{22}}$, then E_3 is a saddle point with one-dimensional stable manifold.

Proof. By Remark 2.1, we can see that all coordinate axes and planes are positively invariant under the map S . Theorem 3.6 in [12] implies that if $\frac{h_2}{a_{21}} > (<) \frac{h_1}{a_{11}}$, $\frac{h_1}{a_{12}} > (<) \frac{h_2}{a_{22}}$, then system (1.2) admits a unique interior fixed point in the coordinate planar $\{x_3 = 0\}$, say $E_3 := (\bar{x}_1, \bar{x}_2, 0)$. Let $F(x)$, $u(t, x)$ and $V(t, x)$ be denoted as in the proof of Lemma 3.1. We also define $u(t, LE_3) := (\bar{u}_1(t), \bar{u}_2(t), 0)$. Then,

$$DF(u(t, LE_3)) = \begin{pmatrix} r_1 - 2a_{11}\bar{u}_1 - a_{12}\bar{u}_2 & -a_{12}\bar{u}_1 & 0 \\ -a_{21} & r_2 - 2a_{22}\bar{u}_2 - a_{21}\bar{u}_1 & -a_{23}\bar{u}_2 \\ 0 & 0 & r_3 - a_{32}\bar{u}_2 \end{pmatrix}.$$

For convenience, we write $DF(u(t, LE_3)) := \begin{pmatrix} A_1 & B_1 \\ 0 & r_3 - a_{32}\bar{u}_2(t) \end{pmatrix}$, where

$$B_1 = \begin{pmatrix} 0 \\ -a_{23}\bar{u}_2(t) \end{pmatrix} \quad \text{and} \quad A_1 = \begin{pmatrix} r_1 - 2a_{11}\bar{u}_1(t) - a_{12}\bar{u}_2(t) & -a_{12}\bar{u}_1(t) \\ -a_{21}\bar{u}_2(t) & r_2 - 2a_{22}\bar{u}_2(t) - a_{21}\bar{u}_1(t) \end{pmatrix}.$$

Since $DS(x) = V(\varphi\omega, Lx) \cdot D(Lx)$, it follows that $DS(E_3)$ has a positive eigenvalue μ_3 given by

$$\mu_3 = \exp\left(\int_0^{\varphi\omega} (r_3 - a_{32}\bar{u}_2(t)) dt - \lambda_3(1 - \varphi)\omega\right).$$

Note that $\bar{u}_1(t)$ and $\bar{u}_2(t)$ satisfy the following equations

$$\begin{cases} \frac{d\bar{u}_1(t)}{dt} = \bar{u}_1(t)(b_1 - a_{11}\bar{u}_1(t) - a_{12}\bar{u}_2(t)), \\ \frac{d\bar{u}_2(t)}{dt} = \bar{u}_2(t)(b_2 - a_{22}\bar{u}_2(t) - a_{21}\bar{u}_1(t)). \end{cases} \quad (3.4)$$

Integrating the equation (3.4) for t from 0 to $\varphi\omega$, one has

$$\int_0^{\varphi\omega} \bar{u}_1(t) dt = \frac{h_2 a_{12} - h_1 a_{22}}{a_{12} a_{21} - a_{11} a_{22}} \quad \text{and} \quad \int_0^{\varphi\omega} \bar{u}_2(t) dt = \frac{h_1 a_{21} - h_2 a_{11}}{a_{12} a_{21} - a_{11} a_{22}}.$$

Then,

$$\mu_3 = \exp\left(h_3 - \frac{(h_1 a_{21} - h_2 a_{11}) a_{32}}{a_{12} a_{21} - a_{11} a_{22}}\right).$$

On the other hand, the other two eigenvalues μ_1 and μ_2 are determined by A_1 . Together with Lemma 2.4 in [11], the conclusion is immediate. \square

Similarly, we also have the following Lemmas 3.6 and 3.7.

Lemma 3.6. (Stability of the planar fixed point E_2) *There admits a unique interior fixed point, say $E_2 := (\hat{x}_1, 0, \hat{x}_3)$, for S in the coordinate plane $\{x_2 = 0\}$. Moreover,*

(i) If $h_2 < \frac{h_1 a_{21}}{a_{11}} + \frac{h_3 a_{23}}{a_{33}}$, then E_2 is an asymptotically stable fixed point of S .

(ii) If $h_2 > \frac{h_1 a_{21}}{a_{11}} + \frac{h_3 a_{23}}{a_{33}}$, then E_2 is a saddle point with two-dimensional stable manifold.

Lemma 3.7. (Stability of the planar fixed point E_1) If $\frac{h_2}{a_{23}} > (<) \frac{h_3}{a_{33}}$, $\frac{h_3}{a_{32}} > (<) \frac{h_2}{a_{22}}$, then there admits a unique interior fixed point, say $E_1 := (0, \check{x}_2, \check{x}_3)$, for S in the coordinate plane $\{x_1 = 0\}$. Moreover,

- (i) If $\frac{h_2}{a_{23}} > \frac{h_3}{a_{33}}$, $\frac{h_3}{a_{32}} > \frac{h_2}{a_{22}}$ and $\frac{h_1}{a_{12}} < \frac{a_{23}h_3 - a_{33}h_2}{a_{23}a_{32} - a_{22}a_{33}}$, then E_1 is an asymptotically stable fixed point of S .
- (ii) If $\frac{h_2}{a_{23}} < \frac{h_3}{a_{33}}$, $\frac{h_3}{a_{32}} < \frac{h_2}{a_{22}}$ and $\frac{h_1}{a_{12}} < \frac{a_{23}h_3 - a_{33}h_2}{a_{23}a_{32} - a_{22}a_{33}}$, then E_1 is a saddle point with two-dimensional stable manifold.
- (iii) If $\frac{h_2}{a_{23}} < \frac{h_3}{a_{33}}$, $\frac{h_3}{a_{32}} < \frac{h_2}{a_{22}}$ and $\frac{h_1}{a_{12}} > \frac{a_{23}h_3 - a_{33}h_2}{a_{23}a_{32} - a_{22}a_{33}}$, then E_1 is a saddle point with one-dimensional stable manifold.

Assume that S has a positive fixed point, say $P := (\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$, we will discuss the dynamics of P in the following. For simplicity, we write

$$\bar{A} := \begin{pmatrix} -a_{11} & a_{12} & 0 \\ a_{21} & -a_{22} & a_{23} \\ 0 & a_{32} & -a_{33} \end{pmatrix}.$$

Lemma 3.8. (Stability of the positive fixed point P)

- (i) If $\det \bar{A} < 0$, then P is an asymptotically stable fixed point of S .
- (ii) If $\det \bar{A} > 0$, then P is an unstable fixed point of S .

Proof. By the expression of L , it follows that

$$LP = (e^{-\lambda_1(1-\varphi)\omega} \tilde{x}_1, e^{-\lambda_2(1-\varphi)\omega} \tilde{x}_2, e^{-\lambda_3(1-\varphi)\omega} \tilde{x}_3).$$

Define $Q_t(LP) := (\tilde{u}_1(t), \tilde{u}_2(t), \tilde{u}_3(t)) = \tilde{u}(t)$, and $V(t, x) := D_x \tilde{u}(t)$. Note that

$DF(u(t), LP) =$

$$\begin{pmatrix} r_1 - 2a_{11}\tilde{u}_1 - a_{12}\tilde{u}_2 & -a_{12}\tilde{u}_1 & 0 \\ -a_{21}\tilde{u}_2 & r_2 - 2a_{22}\tilde{u}_2 - a_{21}\tilde{u}_1 - a_{23}\tilde{u}_3 & -a_{23}\tilde{u}_2 \\ 0 & -a_{32}\tilde{u}_3 & r_3 - 2a_{33}\tilde{u}_3 - a_{32}\tilde{u}_2 \end{pmatrix},$$

the matrix function $V(t) = V(t, LP)$ satisfies

$$\frac{dV(t)}{dt} = DF(\tilde{u}(t))V(t), \quad V(0) = I.$$

Let $w(t) = P(t)V(t)$, where $P(t) = \text{diag}\{\frac{1}{\tilde{u}_1(t)}, \frac{1}{\tilde{u}_2(t)}, \frac{1}{\tilde{u}_3(t)}\}$. Then, one has

$$\frac{dw(t)}{dt} = A(t)w(t) \tag{3.5}$$

where

$$A(t) = \begin{pmatrix} -a_{11}\tilde{u}_1(t) & -a_{12}\tilde{u}_2(t) & 0 \\ -a_{21}\tilde{u}_1(t) & -a_{22}\tilde{u}_2(t) & -a_{23}\tilde{u}_3(t) \\ 0 & -a_{32}\tilde{u}_2(t) & -a_{33}\tilde{u}_3(t) \end{pmatrix}.$$

Let $W(t)$ be the monodromy matrix of the above Eq (3.5), then $W(t)$ satisfies

$$\frac{dW(t)}{dt} = A(t)W(t), \quad W(0) = I,$$

and hence $W(t) = P(t)V(t)P^{-1}(0)$. Thus $V(t) = P^{-1}(t)W(t)P(0)$. In view of

$$P(0) = \text{diag}\left\{\frac{e^{\lambda_1(1-\varphi)\omega}}{\tilde{x}_1}, \frac{e^{\lambda_2(1-\varphi)\omega}}{\tilde{x}_2}, \frac{e^{\lambda_3(1-\varphi)\omega}}{\tilde{x}_3}\right\}$$

and

$$P(\varphi\omega) = \text{diag}\left\{\frac{1}{\tilde{u}_1(\varphi\omega)}, \frac{1}{\tilde{u}_2(\varphi\omega)}, \frac{1}{\tilde{u}_3(\varphi\omega)}\right\} = \text{diag}\left\{\frac{1}{\tilde{x}_1}, \frac{1}{\tilde{x}_2}, \frac{1}{\tilde{x}_3}\right\},$$

we have

$$V(\varphi\omega) = P^{-1}(\varphi\omega)W(\varphi\omega)P(0).$$

Then,

$$\begin{aligned} DS(P) &= V(\varphi\omega, LP) \cdot D(LP) \\ &= P^{-1}(\varphi\omega)W(\varphi\omega)P(0) \cdot D(LP) \\ &= \text{diag}\{\tilde{x}_1, \tilde{x}_2, \tilde{x}_3\} \cdot W(\varphi\omega) \cdot \text{diag}\left\{\frac{1}{\tilde{x}_1}, \frac{1}{\tilde{x}_2}, \frac{1}{\tilde{x}_3}\right\}, \end{aligned}$$

and hence, $DS(P) \sim W(\varphi\omega)$, that is, $r(DS(P)) = r(W(\varphi\omega))$.

Let

$$Z(t) = \text{diag}\{-1, 1, -1\} \cdot W(t) \cdot \text{diag}\{-1, 1, -1\}^{-1},$$

then

$$\frac{dZ(t)}{dt} = \tilde{A}(t)Z(t), \quad Z(0) = I, \quad (3.6)$$

where

$$\tilde{A}(t) = \begin{pmatrix} -a_{11}\tilde{u}_1(t) & a_{12}\tilde{u}_2(t) & 0 \\ a_{21}\tilde{u}_1(t) & -a_{22}\tilde{u}_2(t) & a_{23}\tilde{u}_3(t) \\ 0 & a_{32}\tilde{u}_2(t) & -a_{33}\tilde{u}_3(t) \end{pmatrix}.$$

Note that $Z(\varphi\omega) \sim W(\varphi\omega)$ implies that $DS(P) \sim Z(\varphi\omega)$, then,

$$r(DS(P)) = r(Z(\varphi\omega)).$$

Since the matrix $\tilde{A}(t)$ is cooperative and irreducible, it follows that $Z(t) > Z(0)$ for each $t > 0$, and then $Z(t)$ is a positive matrix (see [25, Theorem B.3]). By Perron-Frobenius theorem, $\rho_3 := r(Z(\varphi\omega))$ is a simple eigenvalue of $Z(\varphi\omega)$ with a positive eigenvector $e = (e_1, e_2, e_3)^T$. If ρ_1 and ρ_2 represent the other two eigenvalues of $Z(\varphi\omega)$, then $|\rho_i| < \rho_3$, $i = 1, 2$. By Liouville's formula, we also obtain that $0 < \rho_1\rho_2\rho_3 = \det Z(\varphi\omega) = e^{\int_0^{\varphi\omega} \text{trace}(\tilde{A}(s))ds} < 1$.

Let $z(t) := Z(t)e = (z_1(t), z_2(t), z_3(t))$, then $z(\varphi\omega) = Z(\varphi\omega)e = \rho_3(e_1, e_2, e_3)$ and $z(0) = Z(0)e = (e_1, e_2, e_3) > 0$. Noticing that $z(t)$ satisfies the equation

$$\frac{dz(t)}{dt} = \tilde{A}(t)z(t),$$

it follows that

$$\begin{cases} \dot{z}_1(t) = -a_{11}\tilde{u}_1(t)z_1(t) + a_{12}\tilde{u}_2(t)z_2(t), \\ \dot{z}_2(t) = a_{21}\tilde{u}_1(t)z_1(t) - a_{22}\tilde{u}_2(t)z_2(t) + a_{23}\tilde{u}_3(t)z_3(t), \\ \dot{z}_3(t) = a_{32}\tilde{u}_2(t)z_2(t) - a_{33}\tilde{u}_3(t)z_3(t). \end{cases}$$

Using the method of elimination, we have

$$a_{21}a_{33}\dot{z}_1(t) + a_{11}a_{33}\dot{z}_2(t) + a_{23}a_{11}\dot{z}_3(t) = (a_{33}(a_{12}a_{21} - a_{11}a_{22}) + a_{11}a_{23}a_{32})\tilde{u}_2(t)z_2(t).$$

Integrating the above equation for t from 0 to $\varphi\omega$, we then obtain

$$\begin{aligned} a_{21}a_{33} \int_0^{\varphi\omega} dz_1(t) + a_{11}a_{33} \int_0^{\varphi\omega} dz_2(t) + a_{23}a_{11} \int_0^{\varphi\omega} dz_3(t) \\ = (a_{33}(a_{12}a_{21} - a_{11}a_{22}) + a_{11}a_{23}a_{32}) \int_0^{\varphi\omega} \tilde{u}_2(t)z_2(t)dt. \end{aligned}$$

Note that

$$a_{33}(a_{12}a_{21} - a_{11}a_{22}) + a_{11}a_{23}a_{32} = \det \bar{A},$$

it follows that

$$(a_{21}a_{33}e_1 + a_{11}a_{33}e_2 + a_{11}a_{23}e_3) \times (\rho_3 - 1) = \det \bar{A} \cdot \int_0^{\varphi\omega} \tilde{u}_2(t)z_2(t)dt.$$

Based on the fact that $\int_0^{\varphi\omega} \tilde{u}_2(t)z_2(t)dt > 0$, we have

- (i) If $\det \bar{A} > 0$, then $\rho_3 > 1$.
- (ii) If $\det \bar{A} < 0$, then $\rho_3 < 1$.

This implies that the proof is completed. □

4. Global stability

To obtain the existence and uniqueness of the positive fixed point for the Poincaré map S , we first give a lemma with respect to the index of fixed points. For the reader's convenience, we recall some known results on the fixed point index of a continuous map (see Amann [19] for a more detailed discussion).

Let $U \subseteq \mathbb{R}_+^n$ be open and $S : U \rightarrow \mathbb{R}_+^n$ be a continuous map such that $\text{Fix}(S, U)$ is compact, where $\text{Fix}(S, U)$ is defined by the set of all fixed points of S in the subset U . The fixed point index of S is denoted by

$$i(S, U, \mathbb{R}_+^n) := \deg(id - S, U, 0),$$

where id is the identity map and $\deg(id - S, U, 0)$ is the Brouwer degree for $id - S$. The fixed point index of S at an isolated fixed point $\theta \in U$ is defined by

$$i(S, \theta) := i(S, B_\delta(\theta), \mathbb{R}_+^n) = \deg(id - S, B_\delta(\theta), 0),$$

where $B_\delta(\theta) := \{x \in \mathbb{R}^n : \|x - \theta\| < \delta\}$ is an open ball in U such that $\text{Fix}(S, B_\delta(\theta)) = \{\theta\}$. In particular, if S is differentiable at $\theta \in \text{Fix}(S, U)$ and 1 is not an eigenvalue of $DS(\theta)$, then

$$i(S, \theta) = (-1)^\beta,$$

where β is the sum of the multiplicities of all the eigenvalues of $DS(\theta)$ which are greater than one. When S has only finitely many fixed points in U , one has

$$i(S, U, \mathbb{R}_+^n) = \sum_{\theta \in \text{Fix}(S, U)} i(S, \theta).$$

Lemma 4.1. *Assume that each fixed point θ of the Poincaré map S deduced by system (1.2) is hyperbolic, then*

- (i) $i(S, \mathbb{R}_+^3, \mathbb{R}_+^3) = 1$;
- (ii) *If $DS(\theta)$ has no eigenvalue whose modulus is larger than 1, then $i(S, \theta) = 1$;*
- (iii) *If $\theta \in \text{Int}H_L^+$, $L \neq \Lambda = \{1, 2, 3\}$, then $DS(\theta)$ has at least one eigenvalue with modulus larger than 1 $\Leftrightarrow i(S, \theta) = 0$.*

Proof. (i) Since \mathbb{R}_+^3 is nonempty closed convex set, it follows from Dugundjis theorem that $i(S, \mathbb{R}_+^3, \mathbb{R}_+^3)$ is well-defined. Let $x_0 \in \mathbb{R}_+^3$ be arbitrary and define a compact map

$$g : [0, 1] \times \mathbb{R}_+^3 \rightarrow \mathbb{R}_+^3 \text{ by } g(\lambda, x) := (1 - \lambda)x_0 + \lambda S(x).$$

Then g maps the product space into \mathbb{R}_+^3 and has no fixed points on the boundary of \mathbb{R}_+^3 (relative to \mathbb{R}_+^3) because this boundary is empty. By the homotopy invariance and the normalization property, we have

$$i(S, \mathbb{R}_+^3, \mathbb{R}_+^3) = i(x_0, \mathbb{R}_+^3, \mathbb{R}_+^3) = 1.$$

- (ii), (iii) See the proof of Lemma 4.2 for $n = 3$ and $U = \mathbb{R}_+^3$ in Liang and Jiang [20]. □

Arguing as the proof of Theorem 2.3 in Niu et al. [12], it can easily be proved that the Poincaré map S also has a carrying simplex.

Lemma 4.2. (The existence of the carrying simplex) *Assume that $h_i > 0, i = 1, 2, 3$, then the Poincaré map S admits a carrying simplex Σ which attracts every nontrivial orbit in \mathbb{R}_+^3 .*

Proof. See the proof of Theorem 2.3 for $n = 3$ in Niu et al. [12]. □

For each coordinate plane, we write $\Pi_i = \{x \in \mathbb{R}_+^3 : x_i = 0\}, i = 1, 2, 3$. We denote by $S|_{\Pi_i}$ the restriction of S to Π_i . To obtain the global stability of the positive fixed point, we also need the following lemma which is a special version of Theorem 2.4 in Balreira et al. [21] for maps defined on \mathbb{R}_+^3 .

Lemma 4.3. ([21, Theorem 2.4]) *Consider the map $S = (x_1 G_1(x), x_2 G_2(x), x_3 G_3(x))$ defined on \mathbb{R}_+^3 with $G_i(x) \geq 0, i = 1, 2, 3$, which has a carrying simplex. Assume that*

- (a) $\det DS(x) > 0$ for all $x \in \mathbb{R}_+^3$;
- (b) $DS(x)^{-1} > 0$ for all $x \in \text{Int}\mathbb{R}_+^3$;

- (c) for each $i = 1, 2, 3$, $S|_{\Pi_i}$ has a unique interior fixed point $E_{(i)}$ that is globally asymptotically stable in $\text{Int}\Pi_i$, but a saddle for S ;
 (d) S has a unique positive fixed point $p \in \text{Int}\mathbb{R}_+^3$.

Then P is globally asymptotically stable in $\text{Int}\mathbb{R}_+^3$ for S .

Next, we will analyze the global dynamics of system (1.2). Applying Lemma 4.1 to Lemma 3.1–3.7, it is not difficult to calculate the index of each of axial and planar fixed points. If S has a positive fixed point, it follows from Lemma 3.8 that $i(S, P) = 1$ under the condition $\det \bar{A} < 0$. By using the index of each boundary fixed point and $i(S, \mathbb{R}_+^3, \mathbb{R}_+^3) = 1$, we can choose appropriate parameter values to ascertain whether the positive fixed point exists. We can further obtain the global dynamics for system (1.2).

Theorem 4.4. (Three species Coexistence) *Suppose that system (1.2) satisfies*

- (i) $h_i > 0$ ($i = 1, 2, 3$), $\det \bar{A} < 0$;
 (ii) $\frac{h_1}{a_{12}} > \frac{h_2}{a_{22}}$; $\frac{h_3}{a_{32}} > \frac{h_2}{a_{22}}$; $h_2 > \frac{a_{21}}{a_{11}}h_1 + \frac{a_{23}}{a_{33}}h_3$;
 (iii) $h_1(a_{22}a_{33} - a_{23}a_{32}) > a_{12}(a_{33}h_2 - a_{23}h_3)$;
 (iv) $h_3(a_{11}a_{22} - a_{12}a_{21}) > a_{32}(a_{11}h_2 - a_{21}h_1)$;

then there admits a unique positive fixed point, say $P = (\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$, for the Poincaré map S . Moreover, P is globally asymptotically stable in $\text{Int}\mathbb{R}_+^3$.

Proof. By Lemma 2.1, $S|_{H_i^+}$ has a unique interior fixed point R_i that is globally asymptotically stable in $\text{Int}H_i^+$ for each $i = 1, 2, 3$ because $h_i > 0$. And yet, R_i is a saddle point for S due to the assumption (ii) and Lemmas 3.2–3.4. Under the assumptions (ii)–(iv), by Lemmas 3.5–3.7 and Theorem 3.6 in [12], $S|_{\Pi_i}$ has a unique interior fixed point $E_{(i)}$ for $i = 1, 2, 3$ that is globally asymptotically stable in $\text{Int}\Pi_i$, but a saddle point for S . Using Lemma 4.1(iii), one has

$$i(S, O) = i(S, R_i) = i(S, E_i) = 0, \quad i = 1, 2, 3.$$

Since $i(S, \mathbb{R}_+^3, \mathbb{R}_+^3) = 1$ (Lemma 4.1(i)), it is easy to see that there exists at least a positive fixed point for S , say P . Note that $\det \bar{A} < 0$ and Lemma 3.8(i), so P is an asymptotically stable fixed point for S . Lemma 4.1(ii) implies that $i(S, P) = 1$. If there also exists another positive fixed point, say P^* , that is, $P^* \neq P$, then P^* is also asymptotically stable, and hence $i(S, P^*) = 1$, which contradicts the fact that $i(S, \mathbb{R}_+^3, \mathbb{R}_+^3) = 1$. Consequently, the positive fixed point P is unique. By the expression (3.1) of $DS(x)$ and the Eq (3.3), we can obtain that for any $x \in \mathbb{R}_+^3$,

$$\begin{aligned} \det DS(x) &= \det V(\varphi\omega, Lx) \cdot \exp\left(-(\lambda_1 + \lambda_2 + \lambda_3)(1 - \varphi)\omega\right) \\ &= \exp\left\{\int_0^{\varphi\omega} \text{trace}\{DF(u(t, Lx))\}dt\right\} \cdot \exp\left(-(\lambda_1 + \lambda_2 + \lambda_3)(1 - \varphi)\omega\right) > 0. \end{aligned}$$

Besides, it follows from the proof of Theorem 2.3 in [12] that $DS(x)^{-1} > 0$, $\forall x \in \text{Int}\mathbb{R}_+^3$. So far, the conditions (a),(b),(c) and (d) of Lemma 4.3 are satisfied for S , which implies that P is globally asymptotically stable in $\text{Int}\mathbb{R}_+^3$. We have completed the proof. \square

Theorem 4.5. (Two species Coexistence) *Suppose that system (1.2) satisfies*

$$(i) \ h_i > 0 \ (i = 1, 2, 3), \ \det \bar{A} < 0;$$

$$(ii) \ \frac{h_2}{a_{21}} < \frac{h_1}{a_{11}}; \ \frac{h_2}{a_{23}} < \frac{h_3}{a_{33}}; \ \frac{h_1}{a_{12}} > \frac{h_2}{a_{22}}; \ \frac{h_3}{a_{32}} > \frac{h_2}{a_{22}}; \ h_2 < \frac{a_{21}}{a_{11}}h_1 + \frac{a_{23}}{a_{33}}h_3;$$

then there is no positive fixed point for the Poincaré map S . Moreover, the planar fixed point E_2 is globally asymptotically stable in $\text{Int}\mathbb{R}_+^3$.

Proof. By Lemma 2.1 and Theorem 3.6 in [12], there exist three axial fixed points R_i ($i = 1, 2, 3$) and one planar fixed point E_2 under the assumptions (i) and (ii). Moreover, R_i is a saddle point for S and E_2 is an asymptotically stable fixed point of S . In view of Lemma 4.1(ii) and (iii), one has

$$i(S, O) = i(S, R_i) = 0, \ i = 1, 2, 3 \ \text{and} \ i(S, E_2) = 1.$$

If there exists a positive fixed point for S , say P , it follows from $\det \bar{A} < 0$ and Lemma 3.8(i) that P is an asymptotically stable fixed point of S . Lemma 4.1(ii) implies that $i(S, P) = 1$. This contradicts the fact that $i(S, \mathbb{R}_+^3, \mathbb{R}_+^3) = 1$. So there is no positive fixed point for S . Lemma 4.2 states that the Poincaré map S has a carrying simplex Σ by which is homeomorphic to the probability simplex Δ^2 . We further get that Σ is a topological disk and $S|_\Sigma$ is an orientation preserving homeomorphism from Σ onto Σ (see [22]). By Corollary 2.1 in [23], every trajectory on Σ converges to some fixed point. In particular, R_1 and R_3 are saddle points, R_2 is an unstable node point, while E_2 is an asymptotically stable node point on Σ . Since $S|_\Sigma$ is an orientation preserving homeomorphism, every trajectory on Σ converges to E_2 . By using the property (P_3) of the carrying simplex, one has

$$\lim_{n \rightarrow +\infty} S^n(x) = E_2, \quad \forall x \in \text{Int}\mathbb{R}_+^3.$$

Thus, E_2 is globally asymptotically stable in $\text{Int}\mathbb{R}_+^3$. The proof is completed. \square

By suitable modification to the proof of Theorem 4.5, we can also obtain the following Theorems 4.6 and 4.7.

Theorem 4.6. (Two species Coexistence) *Suppose that system (1.2) satisfies*

$$(i) \ h_i > 0 \ (i = 1, 2, 3), \ \det \bar{A} < 0;$$

$$(ii) \ \frac{h_3}{a_{32}} < \frac{h_2}{a_{22}}; \ \frac{h_1}{a_{12}} > \frac{h_2}{a_{22}}; \ h_2 > \frac{a_{21}}{a_{11}}h_1 + \frac{a_{23}}{a_{33}}h_3;$$

$$(iii) \ h_3(a_{11}a_{22} - a_{12}a_{21}) < a_{32}(a_{11}h_2 - a_{21}h_1);$$

then there is no positive fixed point for the Poincaré map S . Moreover, the planar fixed point E_3 is globally asymptotically stable in $\text{Int}\mathbb{R}_+^3$.

Theorem 4.7. (Two species Extinction) *Suppose that system (1.2) satisfies*

$$(i) \ h_i > 0 \ (i = 1, 2, 3), \ \det \bar{A} < 0;$$

$$(ii) \ \frac{h_1}{a_{12}} < \frac{h_2}{a_{22}}; \ \frac{h_3}{a_{32}} < \frac{h_2}{a_{22}}; \ h_2 > \frac{a_{21}}{a_{11}}h_1 + \frac{a_{23}}{a_{33}}h_3;$$

then there is no positive fixed point for the Poincaré map S . Moreover, the axial fixed point R_2 is globally asymptotically stable in $\text{Int}\mathbb{R}_+^3$.

Remark 4.1. By using our analytical approach in Theorems 4.5–4.7, we also take different sufficient conditions to obtain the global dynamics of system (1.2).

Next, we will investigate the global extinction for system (1.2). For simplicity, we introduce the following notations

$$\tilde{A} = \begin{pmatrix} a_{11} & a_{12} & 0 \\ a_{21} & a_{22} & a_{23} \\ 0 & a_{32} & a_{33} \end{pmatrix}, \quad \tilde{B} = \begin{pmatrix} h_1 \\ h_2 \\ h_2 \end{pmatrix} \quad \text{and} \quad y = \begin{pmatrix} y_1 \\ y_2 \\ y_2 \end{pmatrix}.$$

Lemma 4.8. If S has a positive fixed point, say P , then $y := \int_0^{\varphi\omega} Q_t(LP)dt$ is a positive solution of the linear algebraic system $\tilde{A}y = \tilde{B}$. In other words, if the linear system $\tilde{A}y = \tilde{B}$ has no positive solution, then S has no positive fixed point in $\text{Int}\mathbb{R}_+^3$.

Proof. We write the positive fixed point $P := (\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$ as Lemma 3.8. Let $Q_t(LP) = u(t, LP) := (\tilde{u}_1(t), \tilde{u}_2(t), \tilde{u}_3(t))$, then $(\tilde{u}_1(t), \tilde{u}_2(t), \tilde{u}_3(t))$ satisfies the following equations

$$\begin{cases} \frac{\tilde{u}_1'(t)}{\tilde{u}_1(t)} = r_1 - a_{11}\tilde{u}_1(t) - a_{12}\tilde{u}_2(t), \\ \frac{\tilde{u}_2'(t)}{\tilde{u}_2(t)} = r_2 - a_{22}\tilde{u}_2(t) - a_{21}\tilde{u}_1(t) - a_{23}\tilde{u}_3(t), \\ \frac{\tilde{u}_3'(t)}{\tilde{u}_3(t)} = r_3 - a_{33}\tilde{u}_3(t) - a_{32}\tilde{u}_2(t). \end{cases} \quad (4.1)$$

Integrating the above equations of (4.1) for t from 0 to $\varphi\omega$, it follows that

$$\begin{cases} a_{11} \int_0^{\varphi\omega} \tilde{u}_1(t)dt + a_{12} \int_0^{\varphi\omega} \tilde{u}_2(t)dt = (r_1\varphi - \lambda_1(1 - \varphi))\omega, \\ a_{21} \int_0^{\varphi\omega} \tilde{u}_1(t)dt + a_{22} \int_0^{\varphi\omega} \tilde{u}_2(t)dt + a_{23} \int_0^{\varphi\omega} \tilde{u}_3(t)dt = (r_2\varphi - \lambda_2(1 - \varphi))\omega, \\ a_{32} \int_0^{\varphi\omega} \tilde{u}_2(t)dt + a_{33} \int_0^{\varphi\omega} \tilde{u}_3(t)dt = (r_3\varphi - \lambda_3(1 - \varphi))\omega. \end{cases}$$

Note that $y_i = \int_0^{\varphi\omega} \tilde{u}_i(t)dt$ and $h_i = (r_i\varphi - \lambda_i(1 - \varphi))\omega$, $i = 1, 2, 3$, we have

$$\begin{cases} a_{11}y_1 + a_{12}y_2 = h_1, \\ a_{21}y_1 + a_{22}y_2 + a_{23}y_3 = h_2, \\ a_{32}y_2 + a_{33}y_3 = h_3. \end{cases} \quad (4.2)$$

From the above equations, it is clear that $\tilde{A}y = \tilde{B}$ with $y_i > 0$ ($i = 1, 2, 3$). Therefore, the conclusion is immediate. The proof is completed. \square

Theorem 4.9. (Global Extinction) Suppose that $h_i < 0$ ($i = 1, 2, 3$). Then the trivial fixed point O is globally asymptotically stable in \mathbb{R}_+^3 .

Proof. Since $h_i < 0$ ($i = 1, 2, 3$), it follows that S has no interior fixed points on the coordinate axes of \mathbb{R}_+^3 due to Lemma 2.1(i). By appealing to Lemma 2.4 in [11], there are also no interior fixed points on the coordinate plane of \mathbb{R}_+^3 . Lemma 4.8 implies that there is no positive fixed point in $\text{Int}\mathbb{R}_+^3$ for S . Therefore, O is a unique fixed point for S in \mathbb{R}_+^3 . By the equations of system (1.4), one has

$$\frac{dx_i(t)}{dt} \leq x_i(t)(r_i(t) - a_{ii}(t)x_i(t)), \quad i = 1, 2, 3.$$

Since $h_i < 0$, by differential inequality theorem and Lemma 2.1(i), it follows that

$$\lim_{t \rightarrow +\infty} x_i(t, x_0) = 0, \quad \forall x_0 \in \mathbb{R}_+^3, \quad i = 1, 2, 3.$$

Note that Lemma 3.1(i), O is an asymptotically stable fixed point of S , and hence, O is globally asymptotically stable in \mathbb{R}_+^3 . We have completed the proof. \square

5. Numerical simulation

In this section, we provide some numerical examples to illustrate our analytic results.

Example 1. (Three species Coexistence) Taking parameter values $\omega = 10$, $\phi = 0.5$, $\lambda_1 = 0.1$, $\lambda_2 = 0.1$, $\lambda_3 = 0.1$, $r_1 = 0.3$, $r_2 = 0.3$, $r_3 = 0.3$, $a_{11} = 0.4$, $a_{12} = 0.2$, $a_{21} = 0.1$, $a_{22} = 0.5$, $a_{23} = 0.1$, $a_{32} = 0.2$, $a_{33} = 0.5$, and initial values $x^0 = (0.8, 0.9, 0.8)$, system (1.2) satisfies the conditions of Theorem 4.4. The numerical simulations for the solution of system (1.2) and the orbit of the Poincaré map S are shown in Figure 1, which imply that three species will be coexistence.




Figure 1. Three species will be coexistence.




Figure 2. Species 1 and Species 3 will be coexistence.

Example 2. (Two species coexistence) Taking parameter values $\omega = 10$, $\phi = 0.5$, $\lambda_1 = 0.1$, $\lambda_2 = 0.1$, $\lambda_3 = 0.1$, $r_1 = 0.3$, $r_2 = 0.3$, $r_3 = 0.3$, $a_{11} = 0.2$, $a_{12} = 0.2$, $a_{21} = 0.3$, $a_{22} = 1$, $a_{23} = 0.22$, $a_{32} = 0.1$, $a_{33} = 0.2$, and initial values $x^0 = (0.8, 0.9, 0.8)$, system (1.2) satisfies the conditions of Theorem 4.5.

The numerical simulations for the solution of system (1.2) and the orbit of the Poincaré map S are shown in Figure 2, which imply that Species 1 and Species 3 will be coexistence, and Species 2 will go to extinction.

Example 3. (Two species coexistence) Taking parameter values $\omega = 10$, $\phi = 0.5$, $\lambda_1 = 0.1$, $\lambda_2 = 0.1$, $\lambda_3 = 0.1$, $r_1 = 0.3$, $r_2 = 0.3$, $r_3 = 0.3$, $a_{11} = 0.6$, $a_{12} = 0.1$, $a_{21} = 0.1$, $a_{22} = 0.2$, $a_{23} = 0.2$, $a_{32} = 0.3$, $a_{33} = 0.5$, and initial values $x^0 = (0.8, 0.9, 0.8)$, system (1.2) satisfies the conditions of Theorem 4.6. The numerical simulations for the solution of system (1.2) and the orbit of the Poincaré map S are shown in Figure 3, which imply that Species 1 and 2 will be coexistence, and Species 3 will go to extinction.




Figure 3. Species 1 and Species 2 will be coexistence.




Figure 4. Species 2 will win the competition.

Example 4. (Two species extinction) Taking parameter values $\omega = 10$, $\phi = 0.5$, $\lambda_1 = 0.1$, $\lambda_2 = 0.1$, $\lambda_3 = 0.1$, $r_1 = 0.3$, $r_2 = 0.3$, $r_3 = 0.3$, $a_{11} = 0.6$, $a_{12} = 0.3$, $a_{21} = 0.1$, $a_{22} = 0.23$, $a_{23} = 0.2$, $a_{32} = 0.25$,

$a_{33} = 0.5$, and initial values $x^0 = (0.8, 0.9, 0.8)$, system (1.2) satisfies the conditions of Theorem 4.7. The numerical simulations for the solution of system (1.2) and the orbit of the Poincaré map S are shown in Figure 4, which imply that Species 2 will win the competition and Species 1 and Species 3 will go to extinction.

Example 5. (Global extinction) Taking parameter values $\omega = 10$, $\phi = 0.5$, $\lambda_1 = 0.3$, $\lambda_2 = 0.3$, $\lambda_3 = 0.3$, $r_1 = 0.1$, $r_2 = 0.1$, $r_3 = 0.1$, $a_{11} = 0.6$, $a_{12} = 0.1$, $a_{21} = 0.1$, $a_{22} = 0.2$, $a_{23} = 0.2$, $a_{32} = 0.3$, $a_{33} = 0.5$, and initial values $x^0 = (0.8, 0.9, 0.8)$, system (1.2) satisfies the conditions of Theorem 4.9. The numerical simulations for the solution of system (1.2) and the orbit of the Poincaré map S are shown in Figure 5, which imply that three species will go to extinction.




Figure 5. Three species will be extinction.

6. Discussion

In this paper, we focus on a tridiagonal three-species competition model with time ω -periodic coefficients (called Seasonal Succession). By the stability analysis of equilibria, we estimate the Floquet multipliers of all nonnegative periodic solutions of system (1.2), and get the local dynamics of these periodic solutions. By using the Brouwer degree theory, we present an index result of the fixed points for the Poincaré map S . Based on this, we can verify the existence and uniqueness of the positive fixed point under appropriate conditions. Sufficient conditions of the global stability for coexistence and extinction of system (1.2) are provided via the local dynamics of all nonnegative fixed points. More precisely, three species will be existence under the assumptions of Theorem 4.4; Species 1 and 3 will be coexistence and Species 2 will go to extinction in the competition under the assumptions of Theorem 4.5; Species 1 and 2 will be coexistence and Species 3 will go to extinction in the competition under the assumptions of Theorem 4.6; Species 2 will win the competition and Species 1 and 3 will go to extinction under the assumptions of Theorem 4.7; three species will go to extinction under the assumptions of Theorem 4.9. From above analytic results, it is not difficult to see that the introduction of seasonal succession may lead to species' extinction.

On the other hand, there is no explicit expression of the Poincaré map S for the time-periodic differential equations, even for the simplest form as system (1.2). This makes the researches on the

dynamics the Poincaré map of the time-periodic Kolmogorov competitive systems become much more difficult and complicated. In future work, we will try to give a complete classification for the dynamics of system (1.2) and explore the influence of parameter values φ and $\lambda_i, i = 1, 2, 3$ related to the seasonal succession on the dynamics of the system (1.2).

Acknowledgments

We are very grateful to the anonymous referees for their valuable comments and suggestions, which led to an improvement of our original manuscript. This work was supported by the National Natural Science Foundation of China (No.11871231) and Natural Science Foundation Fund of Fujian Province (No.2022J01305).

Conflict of interest

The authors declare there is no conflict of interest.

References

1. H. L. Smith, *Monotone Dynamical Systems. An Introduction to the Theory of Competitive and Cooperative Systems*, American Mathematical Society, 1995. <https://doi.org/10.1090/surv/041>
2. J. D. Murray, *Mathematical Biology I: An Introduction*. 3rd edition, Springer-Verlag, Berlin, 2002.
3. M. W. Hirsch, H. L. Smith, Monotone dynamical systems, in *Handbook of Differential Equations: Ordinary Differential Equations*, Elsevier, Amsterdam, 2005. [https://doi.org/10.1016/S1874-5725\(05\)80006-9](https://doi.org/10.1016/S1874-5725(05)80006-9)
4. X. Q. Zhao, *Dynamical systems in population biology*. 2nd edition, Springer, New York, 2017. <https://doi.org/10.1007/978-3-319-56433-3>
5. J. Smillie, Competitive and cooperative tridiagonal systems of differential equations, *SIAM J. Math. Anal.*, **15** (1984), 530–534. <https://doi.org/10.1137/0515040>
6. H. L. Smith, Periodic tridiagonal competitive and cooperative systems of differential equations, *SIAM J. Math. Anal.*, **22**(1991), 1102–1109. <https://doi.org/10.1137/0522071>
7. M. Gyllenberg, Y. Wang, Periodic tridiagonal systems modeling competitive-cooperative ecological interactions, *Discrete Contin. Dyn. Syst. B*, **5** (2005), 289–298. <https://doi.org/10.3934/dcdsb.2005.5.289>
8. C. Fang, M. Gyllenberg, Y. Wang, Floquet bundles for tridiagonal competitive-cooperative systems and the dynamics of time-recurrent systems, *SIAM J. Math. Anal.*, **45** (2013), 2477–2498. <https://doi.org/10.1137/120878021>

9. C. A. Klausmeier, Successional state dynamics: A novel approach to modeling nonequilibrium foodweb dynamics, *J. Theor. Biol.*, **262** (2010), 584–595. <https://doi.org/10.1016/j.jtbi.2009.10.018>
10. C. F. Steiner, A. S. Schwaderer, V. Huber, C. A. Klausmeier, E. Litchman, Periodically forced food chain dynamics: model predictions and experimental validation, *Ecology*, **90** (2009), 3099–3107. <https://doi.org/10.1890/08-2377.1>
11. S. B. Hsu, X. Q. Zhao, A Lotka-Volterra competition model with seasonal succession, *J. Math. Biol.*, **64** (2012), 109–130. <https://doi.org/10.1007/s00285-011-0408-6>
12. L. Niu, Y. Wang, X. Xie, Carrying simplex in the Lotka-Volterra competition model with seasonal succession with applications, *Discrete Contin. Dyn. Syst. B*, **26** (2021), 2161–2172. <https://doi.org/10.3934/dcdsb.2021014>
13. X. Xie, L. Niu, Global stability in a three-species Lotka-Volterra cooperation model with seasonal succession, *Math. Meth. Appl. Sci.*, **44** (2021), 14807–14822. <https://doi.org/10.1002/mma.7744>
14. Y. Zhang, X. Q. Zhao, Bistable travelling waves for reaction and diffusion model with seasonal succession, *Nonlinearity*, **26** (2013), 691–709. <https://doi.org/10.1088/0951-7715/26/3/691>
15. D. Xiao, Dynamics and bifurcations on a class of population model with seasonal constant-yield harvesting, *Discrete Contin. Dyn. Syst. B*, **21** (2016), 699–719. <https://doi.org/10.3934/dcdsb.2016.21.699>
16. P. G. Barrientos, J. A. Rodriguez, A. Ruiz-Herrera, Chaotic dynamics in the seasonally forced SIR epidemic model, *J. Math. Biol.*, **75** (2017), 1655–1668. <https://doi.org/10.1007/s00285-017-1130-9>
17. M. Han, X. Hou, L. Sheng, C. Wang, Theory of rotated equations and applications to a population model, *Discrete Contin. Dyn. Syst.*, **38** (2018), 2171–2185. <https://doi.org/10.3934/dcds.2018089>
18. Y. Tang, D. Xiao, W. Zhang, D. Zhu, Dynamics of epidemic models with asymptomatic infection and seasonal succession, *Math. Biosci. Eng.*, **14** (2017), 1407–1424. <https://doi.org/10.3934/mbe.2017073>
19. H. Amann, Fixed point equations and nonlinear eigenvalue problems in Ordered Banach Spaces, *SIAM Rev.*, **18** (1976), 620–709. <https://doi.org/10.1137/1018114>
20. X. Liang, J. Jiang, On the finite-dimensional dynamical systems with limited competition, *Trans. Am. Math. Soc.*, **354** (2002), 3535–3554. <https://doi.org/10.1090/S0002-9947-02-03032-5>
21. E. C. Balreira, S. Elaydi, R. Luís, Global stability of higher dimensional monotone maps, *J. Differ. Equation Appl.*, **23** (2017), 2037–2071. <https://doi.org/10.1080/10236198.2017.1388375>
22. A. Ruiz-Herrera, Exclusion and dominance in discrete population models via the carrying simplex, *J. Differ. Equation Appl.*, **19** (2013), 96–113. <https://doi.org/10.1080/10236198.2011.628663>
23. A. Ruiz-Herrera, Topological criteria of global attraction with applications in population dynamics, *Nonlinearity*, **25** (2012), 2823–2841. <https://doi.org/10.1088/0951-7715/25/10/2823>
24. H. L. Smith, Competing subcommunities of mutualists and a generalized Kamke theorem, *SIAM J. Math. Anal.*, **46** (1986), 856–874. <https://doi.org/10.1137/0146052>

-
25. H. L. Smith, P. Waltman, *The Theory of the Chemostat*, Cambridge University Press, 1995. <https://doi.org/10.1017/CBO9780511530043>
 26. Y. Wang, J. Jiang, Uniqueness and attractivity of the carrying simplex for discrete-time competitive dynamical systems, *J. Differ. Equations*, **186** (2002), 611–632. [https://doi.org/10.1016/S0022-0396\(02\)00025-6](https://doi.org/10.1016/S0022-0396(02)00025-6)



AIMS Press

©2023 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)